

SYSTEMATIC SEARCH FOR SINGULARITIES

SYSTEMATIC SEARCH FOR SINGULARITIES IN 3D EULER
FLOWS BASED ON THE VOIGT REGULARIZATION

BY

NOAH BENSLER, B.Sc.

A THESIS

SUBMITTED TO THE SCHOOL OF COMPUTATIONAL SCIENCE & ENGINEERING

AND THE SCHOOL OF GRADUATE STUDIES

OF MCMASTER UNIVERSITY

IN PARTIAL FULFILMENT OF THE REQUIREMENTS

FOR THE DEGREE OF

MASTERS OF COMPUTATIONAL SCIENCE & ENGINEERING

© Copyright by Noah Bensler, September 2026

All Rights Reserved

Masters of Computational Science & Engineering (2026) McMaster University
(School of Computational Science & Engineering) Hamilton, Ontario, Canada

TITLE: SYSTEMATIC SEARCH FOR SINGULARITIES IN 3D
 EULER FLOWS BASED ON THE VOIGT REGULAR-
 IZATION

AUTHOR: Noah Bensler
 B.Sc. (Computer Science & Astrophysics),
 University of Calgary, Calgary, Canada

SUPERVISOR: Dr. Bartosz Protas

NUMBER OF PAGES: [xvi](#), [68](#)

Lay Abstract

A lay abstract of not more 150 words must be included explaining the key goals and contributions of the thesis in lay terms that is accessible to the general public.

The Navier-Stokes equations model the behaviour of viscous incompressible fluids. The question of whether or not the Navier-Stokes equations, or the inviscid Euler equations, are a valid description of fluid mechanics remains unanswered. A singularity is the occurrence of an infinite derivative of the velocity at a single point in the fluid flow. If the equations form a singularity from smooth initial conditions, then they are not valid descriptions of fluid mechanics. The Euler-Voigt equations are an inviscid regularization of the Euler equations which are known to not form singularities from smooth initial conditions. However, by decreasing the regularization parameter we approximate Euler flows. We examine the Euler-Voigt equations with a PDE-constrained optimization problem to identify smooth initial conditions that produce 'extreme' behaviour in numerical simulations. ## We present numerical evidence that CONCLUSION ##

Abstract

Abstract here (no more than 300 words). An abstract of not more than 300 words must be included and indicates the major emphasis of the thesis, new discoveries, and its contribution to knowledge. The style of abstract varies from discipline to discipline but the student should follow an appropriate style. This page must be numbered 'iv'.

The Navier-Stokes equations are a system of partial differential equations used to model the behaviour of viscous incompressible fluids. The question of whether or not unique classical solutions of the Navier-Stokes equations, and inviscid Euler equations, exist for all time, given smooth initial conditions, remains unanswered for the three-dimensional problem.

A singularity is the occurrence of an infinite derivative of the velocity at a single point in the fluid flow. The formation of a singularity from smooth initial conditions would invalidate the equations as accurate descriptions of fluid mechanics. The Euler-Voigt equations are an inviscid regularization of the Euler equations which are known to be globally well-posed from smooth initial conditions. However, by decreasing the regularization parameter the solutions of the Euler-Voigt equations converge

to those of the Euler equations. We examine the Euler-Voigt equations with a PDE-constrained optimization problem to identify smooth initial conditions that produce 'extreme' behaviour in numerical simulations. ## We present numerical evidence that CONCLUSION ##

Your Dedication
Optional second line

Acknowledgements

Acknowledgements go here. An expression of thanks for assistance given by the Supervisor of research and by others should be either set forth on a separate page or incorporated into the Preface (if there is one). These and all subsequent preliminary pages listed under (f), (g) and (h) must be numbered in lower case Roman numerals, i.e., ‘iv’, ‘v’, ‘vi’ etc.

Dr. Bartosz Protas, Dr. Xinyu Zhao, Dr. Jörn Davidsen, members of committee,
Parents

I would like to thank Dr. Bartosz Protas for his support and guidance on my research project. I also want to thank Dr. Jörn Davidsen of the Complexity Science Group for supervising my undergraduate research and inspiring me to pursue computational physics.

Contents

Lay Abstract	iii
Abstract	iv
Acknowledgements	vii
Notation, and Abbreviations	xiv
Declaration of Academic Achievement	xv
1 Introduction	1
1.1 Singularity Formation in Models of 3D Fluid Flows	1
1.2 3D Euler-Voigt Equations	4
1.3 Outline of the Thesis	5
2 Properties of the Euler-Voigt Equations	7
2.1 Finite Time Blow-up Criterion	7
2.2 Initial Conditions	11
2.3 Scaling Symmetries of Euler and Euler-Voigt Systems	13
2.4 Initial Guesses	16

3	The Optimization Problem	17
3.1	Formulation of the Problem	17
3.2	Gradient of Objective Function	18
3.3	Riemannian Conjugate Gradient Method	21
4	Numerical Methods	26
4.1	Spatial Discretization & Spectral Methods	26
4.2	Time Stepping	26
4.3	Determination of Gevrey class size	31
4.4	Spectrum Validation	32
4.5	Kappa Test Validation	33
5	Results	36
5.1	Singularity Detection	36
5.2	Initial Guess Results	38
6	Conclusion	43
A	Function Spaces	44
B	Proof of Relation (2.25)	46
C	Rescaling the Euler equations	47
D	Rescaling the Euler-Voigt equations	49
E	Evaluation of $\ \boldsymbol{\eta}\ _{\dot{H}^1}$ for the General Taylor Green Vortex	51
F	Evaluation of $\ \nabla \times \boldsymbol{\eta}\ _{L^\infty}$ for the General Taylor Green Vortex	54

G	Linearization of the Euler-Voigt System	56
H	Derivation of the Adjoint System	58
I	Derivation of the Gâteaux Directional Differential of the Objective Functional	63

List of Figures

2.1	Maximum of vorticity $\ \omega(\cdot, t)\ _\infty$. Results from runs performed at different resolutions are displayed together: 128^3 (orange plus), 256^3 (cyan star), 512^3 (brown triangles), 1024^3 (blue squares).	15
2.2	Isosurfaces of the scaled TGV vorticity. (a) magnitude of ω , vorticity components (b) ω_1 , (c) ω_2 , and (d) ω_3	16
5.1	Isosurfaces of the vorticity at $T=5$ from the TGV. (a) magnitude of ω , vorticity components (b) ω_1 , (c) ω_2 , and (d) ω_3	38
5.2	Isosurfaces of the vorticity at $T=5$ from the TGV. (a) magnitude of ω , vorticity components (b) ω_1 , (c) ω_2 , and (d) ω_3	39
5.3	Nonoptimized TGV with $T = 3$. Log-log plot of $\max_{t \in [0, T']} \ \nabla \mathbf{u}_\alpha(t)\ _{L^2}$ vs α at $T' = 0$ (blue) to $T' = 3$ (red), Steepest slope $p = -0.225$ (black), slope of $p = -1$ (dotted black)	39
5.4	Nonoptimized TGV with $T = 3$. Plot of the slope of $(\max_{t \in [0, T']} \ \nabla \mathbf{u}_\alpha(t)\ _{L^2}$ vs $\alpha)$ vs time at each α	40
5.5	Nonoptimized TGV with $T = 3$. Log-log plot of $\max_{t \in [0, T']} \ \nabla \mathbf{u}_\alpha(t)\ _{L^2}$ vs α at $T' = 0$ (blue) to $T' = 5$ (red), Steepest slope $p = -0.417$ (black), slope of $p = -1$ (dotted black)	41

5.6	Nonoptimized TGV with $T = 5$. Plot of the slope of $(\max_{t \in [0, T]} \ \nabla \mathbf{u}_\alpha(t)\ _{L^2}$ vs α) vs time at each α	41
-----	---	----

List of Tables

4.1	Relative error of Φ_{dt} for each step size to Φ_c . Resolution $N = 128^3$ and time window $T = 5$	28
4.2	Relative error of Φ_{dt} for each step size to Φ_c . Resolution $N = 256^3$ and time window $T = 5$	28
4.3	Relative error of Φ_{dt} for each step size to Φ_c . Resolution $N = 512^3$ and time window $T = 5$	28
4.4	Relative error of Φ_{dt} for each step size to Φ_c . Resolution $N = 1024^3$ and time window $T = 5$	29
4.5	Maximum relative error of $E_\alpha(t)$ to $E_\alpha(0)$ for each step size. Resolution $N = 128^3$ and time window $T = 5$	30
4.6	Maximum relative error of $E_\alpha(t)$ to $E_\alpha(0)$ for each step size. Resolution $N = 256^3$ and time window $T = 5$	30
4.7	Maximum relative error of $E_\alpha(t)$ to $E_\alpha(0)$ for each step size. Resolution $N = 512^3$ and time window $T = 5$	30
4.8	Maximum relative error of $E_\alpha(t)$ to $E_\alpha(0)$ for each step size. Resolution $N = 1024^3$ and time window $T = 5$	31
4.9	Selection of Δt for each resolution and α	31

Notation, and Abbreviations

Notation

x	Plain-text to denote scalar x
$\boldsymbol{x}$	Bold-text to denote vector x
$\boldsymbol{x}^\top$	Transpose of vector x
I	Identity Operator
∇	Gradient Operator
Δ	Laplacian Operator
$:=$	Equal to by definition

Abbreviations

PDEs	Partial Differential Equations
3D	three-dimensional or three dimensions
TGV	Taylor Green Vortex

Declaration of Academic Achievement

Dr. Zhao developed most of the mathematics for the subspace V (2.26), along with Theorem 2.1.3 and its proof. Dr. Zhao also developed the mathematics for much of the Euler-Voigt optimization problem described in Chapter 3, including the objective functional (3.32), the linearization of the Euler-Voigt equations (3.37), the adjoint system (3.38), and the gradient of the objective functional (3.44). The Riemannian conjugate gradient method for the Euler-Voigt optimization problem in Section 3.3 was developed by Dr. Zhao and Dr. Protas. This includes the tangent space (3.45), the projection operator (3.48), the search direction (3.52), and step size (3.55).

The retraction operator (3.49), and the vector transport operation (3.50) were originally developed by Dr. Zhao and Dr. Protas but I modified these formulations to suit the present problem. The formulation of the manifold $\mathcal{M}_C$ (2.28) was based on work by Dr. Zhao, but the calculation of the constraint parameter C was my work. This included the calculations of the $\dot{H}^1$ seminorm (2.30) and vorticity (2.31) of the general Taylor Green Vortex, which are included in Appendices E and F. The results in Appendices B, G, H, and I are my re-derivations of Dr. Zhao's original results. The derivation in Appendices C and D are my re-derivations of the well known Euler and Euler-Voigt scaling properties.

For the numerical models, Dr. Zhao created the original code base which I modified to suit this project. The determination of the numerical model constants, the time and spatial resolutions, regularization parameter α , and Gevrey parameter σ were also my work. These values were validated through analysis of the stability and accuracy of the model, conservation of the α -energy, comparison of the effect of different values of σ , examination of the solutions' Fourier spectrum, and the kappa test, all of which were my work and are described in Chapter 4. I also validated the results of the numerical models through comparison with the relevant findings of [3] and [17]. I performed all of the computations to obtain the final numerical results. The findings and conclusions in Chapters 5 and 6 are my work. The Python code used to perform the data analysis and produce the graphs is also my own work.

1 Introduction

1.1 Singularity Formation in Models of 3D Fluid Flows

The three-dimensional (3D) Navier-Stokes equations are a set of partial differential equations (PDEs) that describe the motion of homogeneous, viscous, incompressible fluids. The Navier-Stokes equations on the domain $\Omega \subseteq \mathbb{R}^3$ are defined as

$$\partial_t \mathbf{u} + \mathbf{u} \cdot \nabla \mathbf{u} + \nabla p - \nu \Delta \mathbf{u} = 0 \quad \text{in } \Omega \times (0, T], \quad (1.1a)$$

$$\nabla \cdot \mathbf{u} = 0 \quad \text{in } \Omega \times [0, T], \quad (1.1b)$$

$$\mathbf{u}(x, 0) = \boldsymbol{\eta} \quad \text{in } \Omega \quad (1.1c)$$

where $\mathbf{u} = [u_1, u_2, u_3]^\top$ is the fluid velocity field, p is the scalar pressure, $\nu > 0$ is the coefficient of kinematic viscosity, and T is the length of the time interval. The field $\boldsymbol{\eta}$ is the initial condition assumed to be divergence-free. Finally, the gradient operator $\nabla = [\partial_1, \partial_2, \partial_3]^\top$ is used to define the gradient of the pressure, the tensor $[\nabla \mathbf{u}]_{ij} = \partial_j u_i$ for $i, j = 1, 2, 3$, the divergence of the velocity field, and the Laplacian operator $\Delta \mathbf{u} = \nabla \cdot (\nabla \mathbf{u})$. Because of their generality, the Navier-Stokes equations are utilized to model the behaviour of systems in fields as disparate as aero/hydrodynamics, weather prediction, biophysics, and astrophysics [8, 10]. Despite the numerous fields

that make use of the 3D Navier-Stokes equations, and extensive research examining their properties, it is only known that weak Leray-Hopf solutions exist globally in time [19]. However, these solutions may contain singularities and their uniqueness remains unproven in most cases. It is also yet to be proven whether or not smooth classical solutions, defined in the pointwise sense, exist globally in time [25]. In fact, this is the basis of one of the seven **Millennium Problems** posed by the Clay Mathematics Institute which offers a substantial monetary prize for the confirmation of the existence, or non-existence, of a classical solution corresponding to smooth initial conditions [11]. The space of smooth functions and all other function spaces utilized in this analysis are **defined** in Appendix A. It is common for scientific research that focuses the fundamental properties of the Navier-Stokes equations to utilize either the unbounded domain $\Omega := \mathbb{R}^3$, the axisymmetric domain, or the periodic 3D torus $\Omega := \mathbb{T}_L^3$ with domain size $L > 0$ [25]. For this analysis, we use the 3D torus of unit size ($L = 1$) with periodic boundary conditions.

A central question in fluid mechanics is whether or not the Navier-Stokes equations are globally well-posed or if singularities can form from smooth initial conditions. A singularity occurs when the derivative of the velocity field becomes infinite at a single point [22]. The formation of a singularity in a fluid flow, from smooth initial conditions, would invalidate the Navier-Stokes equations as an accurate description of fluid flows [25]. The question about singularity formation from smooth initial conditions also remains open for the Euler equations [13]. The Euler equations are the inviscid

form of the Navier-Stokes equations, obtained by setting $\nu = 0$

$$\partial_t \mathbf{u} + \mathbf{u} \cdot \nabla \mathbf{u} + \nabla p = 0 \quad \text{in } \Omega \times (0, T], \quad (1.2a)$$

$$\nabla \cdot \mathbf{u} = 0 \quad \text{in } \Omega \times [0, T], \quad (1.2b)$$

$$\mathbf{u}(x, 0) = \boldsymbol{\eta} \quad \text{in } \Omega. \quad (1.2c)$$

The Euler equations are often studied because they are simpler to analyse than the Navier-Stokes equations due to the absence of the viscous dissipation. The local existence of classical solutions to the Euler equations was proven in [16] for solutions defined in Sobolev spaces.

Theorem 1.1.1 ([16]). *If $\boldsymbol{\eta} \in H^m(\mathbb{T}^3)$ for some $m > 5/2$ and satisfies $\nabla \cdot \boldsymbol{\eta} = 0$, then there exists a time $T = T(\|\boldsymbol{\eta}\|_{H^m}) > 0$ such that (1.2) has a unique solution $\mathbf{u}(\cdot; \boldsymbol{\eta}) \in C([0, T]; H^m) \cap C^1([0, T]; H^{m-1})$.*

If a singularity is to form in the solutions to the Euler equations, it must occur after the minimum local existence time defined in Theorem 1.1.1. There are many different conditional regularity results for determining if a singularity has formed, the best known of which is the Beale-Kato-Majda (BKM) criterion [1] which asserts

Theorem 1.1.2 ([1]). *A smooth solution $\mathbf{u}$ of the Euler system develops a singularity at $t = t_0$ if and only if*

$$\lim_{t \rightarrow t_0} \int_0^t \|\boldsymbol{\omega}\|_{L^\infty} d\tau = \infty, \quad \boldsymbol{\omega} = \nabla \times \mathbf{u}, \quad (1.3)$$

where $\|\cdot\|_{L^\infty}$ is the standard Lebesgue L^∞ norm.

A key difference between the Navier-Stokes and Euler equations is that there is more numerical evidence that indicates singularity formation in Euler is possible from smooth initial conditions. For example, [2] and [3] presented numerical evidence for singularity formation in the Euler equations on a 3D torus ($L = 2\pi$). Hou et al. produced several papers which provide numerical evidence for singularity formation in the 3D axisymmetric Euler equations [21][15][14] and which proved singularity formation in the 3D axisymmetric Euler equations with rigid boundaries and periodic boundary conditions in the axial direction [6][7]. We make special note of the optimization-based approach of [27], which utilized a Riemannian conjugate gradient method to identify 'extreme' initial conditions, and **reports** numerical results consistent with singularity formation in Euler flows on the 3D unit torus. Finally, [17] which provides numerical evidence for singularity formation in the Euler equations based on the Voigt regularization. These final two papers are of special significance for us because they form the primary basis for the motivation and methodology of this analysis.

1.2 3D Euler-Voigt Equations

The Euler-Voigt equations are an inviscid regularization of the Euler equations (1.2) which have been shown to be globally well-posed and so do not form singularities from smooth initial conditions [5]. The Euler-Voigt equations are defined as

$$-\alpha^2 \partial_t \Delta \mathbf{u}_\alpha + \partial_t \mathbf{u}_\alpha + \mathbf{u}_\alpha \cdot \nabla \mathbf{u}_\alpha + \nabla p_\alpha = 0 \quad \text{in } \Omega \times (0, T], \quad (1.4a)$$

$$\nabla \cdot \mathbf{u}_\alpha = 0 \quad \text{in } \Omega \times (0, T], \quad (1.4b)$$

$$\mathbf{u}_\alpha(x, 0) = \boldsymbol{\eta}_\alpha \quad \text{in } \Omega, \quad (1.4c)$$

where the regularization is accomplished with the addition of the term $-\alpha^2 \partial_t \Delta \mathbf{u}_\alpha$ in which $\alpha > 0$ is the regularization parameter with units of length. This term stabilizes the system without dissipation, which would remove energy from the system [17]. The regularization term acts as a spectral filter and suppresses the development of small scale structures with length-scale smaller than α [23]. We introduce the subscript α to distinguish between solutions of (1.2) and (1.4) i.e., $\mathbf{u}$ and $\mathbf{u}_\alpha$.

For (1.2) the total kinetic energy is conserved so long as the solution remains well-posed in the classical sense

$$E_k(t) := \frac{1}{2} \|\mathbf{u}\|_{L^2}^2 = E_k(0), \quad (1.5)$$

where the norm $\|\cdot\|_{L^2}$ here is the standard Lebesgue L^2 norm. For (1.4) the conserved quantity is known to be the α -energy. The α -energy is defined as

$$E_\alpha(t) := \|\mathbf{u}_\alpha(t)\|_{L^2}^2 + \alpha^2 \|\nabla \mathbf{u}_\alpha(t)\|_{L^2}^2 = E_\alpha(0). \quad (1.6)$$

1.3 Outline of the Thesis

The goal of this analysis is to search for possible singularities in Euler flows corresponding to smooth initial conditions. This search will be performed based on a PDE-constrained optimization problem in which we will identify initial condition that produce the most 'extreme' behaviour in Euler-Voigt flows. In Chapter 2 we present the properties of the Euler-Voigt equations, including the blow-up criteria, scaling property, and manifold over which we search for initial conditions. In Chapter 3 we

present the formulation of the optimization problem. This includes the description of our objective functional, the gradient calculation, and the Riemannian conjugate gradient method used to compute our search direction and iterative relation. In Chapter 4 we describe the numerical methods used to perform the calculations, including approximation in space and time, Gevrey parameters, and the regularization parameter. In addition, we validate the computations using the Fourier spectrum, kappa test, and conservation of α -energy. In Section 5 we present our results. We begin with the solutions of the Euler-Voigt equations corresponding to the Taylor Green Vortex without optimization as $\alpha \rightarrow 0$. These results indicate that singularity formation does not occur within the time window. We then present our optimization results as $\alpha \rightarrow 0$, which indicate that RESULTS RESULTS RESULTS. Finally, in Chapter 6 we summarize our findings and present our conclusion.

2 Properties of the Euler-Voigt Equations

In this chapter we prove ~~Theorem 2.1.3~~ which provides the criterion used to determine if a singularity forms in Euler flows from smooth initial conditions. Additionally, we construct ~~the manifold (2.28)~~ in which we search for optimal smooth initial conditions. Finally, we examine the structure of the initial guesses including the Taylor Green Vortex, Together, these form the basis for our optimization problem.

2.1 Finite Time Blow-up Criterion

In order to begin our search for finite-time singularity formation, we must first describe the criteria used to detect singularity formation in Euler flows. In [18] a new criterion for identifying singularity formation in Euler flows was developed which utilizes the Euler-Voigt equations. This criterion relies on the convergence of the solutions of (1.4) to the solutions of (1.2) as $\alpha \rightarrow 0$.

Theorem 2.1.1 ([18]). *Given initial data $\boldsymbol{\eta}, \boldsymbol{\eta}_\alpha \in H^s(\mathbb{T}^3)$ for some $s \geq 3$, let $\boldsymbol{u}, \boldsymbol{u}_\alpha \in C([0, T], H^s(\Omega)) \cap C^1([0, T], H^{s-1}(\Omega))$ be the corresponding solutions to (1.2) and (1.4), respectively, where $0 < T < T_{max}$ and T_{max} is the maximal time for which*

a solution to (1.2) exists and is unique. For all $t \in [0, T]$, we have

$$\begin{aligned} & \|\mathbf{u}(t) - \mathbf{u}_\alpha(t)\|_{L^2}^2 + \alpha^2 \|\nabla(\mathbf{u}(t) - \mathbf{u}_\alpha(t))\|_{L^2}^2 \leq \\ & (\|\boldsymbol{\eta} - \boldsymbol{\eta}_\alpha\|_{L^2}^2 + \alpha^2 \|\nabla(\boldsymbol{\eta} - \boldsymbol{\eta}_\alpha)\|_{L^2}^2) e^{Ct} + C\alpha^2(e^{Ct} - 1). \end{aligned} \quad (2.7)$$

Based on this theorem we know that if

$$\|\boldsymbol{\eta} - \boldsymbol{\eta}_\alpha\|_{L^2}^2 + \alpha^2 \|\nabla(\boldsymbol{\eta} - \boldsymbol{\eta}_\alpha)\|_{L^2}^2 \rightarrow 0, \quad \text{as } \alpha \rightarrow 0, \quad (2.8)$$

then for all $t \in [0, T]$,

$$\|\mathbf{u}(t) - \mathbf{u}_\alpha(t)\|_{L^2}^2 + \alpha^2 \|\nabla(\mathbf{u}(t) - \mathbf{u}_\alpha(t))\|_{L^2}^2 \rightarrow 0. \quad (2.9)$$

Therefore, solutions of the Euler-Voigt system (1.4) converge to the corresponding Euler (1.2) flows in H^1 uniformly in time as $\alpha \rightarrow 0$. Using this result, the following theorem was proved in [17].

Theorem 2.1.2. *Let $\boldsymbol{\eta}_\alpha \in H^s$, $s \geq 3$, with $\nabla \cdot \boldsymbol{\eta}_\alpha = 0$ and let $\mathbf{u}_\alpha$ be the corresponding unique solution of (1.4). Suppose that there exists a time $T^* > 0$ such that*

$$\limsup_{\alpha \rightarrow 0^+} \left(\alpha \sup_{t \in [0, T^*]} \|\nabla \mathbf{u}_\alpha(t)\|_{L^2} \right) > 0. \quad (2.10)$$

Then, the corresponding solution of the Euler equations (1.2) with the initial condition $\boldsymbol{\eta}_\alpha$ must develop a singularity within the interval $[0, T^]$.*

We note that this theorem relies on the solutions of (1.2) and (1.4) starting from the same initial condition, i.e., $\boldsymbol{\eta} = \boldsymbol{\eta}_\alpha$. This necessarily satisfies (2.8) as $\alpha \rightarrow 0$. However, in order to use this result in our study, we need to generalize Theorem 2.1.2

for the case (2.8) where $\boldsymbol{\eta} \neq \boldsymbol{\eta}_\alpha$. Therefore, we modify Theorem 5.2 from [18] as follows.

Theorem 2.1.3 (X. Zhao, private communication). *Given initial data $\boldsymbol{\eta}, \boldsymbol{\eta}_\alpha \in H^s(\mathbb{T}^3)$ for some $s \geq 3$ satisfying*

$$\|\boldsymbol{\eta} - \boldsymbol{\eta}_\alpha\|_{L^2}^2 + \alpha^2 \|\nabla(\boldsymbol{\eta} - \boldsymbol{\eta}_\alpha)\|_{L^2}^2 \rightarrow 0, \quad \text{as } \alpha \rightarrow 0, \quad (2.11)$$

let $\mathbf{u}, \mathbf{u}_\alpha \in C([0, T], H^s(\mathbb{T}^3)) \cap C^1([0, T], H^{s-1}(\mathbb{T}^3))$ be the corresponding solutions to (1.2) and (1.4), respectively. If there exists a finite time $T^ > 0$ such that the solutions $\mathbf{u}_\alpha$ satisfy*

$$\limsup_{\alpha \rightarrow 0} \left(\alpha^2 \sup_{t \in [0, T^*]} \|\nabla \mathbf{u}_\alpha(t)\|_{L^2}^2 \right) > 0, \quad (2.12)$$

then $\mathbf{u}$ develops a singularity in the interval $[0, T^]$.*

Proof. Suppose that the solution $\mathbf{u}$ of the Euler equations (1.2) with the initial condition $\boldsymbol{\eta}$ is regular on $[0, T^*]$. Since the Euler-Voigt equations are globally well-posed, we have, cf. (1.6)

$$\|\mathbf{u}_\alpha(t)\|_{L^2}^2 + \alpha^2 \|\nabla \mathbf{u}_\alpha(t)\|_{L^2}^2 = \|\boldsymbol{\eta}_\alpha\|_{L^2}^2 + \alpha^2 \|\nabla \boldsymbol{\eta}_\alpha\|_{L^2}^2, \quad (2.13)$$

thus

$$\alpha^2 \|\nabla \mathbf{u}_\alpha(t)\|_{L^2}^2 = \|\boldsymbol{\eta}_\alpha\|_{L^2}^2 + \alpha^2 \|\nabla \boldsymbol{\eta}_\alpha\|_{L^2}^2 - \|\mathbf{u}_\alpha(t)\|_{L^2}^2. \quad (2.14)$$

Taking the supremum over $t \in [0, T^*]$ of the above equation, we obtain that

$$\sup_{t \in [0, T^*]} \alpha^2 \|\nabla \mathbf{u}_\alpha(t)\|_{L^2}^2 = \|\boldsymbol{\eta}_\alpha\|_{L^2}^2 + \alpha^2 \|\nabla(\boldsymbol{\eta}_\alpha)\|_{L^2}^2 - \inf_{t \in [0, T^*]} \|\mathbf{u}_\alpha(t)\|_{L^2}^2. \quad (2.15)$$

Taking the limsup as $\alpha \rightarrow 0$, we obtain

$$\limsup_{\alpha \rightarrow 0} \left(\sup_{t \in [0, T^*]} \alpha^2 \|\nabla \mathbf{u}_\alpha(t)\|_{L^2}^2 \right) = \limsup_{\alpha \rightarrow 0} \left(\|\boldsymbol{\eta}_\alpha\|_{L^2}^2 + \alpha^2 \|\nabla(\boldsymbol{\eta}_\alpha)\|_{L^2}^2 - \inf_{t \in [0, T^*]} \|\mathbf{u}_\alpha(t)\|_{L^2}^2 \right). \quad (2.16)$$

We now make use of Theorem 2.1.1, which demonstrates that the convergence of the solutions of the Euler-Voigt equations (1.4) to the solution of the Euler equations (1.2) is uniform on $[0, T^*]$. Therefore we can reduce the above equation to

$$\limsup_{\alpha \rightarrow 0} \left(\sup_{t \in [0, T^*]} \alpha^2 \|\nabla \mathbf{u}_\alpha(t)\|_{L^2}^2 \right) = \|\boldsymbol{\eta}\|_{L^2}^2 - \|\mathbf{u}(t)\|_{L^2}^2. \quad (2.17)$$

Since the energy of regular solutions of the Euler equations is conserved, we have

$$\limsup_{\alpha \rightarrow 0} \left(\sup_{t \in [0, T^*]} \alpha^2 \|\nabla \mathbf{u}_\alpha(t)\|_{L^2}^2 \right) = 0, \quad (2.18)$$

which contradicts the hypothesis in the theorem that the above expression is positive.

Equality (2.17) holds because

$$\begin{aligned} \limsup_{\alpha \rightarrow 0} \left| \inf_{t \in [0, T^*]} \|\mathbf{u}_\alpha(t)\|_{L^2} - \|\boldsymbol{\eta}\|_{L^2} \right| &\leq \limsup_{\alpha \rightarrow 0} \|\mathbf{u}_\alpha(t_\alpha) - \boldsymbol{\eta}\|_{L^2}^2 \\ &\leq \limsup_{\alpha \rightarrow 0} \|\mathbf{u}_\alpha(t_\alpha) - \mathbf{u}(t_\alpha)\|_{L^2}^2 + \|\mathbf{u}(t_\alpha) - \boldsymbol{\eta}\|_{L^2}^2 \\ &= 0, \end{aligned}$$

where the reverse triangle and triangle inequalities were used and t_α is the time at which

$$\inf_{t \in [0, T^*]} \|\mathbf{u}_\alpha(t)\|_{L^2} = \|\mathbf{u}_\alpha(t_\alpha)\|_{L^2}. \quad (2.20)$$

□

2.2 Initial Conditions

We have established Theorem 2.1.3 which we will use to identify singularity formation in Euler flows corresponding to smooth initial conditions. In order to formulate our optimization problem we must now describe the function space we perform our search in. The Theorems (2.1.1), (2.1.2), and (2.1.3) presented in Section (2.1) require initial conditions $\boldsymbol{\eta}, \boldsymbol{\eta}_\alpha \in H^s(\mathbb{T}^3)$ for some $s \geq 3$. However, the pseudo-spectral methods used in this study to numerically solve system (1.4) require the functions to be analytic in order to have exponential convergence [4]. For the Navier-Stokes equations (1.1), the viscous term acts to instantaneously regularize the solution in space. Therefore, if the initial condition has only Sobolev regularity, the resulting flow becomes real-analytic (with Gevrey regularity defined below) at $t = 0^+$. This regularity allows the solution to achieve exponential convergence.

The Euler equations (1.2) do not include any viscous effects and so the solution simply stays at the same level of smoothness as the initial condition, unless blow-up occurs. For Euler-Voigt the regularization term keeps the solution globally well-posed but does not improve the regularity [18]. Therefore, we must have analytic initial conditions to make use of the pseudo-spectral methods. For all the models considered here, a requirement is that the flow is divergence free at all times, including the initial conditions. Finally, on a periodic domain the quantity $\int_\Omega \boldsymbol{\eta}_\alpha d\mathbf{x}$ is an invariant of motion for solutions of the Euler-Voigt equations. Due to this, we assume without loss of generality that the initial conditions have a mean of zero. In order to identify initial

conditions that satisfy these criteria, we first consider analytic initial conditions that belong to the Gevrey class

$$G^\sigma := \left\{ \mathbf{v} \in H^s(\mathbb{T}^3) : \|\mathbf{v}\|_{G^\sigma} = \langle \mathbf{v}, \mathbf{v} \rangle_{G^\sigma}^{1/2} < \infty \right\}, \quad s \geq 3, \quad \sigma > 0, \quad (2.21)$$

with the norm defined as

$$\forall \mathbf{v}, \mathbf{u} \in G^\sigma, \quad \langle \mathbf{v}, \mathbf{u} \rangle_{G^\sigma} := \sum_{\mathbf{j} \in \mathbb{Z}^3} (1 + |2\pi \mathbf{j}|^2)^s e^{4\pi\sigma|\mathbf{j}|} \widehat{\mathbf{v}}_{\mathbf{j}} \cdot \overline{\widehat{\mathbf{u}}_{\mathbf{j}}} \quad (2.22a)$$

$$= \int_{\mathbb{T}^3} (1 + |D|^2)^s e^{2\sigma|D|} \mathbf{v} \cdot \mathbf{u} \, d\mathbf{x}, \quad (2.22b)$$

where hat denotes the Fourier coefficient

$$\widehat{\mathbf{u}}_k = \int_{\Omega} e^{-i\mathbf{k} \cdot \mathbf{x}} \mathbf{u}(\mathbf{x}) d\mathbf{x}, \quad (2.23)$$

overbar denotes complex conjugation, and the operators $|D|$ and $e^{\sigma|D|}$ are defined via

$$\left[\widehat{|D|\mathbf{v}} \right]_{\mathbf{j}} := 2\pi|\mathbf{j}| \widehat{\mathbf{v}}_{\mathbf{j}}, \quad \left[\widehat{e^{\sigma|D|}\mathbf{v}} \right]_{\mathbf{j}} := e^{2\pi|\mathbf{j}|} \widehat{\mathbf{v}}_{\mathbf{j}}. \quad (2.24)$$

The parameter σ determines the relative 'size' of the Gevrey space

$$G^{\sigma_2} \subset G^{\sigma_1} \subset G^0 = H^s, \quad 0 < \sigma_1 < \sigma_2, \quad (2.25)$$

which is proved in Appendix B. The value of σ is chosen in Section 4.3. In order to satisfy the divergence free condition, we introduce the following subspace in which

optimal initial conditions will be sought

$$V := \{\mathbf{v} \in G^\sigma : \nabla \cdot \mathbf{v} = 0, \quad \int_{\mathbb{T}^3} \mathbf{v} \, d\mathbf{x} = 0\}. \quad (2.26)$$

2.3 Scaling Symmetries of Euler and Euler-Voigt Systems

Both (1.2) and (1.4) possess a scaling property. Given a solution $\mathbf{u}$, or $\mathbf{u}_\alpha$, we can construct a new solution

$$\mathbf{v}(\mathbf{y}, \tau) = \frac{1}{\gamma} \mathbf{u}(\beta \mathbf{x}, \lambda t), \quad \mathbf{y} = \beta \mathbf{x}, \quad \tau = \lambda t \quad (2.27)$$

where $\lambda, \gamma, \beta > 0$ satisfy $\beta/(\lambda\gamma) = 1$, which is proven in Appendices C and D, respectively for the Euler (1.2) and Euler-Voigt (1.4) equations. Our goal is to find a time interval $[0, T]$ that is sufficiently large so that it is outside the interval of local existence established by Kato's theorem 1.1.1 and so allows for the potential of singularity formation in Euler flows. Due to this scaling property of the Euler-Voigt equations (2.27), the blow-up time of the solution will depend on the 'size' of our solution (γ) and the spatial scale (β). Our domain size will not change, and so in order to keep consistent time scale we must fix γ between all initial conditions. To accomplish this, we restrict our discussion to initial conditions with a fixed $\dot{H}^1$ seminorm which belong to a closed manifold $\mathcal{M}_C \subset V$ defined as cf. (2.26)

$$\mathcal{M}_C = \{\mathbf{v} \in V : \|\mathbf{v}\|_{\dot{H}^1} = C\}, \quad (2.28)$$

where $C > 0$ is a constant. We construct the manifold by fixing the $\dot{H}^1$ seminorm because of the form of our objective functional used in the optimization search, which

will be described in Chapter 3. The question we must answer now is what value of the constraint parameter C we use. To determine this we look to our first initial condition, the 3D Taylor Green Vortex (TGV). The TGV in the domain $\Omega = [0, L]^3$ has the form

$$\boldsymbol{\eta}_{TGV}(x) = \begin{bmatrix} V_0 \sin(2\pi x/L) \cos(2\pi y/L) \cos(2\pi z/L) \\ -V_0 \cos(2\pi x/L) \sin(2\pi y/L) \cos(2\pi z/L) \\ 0 \end{bmatrix}, \quad (2.29)$$

for some $V_0 \neq 0$. The $\dot{H}^1$ -seminorm of (2.29) is

$$\|\boldsymbol{\eta}_{TGV}\|_{\dot{H}^1} = \sqrt{3L\pi}V_0, \quad (2.30)$$

which is proven in Appendix E. In addition, the BKM criterion (1.1.2) uses the quantity $\|\boldsymbol{\omega}\|_{L^\infty}$ to determine singularity formation. For the TGV

$$\|\boldsymbol{\omega}\|_{L^\infty} = \|\nabla \times \boldsymbol{\eta}_{TGV}\|_{L^\infty} = 4\pi V_0/L, \quad (2.31)$$

which is proven in Appendix F. In [3], the authors provided numerical evidence that blow-up of (1.2) may occur around $T^* \approx 4$ for initial condition (2.29) with parameters $L = 1$ and $V_0 = 1$. This result provides the baseline for our study and so we must match this time window so that singularity formation identified by (2.1.3), if it happens, occurs at approximately the same time as in [3]. Therefore, we set $\lambda = 1$. Our Euler-Voigt system is defined on the domain $\Omega = [0, 1]^3$ and so we set $\beta = 1/2\pi$. Therefore, from (2.27) we set $\gamma = 1/2\pi$. The model we are aiming to match uses $V_0 = 1$ and so to match the time window we use $V_0 = 1/2\pi$.

These considerations give $\|\boldsymbol{\eta}_{TGV}\|_{\dot{H}^1} = \sqrt{3}/2$ and $\|\nabla \times \boldsymbol{\eta}_{TGV}\|_{L^\infty} = 2$ for the initial condition (2.29). Therefore, we can restrict our discussion of the TGV to initial conditions with $\dot{H}^1$ seminorm which belong to a closed manifold (2.28) with $C = \sqrt{3}/2$. In order to select initial conditions within this manifold, we will first select initial conditions $\boldsymbol{\eta} \in V$ and then apply a retraction function to ensure it has $\|\cdot\|_{\dot{H}^1} = \sqrt{3}/2$. To verify that this selection of the $\dot{H}^1$ seminorm of the initial condition is correct, we compare this scaling to the results shown in Figure 1b of [3]. This figure shows the evolution of the L^∞ norm of the vorticity over time.




Figure 2.1: Maximum of vorticity $\|\omega(\cdot, t)\|_\infty$. Results from runs performed at different resolutions are displayed together: 128^3 (orange plus), 256^3 (cyan star), 512^3 (brown triangles), 1024^3 (blue squares).

We present our reproduction in Figure 2.1 which shows that the evolution of the vorticity at each resolution in our scaled model matches exactly to the corresponding computations shown in Figure 1b in [3]. Therefore, our construction of the manifold $\mathcal{M}_C$ is correct.

2.4 Initial Guesses

Finally, before we begin our search for singularities, we examine the structure of the TGV and the other initial guesses used in this study. Figure 2.2 shows the isosurfaces of the vorticity of the scaled TGV.




Figure 2.2: Isosurfaces of the scaled TGV vorticity. (a) magnitude of $\boldsymbol{\omega}$, vorticity components (b) ω_1 , (c) ω_2 , and (d) ω_3 .

ADD AND COMPARE THE OTHER INITIAL GUESSES HERE

3 The Optimization Problem

In this chapter we ~~construct the~~ optimization problem (3.1.1) and the Riemannian conjugate gradient method used to solve it.

3.1 Formulation of the Problem

With our initial condition and blowup criterion defined by Theorem 2.1.3, we now construct the PDE-constrained optimization problem. We begin with the ~~objection~~ functional $\Phi_{\alpha,T} : V \rightarrow \mathbb{R}^+$, which is defined as

$$\Phi_{\alpha,T}(\boldsymbol{\eta}_\alpha) := \|\nabla \mathbf{u}_\alpha(T; \boldsymbol{\eta}_\alpha)\|_{L^2}^2, \quad (3.32a)$$

$$\doteq \|\mathbf{u}_\alpha(T; \boldsymbol{\eta}_\alpha)\|_{H^1}^2, \quad (3.32b)$$

where $\mathbf{u}_\alpha(T; \boldsymbol{\eta}_\alpha)$ is the solution of the Euler-Voigt equations (1.4) obtained at time T with the initial condition $\boldsymbol{\eta}_\alpha \in V$ and regularization parameter α . Using this objective functional, we define the optimization problem.

Problem 3.1.1. *Given $\alpha, T \in \mathbb{R}^+$, find*

$$\tilde{\boldsymbol{\eta}}_{\alpha,T} = \operatorname{argmax}_{\boldsymbol{\eta}_\alpha \in M_C} \Phi_{\alpha,T}(\boldsymbol{\eta}_\alpha). \quad (3.33)$$

Our goal is to first fix T and solve the optimization problem for a sequence of decreasing values of α that converge to 0. From Figure 1b of [3], the expected time of singularity formation in the Euler flow is $T^* \approx 4$. From Theorem 2.1.3 if the initial condition criterion (2.8) and objective functional criteria (2.12) are both satisfied, then the corresponding Euler flow develops a singularity in the interval $[0, T^*]$ from initial condition η . We examine the results for a time interval before the expected point singularity formation should occur and again for a time interval after. Therefore, we consider the optimization problem for $T = 3$, where blow-up should not occur, and $T = 5$ where it may occur.

3.2 Gradient of Objective Function

The optimization problem is solved using a Riemannian conjugate gradient method which uses an optimal search direction and distance to create the next approximation of the optimal initial condition. This method is utilized because it accelerates the gradient ascent method which is necessary given the computational cost of each iteration. To solve this problem we utilize a 'optimize-then-discretize' approach where the optimization problem, and the Riemannian conjugate gradient method, are derived in an infinite dimensional function space and then discretized to solve numerically. The same method was used in [27] to solve a PDE-constrained optimization problem for the Euler equations to search for initial conditions η that produce 'extreme' flows.

We compute the optimal search direction using the gradient of the objection functional $\nabla \Phi_{\alpha, T}(\eta_\alpha)$. It is obtained using an adjoint method which depends on solving a properly defined adjoint system backwards in time. To derive an expression for

the gradient, we first introduce the Gâteaux (directional) differential $\Phi'_{\alpha,T}(\boldsymbol{\eta}_\alpha; \boldsymbol{\eta}'_\alpha) : V \times V \rightarrow \mathbb{R}$. **The Gateaux differential is defined as,**

$$\Phi'_{\alpha,T}(\boldsymbol{\eta}_\alpha; \boldsymbol{\eta}'_\alpha) := \lim_{\epsilon \rightarrow 0} \frac{1}{\epsilon} [\Phi_{\alpha,T}(\boldsymbol{\eta}_\alpha + \epsilon \boldsymbol{\eta}'_\alpha) - \Phi_{\alpha,T}(\boldsymbol{\eta}_\alpha)]. \quad (3.34)$$

This represents the change in $\Phi_{\alpha,T}$ from **an infinitesimal perturbation proportional to ϵ** in the direction $\boldsymbol{\eta}'_\alpha$ from the initial condition $\boldsymbol{\eta}_\alpha$. By fixing the initial condition $\boldsymbol{\eta}_\alpha$, we can view the Gâteaux differential as a bounded linear functional on V . Therefore, it can also be represented as an inner product on the Gevrey space using the Riesz representation theorem [27, 20]

$$\Phi'_{\alpha,T}(\boldsymbol{\eta}_\alpha; \boldsymbol{\eta}'_\alpha) = \langle \nabla \Phi_{\alpha,T}(\boldsymbol{\eta}_\alpha), \boldsymbol{\eta}'_\alpha \rangle_{G^\sigma}. \quad (3.35)$$

Substituting the second definition of the objective functional (3.32) into **the Gâteaux differential (3.34)**, we derive the Gâteaux differential in the form of the $\dot{H}^1$ inner product,

$$\Phi'_{\alpha,T}(\boldsymbol{\eta}_\alpha; \boldsymbol{\eta}'_\alpha) = 2 \langle \mathbf{u}_\alpha(\cdot, T), \mathbf{u}'_\alpha(\cdot, T) \rangle_{\dot{H}^1} \quad (3.36)$$

The derivation for this is shown in Appendix I. In (3.36) the value $\mathbf{u}'_\alpha$ is the solution of the linearization of the Euler-Voigt system (1.4) around its solution **$\mathbf{u}_\alpha$ corresponding to the initial condition $\boldsymbol{\eta}_\alpha$** . The linearization of the Euler-Voigt system is defined as

$$\mathcal{L} \begin{bmatrix} \mathbf{u}'_\alpha \\ p' \end{bmatrix} := \begin{bmatrix} \partial_t \mathbf{u}'_\alpha + (I - \alpha^2 \Delta)^{-1} (\mathbf{u}_\alpha \cdot \nabla \mathbf{u}'_\alpha + \mathbf{u}'_\alpha \cdot \nabla \mathbf{u}_\alpha + \nabla p') = 0 \\ \nabla \cdot \mathbf{u}'_\alpha = 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \quad (3.37a)$$

$$\mathbf{u}'_\alpha(x, 0) = \boldsymbol{\eta}'_\alpha \quad (3.37b)$$

where $\boldsymbol{\eta}'_\alpha$ and p' are the perturbations of the initial condition and pressure, respectively. The derivation of the linearized form of the Euler-Voigt equations is shown in Appendix G. In order to obtain the gradient of the objective functional from the Gâteaux differential we must first introduce the fields $\{\mathbf{u}_\alpha^*, p_\alpha^*\}$ which are the solution of the adjoint system

$$\mathcal{L}^* \begin{bmatrix} \mathbf{u}_\alpha^* \\ p_\alpha^* \end{bmatrix} := \begin{bmatrix} -\partial_t \mathbf{u}_\alpha^* - (\nabla(I - \alpha^2)^{-1} \mathbf{u}_\alpha^* + (\nabla(I - \alpha^2)^{-1} \mathbf{u}_\alpha^*)^\top) \mathbf{u}_\alpha - \nabla p_\alpha^* \\ -\nabla \cdot \mathbf{u}_\alpha^* \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \quad (3.38a)$$

$$\mathbf{u}_\alpha^*(\mathbf{x}, T) = 2|D|^2 \mathbf{u}_\alpha(\mathbf{x}, T) \quad (3.38b)$$

which is derived in Appendix H by performing integration by parts, in both space and time, on the following duality-pairing relation

$$\begin{aligned} \left(\mathcal{L} \begin{bmatrix} \mathbf{u}'_\alpha \\ p' \end{bmatrix}, \begin{bmatrix} \mathbf{u}_\alpha^* \\ p^* \end{bmatrix} \right) &:= \int_0^T \int_{\mathbb{T}^3} \mathcal{L} \begin{bmatrix} \mathbf{u}'_\alpha \\ p' \end{bmatrix}, \begin{bmatrix} \mathbf{u}_\alpha^* \\ p^* \end{bmatrix} d\mathbf{x} dt \\ &= \left(\begin{bmatrix} \mathbf{u}'_\alpha \\ p' \end{bmatrix}, \mathcal{L}^* \begin{bmatrix} \mathbf{u}_\alpha^* \\ p^* \end{bmatrix} \right) + \int_{\mathbb{T}^3} \mathbf{u}'_\alpha(\mathbf{x}, T) \cdot \mathbf{u}_\alpha^*(\mathbf{x}, T) d\mathbf{x} \\ &\quad - \int_{\mathbb{T}^3} \boldsymbol{\eta}'_\alpha(\mathbf{x}) \cdot \mathbf{u}_\alpha^*(\mathbf{x}, 0) d\mathbf{x} \\ &= 0. \end{aligned} \quad (3.39)$$

Using the solutions of the adjoint system we can now obtain a convenient expression for the gradient. We take the Gâteaux differential (3.36) in terms of the L^2 inner product

$$\Phi'_{\alpha, T}(\boldsymbol{\eta}_\alpha; \boldsymbol{\eta}'_\alpha) = \langle \mathbf{u}_\alpha^*(\cdot, 0), \mathbf{u}'_\alpha(\cdot, 0) \rangle_{L^2}, \quad (3.40)$$

which is derived in Appendix I. We take the Riesz representation of the Gâteaux

differential in terms of the L^2 inner product (3.40) and in the Riesz representation form (3.35)

$$\Phi'_{\alpha,T}(\boldsymbol{\eta}_\alpha; \boldsymbol{\eta}'_\alpha) = \langle \mathbf{u}_\alpha^*(\cdot, 0), \mathbf{u}'_\alpha(\cdot, 0) \rangle_{L^2} = \langle \nabla \Phi_{\alpha,T}(\boldsymbol{\eta}_\alpha), \boldsymbol{\eta}'_\alpha \rangle_{G^\sigma}. \quad (3.41)$$

Using the definition of the Gevrey inner product (2.22), the L^2 inner product can be converted to the Gevrey inner product,

$$\langle \mathbf{u}_\alpha^*(\cdot, 0), \mathbf{u}'_\alpha(\cdot, 0) \rangle_{L^2} = \langle (1 + |D|^2)^{-3} e^{-2\sigma|D|} \mathbf{u}_\alpha^*(x, 0), \boldsymbol{\eta}'_\alpha(x) \rangle_{G^\sigma}. \quad (3.42)$$

Therefore, we have two forms of the Gâteaux differential as a Gevrey inner product

$$\Phi'_{\alpha,T}(\boldsymbol{\eta}_\alpha; \boldsymbol{\eta}'_\alpha) = \langle \nabla \Phi_{\alpha,T}(\boldsymbol{\eta}_\alpha), \boldsymbol{\eta}'_\alpha \rangle_{G^\sigma} = \langle (1 + |D|^2)^{-3} e^{-2\sigma|D|} \mathbf{u}_\alpha^*(x, 0), \boldsymbol{\eta}'_\alpha(x) \rangle_{G^\sigma}. \quad (3.43)$$

From this we obtain the gradient of the objective functional

$$\nabla \Phi_{\alpha,T}(\boldsymbol{\eta}_\alpha) = (1 + |D|^2)^{-3} e^{-2\sigma|D|} \mathbf{u}_\alpha^*(x, 0). \quad (3.44)$$

The adjoint system (3.38) is linear and a terminal value problem. Therefore, its coefficients are determined by evolving the Euler-Voigt system forward from the initial condition $\boldsymbol{\eta}_\alpha$ to compute $\mathbf{u}_\alpha(\mathbf{x}, T)$, from which we obtain the terminal value of (3.38). Finally, we evolve the adjoint system backwards in time to obtain the initial value $\mathbf{u}_\alpha^*(\mathbf{x}, 0)$ which is used to compute the gradient (3.44).

3.3 Riemannian Conjugate Gradient Method

When solving the optimization problem, we look for an initial condition $\boldsymbol{\eta}_\alpha \in M_C$ that maximizes our objective functional. Therefore, not only must our initial guess $\boldsymbol{\eta}_{\alpha,T}^0$ have a $\dot{H}^1$ seminorm of $\sqrt{3}/2$, but so too must each following iterate $\boldsymbol{\eta}_{\alpha,T}^n$. We ensure this constraint is satisfied using a Riemannian conjugate gradient method modified from the method used in [25] and [9]. This method has three steps:

1. Project the gradient $\nabla\Phi_T(\boldsymbol{\eta}_\alpha)$ onto the tangent space to $\mathcal{M}_C$ at $\boldsymbol{\eta}_\alpha$.
2. Perform a suitable vector transport operation in order to construct a Riemannian conjugate ascent direction using the previous search direction.
3. Retract the resulting state back to the constraint manifold $\mathcal{M}_c$.

We first define the space tangent to $\mathcal{M}_C$ at $\boldsymbol{v}$ as

$$T_{\boldsymbol{v}}\mathcal{M}_C = \{\boldsymbol{z} \in V : \langle \boldsymbol{v}, \boldsymbol{z} \rangle_{\dot{H}^1} = 0\}. \quad (3.45)$$

For any $\boldsymbol{z} \in T_{\boldsymbol{v}}\mathcal{M}_C$, this condition can be expressed as

$$\langle \boldsymbol{v}, \boldsymbol{z} \rangle_{\dot{H}^1} = \int_{\mathbb{T}^3} |D| \boldsymbol{v} \cdot |D| \boldsymbol{z} d\boldsymbol{x} \quad (3.46a)$$

$$= \int_{\mathbb{T}^3} (1 + |D|^2)^3 e^{2\sigma|D|} \left((1 + |D|^2)^{-3} e^{-2\sigma|D|} |D|^2 \boldsymbol{v} \right) \cdot \boldsymbol{z} d\boldsymbol{x} \quad (3.46b)$$

$$= \langle (1 + |D|^2)^{-3} e^{-2\sigma|D|} |D|^2 \boldsymbol{v}, \boldsymbol{z} \rangle_{G^\sigma} = 0. \quad (3.46c)$$

Thus, at each point $\boldsymbol{v} \in M_C$, the tangent space $T_{\boldsymbol{v}}\mathcal{M}_C$ can be characterized using the unit 'normal' vector given by

$$\boldsymbol{n}_{\boldsymbol{v}} = \frac{(1 + |D|^2)^{-3} e^{-2\sigma|D|} |D|^2 \boldsymbol{v}}{\|(1 + |D|^2)^{-3} e^{-2\sigma|D|} |D|^2 \boldsymbol{v}\|_{G^\sigma}} \quad (3.47)$$

which allows us to construct the projection operator $\mathcal{P}_{\mathcal{T}_v \mathcal{M}_C} : V \rightarrow \mathcal{T}_v \mathcal{M}_C$,

$$\forall \boldsymbol{\omega} \in V, \quad \mathcal{P}_{\mathcal{T}_v \mathcal{M}_C} \boldsymbol{\omega} = \boldsymbol{\omega} - \langle \mathbf{n}_v, \boldsymbol{\omega} \rangle_{G^\sigma} \mathbf{n}_v. \quad (3.48)$$

As stated previously, the initial conditions at each iteration must have a $\dot{H}^1$ -seminorm of $\sqrt{3}/2$. In order to ensure this we introduce the retraction operator

$$R(\mathbf{v}) = \frac{\sqrt{3}}{2} \frac{\mathbf{v}}{\|\mathbf{v}\|_{\dot{H}^1}}, \quad \mathbf{v} \neq \mathbf{0}. \quad (3.49)$$

The **conjugate-gradient method** uses the search direction obtained at the n -th iteration to compute the search direction at the $(n+1)$ -th iteration. This is done by **computing a linear combination of the gradient at the current iteration and the search direction from the previous iteration, weighted with a 'momentum' coefficient**. However, the two directions belong to different tangent spaces to $\mathcal{M}_C$ and thus cannot be directly added. In order to allow algebraic operations to be performed on the vectors of the two tangent spaces we introduce the vector transport operation defined by [?]]

$$\mathcal{T}\mathcal{M}_C \oplus \mathcal{T}\mathcal{M}_C \rightarrow \mathcal{T}\mathcal{M}_C \quad : \quad (\boldsymbol{\varphi}, \boldsymbol{\xi}) \mapsto \Gamma_{\boldsymbol{\varphi}}(\boldsymbol{\xi}) \in \mathcal{T}\mathcal{M}_C,$$

where $\mathcal{T}\mathcal{M}_C = \bigcup_{v \in \mathcal{M}_C} \mathcal{T}_v \mathcal{M}_C$ is the tangent bundle on $\mathcal{M}_C$. The operation transports the vector field $\boldsymbol{\xi}$ along the manifold $\mathcal{M}_C$ in the direction of $\boldsymbol{\varphi}$. This allows us to map search directions between the tangent spaces of two consecutive iterations.

This operator is defined via differentiated retraction

$$\begin{aligned}\Gamma_{\varphi_u}(\xi_u) &:= \frac{d}{dt} \mathbf{R}(\varphi_u + t\xi_u) \Big|_{t=0} \\ &:= \frac{\sqrt{3}}{2} \frac{1}{\|\mathbf{u} + \varphi_u\|_{\dot{H}^1}} \left[\xi_u - \frac{\langle \mathbf{u} + \varphi_u, \xi_u \rangle_{\dot{H}^1}}{\|\mathbf{u} + \varphi_u\|_{\dot{H}^1}^2} (\mathbf{u} + \varphi_u) \right], \mathbf{u} \in M_C\end{aligned}$$

With our operators defined we can now construct the iterative relation that will be used to maximization our objective functional (3.32)

$$\boldsymbol{\eta}_\alpha^{(n+1)} = R \left[\boldsymbol{\eta}_\alpha^{(n)} + \tau_n \mathbf{d}^{(n)} \right]. \quad (3.51)$$

To simply notation, let $\mathbf{G}^{(n)} := \nabla \Phi_{\alpha,T}(\boldsymbol{\eta}_\alpha^{(n)})$ and let $\mathcal{T}_n := \mathcal{T}_{\boldsymbol{\eta}_\alpha^{(n)}} \mathcal{M}_C$. The search direction at iteration n ($\mathbf{d}^{(n)}$) is computed as

$$\begin{aligned}\mathbf{d}^{(0)} &= P_{\mathcal{T}_0} \mathbf{G}^{(0)} \\ \mathbf{d}^{(n)} &= P_{\mathcal{T}_n} \mathbf{G}^{(n)} + \beta_n \Gamma_{\tau_{n-1}} \mathbf{d}^{(n-1)}(\mathbf{d}^{(n-1)}), \quad n \geq 1,\end{aligned}$$

where the 'momentum' term is computed using the Polak-Ribière approach [24].

$$\beta_n = \frac{\left\langle P_{\mathcal{T}_n} \mathbf{G}^{(n)}, \left(P_{\mathcal{T}_n} \mathbf{G}^{(n)} - \Gamma_{\tau_{n-1}} \mathbf{d}^{(n-1)} P_{\mathcal{T}_{n-1}} \mathbf{G}^{(n-1)} \right) \right\rangle_{G^\sigma}}{\|P_{\mathcal{T}_{n-1}} \mathbf{G}^{(n-1)}\|_{G^\sigma}^2}. \quad (3.53)$$

In order to remove old data that may slow down convergence, the algorithm is periodically restarted by setting $\beta_n = 0$. For our optimization search, we restart the algorithm every 20 iterations, and when the search direction is not an ascent direction, i.e, when

$$\frac{\left\langle \mathbf{d}^{(n)}, P_{\mathcal{T}_n} \mathbf{G}^{(n)} \right\rangle_{G^\sigma}}{\|\mathbf{d}^{(n)}\|_{G^\sigma} \|P_{\mathcal{T}_n} \mathbf{G}^{(n)}\|_{G^\sigma}} < 1, \quad (3.54)$$

Finally, our optimal step size is determined by the distance along $\mathbf{d}^{(n)}$ which maximizes the objective functional, which is computed by performing an arc search along this direction using a modification of Brent's method.

$$\tau_n = \operatorname{argmax}_{\tau > 0} \Phi_T \left(R \left[\eta^{(n)} + \tau \mathbf{d}^{(n)} \right] \right). \quad (3.55)$$

The optimization search is terminated when the relative change in the objective functional from (3.51) is below a chosen tolerance.

$$0 \leq \frac{\Phi_{\alpha,T}(\boldsymbol{\eta}_\alpha^{(n+1)}) - \Phi_{\alpha,T}(\boldsymbol{\eta}_\alpha^{(n)})}{\Phi_{\alpha,T}(\boldsymbol{\eta}_\alpha^{(n)})} < tol, \quad (3.56)$$

where $tol = 1E-10$.

4 Numerical Methods

4.1 Spatial Discretization & Spectral Methods

Simulations were computed using a standard pseudospectral fourier method on a periodic unit cube, discretized in a uniform grid of spatial resolutions $N = 128^3, 256^3, 512^3, 1024^3$. Derivatives are evaluated in the Fourier space and nonlinear products are computed in the physical space. The standard 2/3 dealiasing rule is also applied. Our interest is in the limit $\alpha \rightarrow 0$. Therefore, we use the regularization parameters $\alpha = 2^{-6}, 2^{-7}, 2^{-8}, 2^{-9}, 2^{-10}$.

4.2 Time Stepping

Time stepping is performed with the explicit fourth-order Runge-Kutta (RK4) method. The determination of our time step size Δt is based on three factors. First, we consider the numerical stability of the models. Second, we consider the accuracy of the solution, determined by the relative error of the objective functional (3.32) to the results for a minimum step size. Finally, we consider the conservation of the α -energy (1.6), determined by the maximum relative error of $E_\alpha(t)$ to $E_\alpha(0)$. These factors are analysed for each resolution and value of α for the Taylor Green Vortex.

The goal of our optimization problem is to identify initial conditions $\tilde{\eta}_{\alpha,T}$ that produce 'extreme' behaviour in solutions to the Euler-Voigt equations. Therefore, the Taylor Green Vortex will by definition be the least extreme initial condition examined. To ensure stability, accuracy, and conservation of α -energy are maintained upon optimization we must select a step size smaller than the minimum step size identified by these criteria.

4.2.1 Numerical Stability & Accuracy

We begin with the stability and accuracy of the Euler-Voigt model. To determine the stability and accuracy of the model at each step size we start from the iteration zero initial conditions and evaluate the objective functional (3.32) for the time window $T = 5$. First we do this for $\Delta t = 2^{-8}$ and designate the result as our 'correct' value Φ_c . We then compute the relative error of the objective functional for each value of Δt (Φ_{dt}) to Φ_c

$$\epsilon_{rel} = \left| \frac{\Phi_c - \Phi_{dt}}{\Phi_c} \right|. \quad (4.57)$$

For models that experienced numerical instability in the time window $[0, T]$ we set ϵ_{rel} to infinity.

Tables (4.1) to (4.4) show the results for ϵ_{rel} at the four resolutions utilized in this analysis. The tables shows that the accuracy of the objective functional decreases as $\alpha \rightarrow 0$ and increases as $\Delta t \rightarrow 0$. In addition, the accuracy of the model overall increases as the resolution increases. Finally, the size of the instability region does increase with resolution, but only for the smallest value of α . For all other values of α , the stability is unaffected over the time window $t \in [0, T]$.

		α				
		1/1024	2/1024	4/1024	8/1024	16/1024
Δt	2^{-2}	Inf	Inf	Inf	0.0045	7.00E-05
	2^{-3}	Inf	Inf	0.0572	0.0011	3.64E-06
	2^{-4}	0.1662	0.1157	0.0332	4.85E-04	2.02E-07
	2^{-5}	0.1093	0.0715	0.0181	2.33E-04	1.21E-08
	2^{-6}	0.0593	0.0369	0.0085	1.02E-04	8.63E-10
	2^{-7}	0.0228	0.0137	0.003	3.45E-05	9.44E-11
	2^{-8}	0	0	0	0	0

Table 4.1: Relative error of Φ_{dt} for each step size to Φ_c . Resolution $N = 128^3$ and time window $T = 5$.

		α				
		1/1024	2/1024	4/1024	8/1024	16/1024
Δt	2^{-2}	Inf	Inf	Inf	0.003	7.00E-05
	2^{-3}	Inf	Inf	0.0112	1.82E-04	3.64E-06
	2^{-4}	Inf	0.0652	0.0021	9.46E-06	2.01E-07
	2^{-5}	0.1097	0.0348	7.13E-04	5.12E-07	1.17E-08
	2^{-6}	0.0619	0.0171	3.08E-04	3.00E-08	7.01E-10
	2^{-7}	0.025	0.0157	1.05E-04	1.96E-09	4.04E-11
	2^{-8}	0	0	0	0	0

Table 4.2: Relative error of Φ_{dt} for each step size to Φ_c . Resolution $N = 256^3$ and time window $T = 5$.

		α				
		1/1024	2/1024	4/1024	8/1024	16/1024
Δt	2^{-2}	Inf	Inf	Inf	0.003	7.00E-05
	2^{-3}	Inf	Inf	0.009	1.82E-04	3.64E-06
	2^{-4}	Inf	0.0174	7.61E-04	9.45E-06	2.01E-07
	2^{-5}	Inf	0.0036	3.63E-05	5.10E-07	1.17E-08
	2^{-6}	0.0306	7.85E-04	1.72E-06	2.90E-08	7.01E-10
	2^{-7}	0.0115	2.42E-04	8.42E-08	1.63E-09	4.04E-11
	2^{-8}	0	0	0	0	0

Table 4.3: Relative error of Φ_{dt} for each step size to Φ_c . Resolution $N = 512^3$ and time window $T = 5$.

		α				
		1/1024	2/1024	4/1024	8/1024	16/1024
Δt	2^{-2}	Inf	Inf	Inf	0.003	7.00E-05
	2^{-3}	Inf	Inf	0.009	1.82E-04	3.64E-06
	2^{-4}	Inf	0.015	7.61E-04	9.45E-06	2.01E-07
	2^{-5}	Inf	0.0022	3.64E-05	5.10E-07	1.17E-08
	2^{-6}	0.0039	2.64E-04	1.71E-06	2.90E-08	7.01E-10
	2^{-7}	5.17E-04	4.92E-06	8.36E-08	1.63E-09	4.04E-11
	2^{-8}	0	0	0	0	0

Table 4.4: Relative error of Φ_{dt} for each step size to Φ_c . Resolution $N = 1024^3$ and time window $T = 5$.

4.2.2 Conservation of α -Energy

The conservation of α -energy is also used to verify the model for a given resolution, regularization parameter, and time window. The RK4 method does not conserve energy, and so the α -energy of our model gradually decreases over time. The tables below show the maximum relative error of the α -energy for each α and Δt ,

$$\epsilon_\alpha = \max_{t \in [0, T]} \left| \frac{E_\alpha(t) - E_\alpha(0)}{E_\alpha(t)} \right|. \quad (4.58)$$

For models that experienced numerical instability in the time window $[0, T]$ we set ϵ_α to infinity.

The Tables show that the conservation of α -energy error ϵ_α increases as $\alpha \rightarrow 0$ and decreases as $\Delta t \rightarrow 0$. In addition, the error decreases as the resolution increases.

Based on the results of the stability, accuracy, and α -energy conservation we can now select the appropriate time step size for each α and resolution, shown in Table (4.9). These step sizes are selected because they are the largest step sizes that ensure

		α				
		1/1024	2/1024	4/1024	8/1024	16/1024
Δt	2^{-2}	Inf	Inf	Inf	3.82E-04	8.73E-06
	2^{-3}	Inf	Inf	1.94E-03	6.91E-05	3.72E-07
	2^{-4}	3.38E-03	2.74E-03	1.11E-03	3.05E-05	1.76E-08
	2^{-5}	2.38E-03	1.81E-03	6.49E-04	1.57E-05	9.59E-10
	2^{-6}	1.51E-03	1.09E-03	3.56E-04	8.05E-06	7.55E-11
	2^{-7}	8.67E-04	6.06E-04	1.87E-04	4.08E-06	1.47E-11
	2^{-8}	4.69E-04	3.21E-04	9.59E-05	2.05E-06	6.00E-12

Table 4.5: Maximum relative error of $E_\alpha(t)$ to $E_\alpha(0)$ for each step size. Resolution $N = 128^3$ and time window $T = 5$.

		α				
		1/1024	2/1024	4/1024	8/1024	16/1024
Δt	2^{-2}	Inf	Inf	Inf	2.91E-04	8.73E-06
	2^{-3}	Inf	Inf	4.34E-04	1.49E-05	3.72E-07
	2^{-4}	Inf	8.36E-04	5.61E-05	6.85E-07	1.75E-08
	2^{-5}	9.63E-04	4.35E-04	1.80E-05	3.40E-08	9.09E-10
	2^{-6}	6.33E-04	2.48E-04	8.97E-06	1.89E-09	4.95E-11
	2^{-7}	3.83E-04	1.37E-04	4.59E-06	1.38E-10	2.89E-12
	2^{-8}	2.14E-04	7.23E-05	2.33E-06	2.37E-11	2.95E-12

Table 4.6: Maximum relative error of $E_\alpha(t)$ to $E_\alpha(0)$ for each step size. Resolution $N = 256^3$ and time window $T = 5$.

		α				
		1/1024	2/1024	4/1024	8/1024	16/1024
Δt	2^{-2}	Inf	Inf	Inf	2.91E-04	8.73E-06
	2^{-3}	Inf	Inf	3.85E-04	1.49E-05	3.72E-07
	2^{-4}	Inf	2.72E-04	2.58E-05	6.85E-07	1.75E-08
	2^{-5}	Inf	2.89E-05	1.12E-06	3.38E-08	9.09E-10
	2^{-6}	1.58E-04	8.18E-06	5.00E-08	1.82E-09	4.95E-11
	2^{-7}	9.02E-05	3.74E-06	2.47E-09	1.03E-10	3.44E-12
	2^{-8}	4.98E-05	1.91E-06	1.42E-10	6.31E-12	2.69E-12

Table 4.7: Maximum relative error of $E_\alpha(t)$ to $E_\alpha(0)$ for each step size. Resolution $N = 512^3$ and time window $T = 5$.

		α				
		1/1024	2/1024	4/1024	8/1024	16/1024
Δt	2^{-2}	Inf	Inf	Inf	2.91E-04	8.73E-06
	2^{-3}	Inf	Inf	3.85E-04	1.49E-05	3.72E-07
	2^{-4}	Inf	2.53E-04	2.58E-05	6.85E-07	1.75E-08
	2^{-5}	Inf	2.67E-05	1.12E-06	3.38E-08	9.09E-10
	2^{-6}	1.46E-05	1.26E-06	4.99E-08	1.82E-09	4.95E-11
	2^{-7}	2.53E-06	5.12E-08	2.45E-09	1.03E-10	3.25E-12
	2^{-8}	1.06E-06	2.35E-09	1.30E-10	6.31E-12	2.78E-12

Table 4.8: Maximum relative error of $E_\alpha(t)$ to $E_\alpha(0)$ for each step size. Resolution $N = 1024^3$ and time window $T = 5$.

		α				
		1/1024	2/1024	4/1024	8/1024	16/1024
N	128^3	2^{-8}	2^{-8}	2^{-6}	2^{-5}	2^{-3}
	256^3	2^{-8}	2^{-8}	2^{-6}	2^{-4}	2^{-3}
	512^3	2^{-8}	2^{-7}	2^{-5}	2^{-4}	2^{-3}
	1024^3	2^{-8}	2^{-7}	2^{-5}	2^{-4}	2^{-3}

Table 4.9: Selection of Δt for each resolution and α .

the stability, accuracy and α -energy conservation of the model from the Taylor Green Vortex.

4.3 Determination of Gevrey class size

In Section (2.2) we introduce the Gevrey space in which we search for initial conditions (2.21). The size of the space is determined by two variables, $s \geq 3$ and $\sigma > 0$. We use a fixed value of $s = 3$. In order to determine the value of σ we start with the optimization problem for $\alpha = 16/1024$, $T = 5$, and resolution $N = 128^3$. The problem is repeated for $\sigma = 1\text{E-}1, 1\text{E-}2, 1\text{E-}3, 1\text{E-}4, 1\text{E-}5$.

4.4 Spectrum Validation

The energy spectrum of the solution $\mathbf{u}_\alpha(\mathbf{x}, t)$ is defined as,

$$e(k, t) := \frac{1}{2} \sum_{j \leq |j| < j+1} |\hat{\mathbf{u}}_{\alpha j}(t)|^2, \quad k = 2\pi j, \quad j \in \mathbb{N}. \quad (4.59)$$

The regularization term of (1.4) acts as a filter which suppresses the growth of small-scale structures of length-scale smaller than the regularization parameter α . Therefore, the spectrum (4.59) is suppressed for wavenumbers $k > k_\alpha$ where

$$k_\alpha := \frac{2\pi}{\alpha}. \quad (4.60)$$

Due to this filter, the solution to (1.4) can remain well resolved. This is ideal for our analysis because it ensures that the model fully captures the small scale structures of the fluid. When this occurs we can be certain that our objective functional (3.32) is accurate. We begin by examining the spectrum of the solution from the Taylor Green Vortex on the time window $[0, T]$ for $T = 5$.

The plots () to () show the semi-log energy vs wavenumber graph of the solution over time.

initial condition shown below demonstrates the exponential decay expected by a analytic function. As the solution evolves in time the rate of decay decreases due to aliasing. While this can be reduced by anti-aliasing techniques, it cannot be entirely prevented

Low resolution and small alpha simulations are especially vulnerable to aliasing errors and the solutions stop being analytic before the chosen timescale.

The chosen timescales, alphas, and resolutions are validated in the graphs below which show that the spectral decay remains exponential and thus the solutions remain analytic.

4.5 Kappa Test Validation

The simulations are discretized in space using a psuedospectral method and time using a fourth order Runge Kutta method. The kappa test is a means of validating these discretization methods based on the validation method of [?]. We define the value κ as

$$\kappa(\epsilon) = \frac{\epsilon^{-1} [\Phi_{\alpha,T}(\boldsymbol{\eta} + \epsilon\boldsymbol{\eta}') - \Phi_{\alpha,T}(\boldsymbol{\eta})]}{\langle \nabla \Phi_{\alpha,T}(\boldsymbol{\eta}), \boldsymbol{\eta}' \rangle} \quad (4.61)$$

where $\Phi_{\alpha,T}$ is the objective functional (3.32), $\boldsymbol{\eta}$ is the initial condition, and $\nabla \Phi_{\alpha,T}$ is the gradient of the objective functional (3.44). The numerator is the first order finite difference approximation of the Gateaux differential. The denominator is the Reisz form of the differential that utilizes the adjoint system (??). If the numerical approximation methods are accurate, then we expect $\kappa \approx 1$.

A log-log plot of $|\kappa - 1|$ vs ϵ can be broken up into three regions, each dominated by different error sources. These are the discretization errors, round off errors, and truncation errors. The round off errors are caused by finite precision numerical representation and so are approximately of order $O(\epsilon^{-1})$. Therefore, we expect the value

of κ to be dominated by round off errors for small ϵ values. The truncation errors are caused by using a first order finite difference formula to calculate the derivative. The truncation errors are of order $O(\epsilon)$. Therefore, we expect the value of κ to be dominated by the truncation errors for large values of ϵ . Between these two regions the value of κ will be determined by the discretization errors in the numerical solutions, which are themselves a product of the spatial and temporal resolution. With smaller values of Δx and Δt causing the error to decrease. This will cause the value of $\log_{10} |\kappa - 1|$ to decrease as the resolution increases. Additionally, an increase in the resolution will decrease the range of ϵ over which the discretization errors dominate. Therefore, we expect the plot of κ vs ϵ to be of the general form shown in the graph below.

The following κ test graphs were generated using a fixed spacial resolution of 128 and with varying temporal resolutions $\Delta t \in [2^{-4}, 2^{-12}]$. The initial conditions used are the 3D Taylor Green, 3D ABC, and 3D ABC Mixture, each with a random initial perturbation. The simulations were run for timescales of $T = 1$ and $T = 3$. The results are shown below.

The results from the 3D Taylor-Green κ test show the expected behaviour from the theoretical kappa test graph. The value of kappa increases at very small and large values of epsilon. The value of kappa approaches 1 as Δt decreases, and so the plateau region lowers and decreases in width. The $T=1$ time scale results follow very closely

to the expected results while the $T=3$ time scale simulations show deviations from the the expected behaviour. These differences are due to the random initial direction of each simulation having more time to deviate from one another.

The kappa test results for the 3D ABC initial condition exhibit the same lowering of kappa as the time step decreases. However, unlike the Taylor Green initial condition, the results from the 3D ABC κ test have little to no plateau region. This indicates that the 3D ABC initial condition simulations are much more sensitive to the round off and truncation errors.

The results from the 3D ABC Mixture κ test show the same expected behaviour as the 3D Taylor Green kappa test.

The three kappa test results align well with the expected theoretical result. Therefore, we conclude that the discretization and time stepping techniques used in this analysis are valid and accurate.

5 Results

Utilizing the methods described in Sections 2, 3, and 4, we present numerical evidence that a singularity does not form in the Euler flow from the Taylor Green Vortex initial condition. In addition, we present numerical evidence that upon optimization ... RESULTS!

5.1 Singularity Detection

Theorem 2.1.3 provides the criteria we use to determine if a singularity forms in the Euler flow corresponding to initial condition $\boldsymbol{\eta}$, where $\boldsymbol{\eta}$ is the optimal initial condition found for the smallest value of α . First we check that the initial conditions $\boldsymbol{\eta}_\alpha$ of Euler-Voigt converge to this optimal initial condition $\boldsymbol{\eta}$ and so satisfy (2.8). If this criterion is satisfied, we then check that (2.12) is satisfied and $\alpha^2\Phi_{\alpha,T}$ converges to a positive value as $\alpha \rightarrow 0$.

Simply measuring the values of (2.8) and (2.12) as α decreases will not be sufficient for determining if each condition is satisfied. This is because it is not necessarily clear whether or not the value is converging to zero or to some small positive value. In order to make this determination we use the method of [17], which proposes the

following ansatz,

$$\sup_{t \in [0, T^*]} \|\nabla \mathbf{u}_\alpha(t)\|_{L^2} \approx \mathcal{O}(\alpha^p), \quad (5.62)$$

for $T^* > 0$ sufficiently large and for some power p . From this and Theorem 2.1.3, in order for a singularity to form in the Euler flow, it must be the case that the increase in the objective functional is larger than the diminishing of α . Therefore, for a singularity to form, $p \leq -1$. To find the value of p , we create a log-log plot of $\max_{t \in [0, T^*]} \|\nabla \mathbf{u}_\alpha(t)\|_{L^2}$ vs α . In order to simplify the visualization of the evolution of this quantity over time, the figures included here will only include the results at $T = 0, 1, \dots, T^*$. The value of p is defined as the steepest slope between any two values of α at any point in the time window.

We utilize the same method to determine if (2.8) is satisfied. However, in this case the condition is satisfied if the quantity does converge to zero. Therefore, we propose the following two ansatze.

$$\|\boldsymbol{\eta} - \boldsymbol{\eta}_\alpha\|_{L^2} \approx \mathcal{O}(\alpha^q), \quad (5.63)$$

and

$$\|\nabla(\boldsymbol{\eta} - \boldsymbol{\eta}_\alpha)\|_{L^2} \approx \mathcal{O}(\alpha^w). \quad (5.64)$$

By creating log-log plots for (5.63) and (5.64) vs α , the values of q and w are given by the steepest slope between any two values of α at any point in time for (5.63) and (5.64) respectively. For (2.8) to be satisfied, it must be the case that $q > 0$ and $w > -1$.

5.2 Initial Guess Results

We begin by presenting our results of for the initial guess of the TGV without optimization. A visualization of the vorticity of the TGV is shown in Figure 2.2. Figure 5.1 shows the vorticity of the solution at $T = 3$ for $\alpha = 8/1024$ and Figure 5.2 shows the vorticity of the solution at $T = 5$. The vorticity field in these figures are shifted along the x axis to bring together areas of significance that were split along the periodic boundaries. The figures show that the areas of high vorticity are concentrated into smaller area as the flow evolves. They also show that the peak of the vorticity increases in magnitude.




Figure 5.1: Isosurfaces of the vorticity at $T=5$ from the TGV. (a) magnitude of ω , vorticity components (b) ω_1 , (c) ω_2 , and (d) ω_3 .

Without optimization, the initial conditions for all values of α are simply the TGV. In this case, it is trivially true that condition (2.8) is satisfied, and so we only need




Figure 5.2: Isosurfaces of the vorticity at $T=5$ from the TGV. (a) magnitude of ω , vorticity components (b) ω_1 , (c) ω_2 , and (d) ω_3 .




Figure 5.3: ~~Nonoptimized TGV~~ with $T = 3$. Log-log plot of $\max_{t \in [0, T']} \|\nabla \mathbf{u}_\alpha(t)\|_{L^2}$ vs α at $T' = 0$ (blue) to $T' = 3$ (red), Steepest slope $p = -0.225$ (black), slope of $p = -1$ (dotted black)




Figure 5.4: Nonoptimized TGV with $T = 3$. Plot of the slope of $(\max_{t \in [0, T']} \|\nabla \mathbf{u}_\alpha(t)\|_{L^2} \text{ vs } \alpha)$ vs time at each α .

to compute the value of p to determine if a singularity forms in the Euler flow corresponding to the TGV.

Figures 5.3 and 5.4 show that without optimization the steepest slope achieved at any point in the time interval is $p = -0.225$ and this slope occurs at $T = 3$. Therefore, these results indicate that a singularity does not form in Euler within the time interval $[0, 3]$ from the TGV initial condition. They also show that the slope is still decreasing at the end of the time window, indicating that they may reach a slope of $p < -1$ at some later point in time. This result is consistent with the results of [3, 17] as the time interval is too short.

Figures 5.5 and 5.6 show that without optimization the steepest slope achieved at any point in the time is $p = -0.417$ and that this slope occurs at $T = 5$. This slope is larger than the p value for the shorter time window. However, it is still not close to the $p = -1$ required by (5.62). Therefore, these results indicate that a singularity does not form in Euler within the time interval $[0, 5]$ from the TGV initial condition. This result is unexpected because the interval is longer than the blow-up time of




Figure 5.5: Nonoptimized TGV with $T = 3$. Log-log plot of $\max_{t \in [0, T']} \|\nabla \mathbf{u}_\alpha(t)\|_{L^2}$ vs α at $T' = 0$ (blue) to $T' = 5$ (red), Steepest slope $p = -0.417$ (black), slope of $p = -1$ (dotted black)




Figure 5.6: Nonoptimized TGV with $T = 5$. Plot of the slope of $(\max_{t \in [0, T']} \|\nabla \mathbf{u}_\alpha(t)\|_{L^2} \text{ vs } \alpha)$ vs time at each α .

$T^* = 4$ from [3].

5.2.1 Comparison to Larios et al. 2018

Our results for the formation of a singularity in the Euler flow corresponding to the TGV initial condition conflict with those of [17, 3]. In [17] their analysis also examined the Euler-Voigt equations in order to search for singularity formation in Euler from the TGV. As shown in Figure 3 of [17], they obtained a result of $p \ll -1$, which indicated that blow-up does occur in Euler from the TGV. Additionally, the minimum of p occurred at a time consistent with the results of [3]. However, their model used a 3D unit torus ($L = 1$) and a coefficient of $V_0 = 1$, rather than the $V_0 = 1/2\pi$ we use to match the scaling of [3]. Because of this the time scale of their result does not match the time scale of [3]. Therefore, the results of [17] do indicate a singularity forms in Euler from the TGV initial condition. However, the blow-up time is much longer than the $T^* \approx 4$ of [3] which is consistent with our result.

6 Conclusion

Every thesis also needs a concluding chapter

A Function Spaces

In this Appendix we introduce the Lebesgue space, ~~continuous function space~~, and Sobolev space. These function spaces are used to define the initial conditions and solutions of the Euler and Euler-Voigt equations, the criteria for singularity formation, the objective functional, the gradient of the objective functional, and the Riemannian conjugate gradient method.

The Lebesgue space $L^p(\Omega)$ consists of real-valued measurable functions $\mathbf{u}$, defined on Ω , and $p \in \mathbb{R}^+$ for which the norm is bounded. The L^p norm is defined as

$$\|\mathbf{u}\|_{L^p} = \left(\int_{\Omega} |\mathbf{u}|^p d\mathbf{x} \right)^{1/p},$$

where $1 \leq p < \infty$. ~~In Fourier space the norm is defined as~~

$$\|\mathbf{u}\|_{L^p} = \left(\sum_{\mathbf{k} \in \mathbb{Z}^3} |\widehat{\mathbf{u}}_{\mathbf{k}}|^p \right)^{1/p},$$

where $\widehat{\mathbf{u}}_{\mathbf{k}}$ is the Fourier coefficient of $\mathbf{u}$, as defined in (2.23). For $p = \infty$, the norm is given by

$$\|\mathbf{u}\|_{L^\infty} = \operatorname{ess\,sup}_{\mathbf{x} \in \Omega} |\mathbf{u}|,$$

where ess sup is the essential supremum of $|\mathbf{u}|$ on Ω [12, 26].

Let $\alpha = (\alpha_1, \dots, \alpha_n)$ be an n -tuple of nonnegative integers α_i . For any nonnegative integer k , the space $C^k(\Omega)$ is the space of functions $\mathbf{u}$ which, together with all partial derivatives $\partial^\alpha \mathbf{u}$ of orders $|\alpha| \leq k$, are continuous on Ω [12].

For $s \in \mathbb{R}^+$, the L^2 -based Sobolev space $H^s(\Omega)$ is the space of all functions for which the norm is finite [26]. The norm is defined as

$$\|\mathbf{u}\|_{H^s}^2 := (2\pi)^3 \sum_{k \in \mathbb{Z}^3} (1 + |k|^{2s}) |\widehat{\mathbf{u}}_k|^2.$$

In physical space, the norm is defined as

$$\|\mathbf{u}\|_{H^s}^2 = \sum_{0 \leq |\alpha| \leq s} \|\partial^\alpha \mathbf{u}\|^2.$$

For any function $\mathbf{u} \in H^s(\mathbb{T}^3)$ the $\dot{H}^s$ seminorm is defined by

$$\|\mathbf{u}\|_{\dot{H}^s}^2 := (2\pi)^3 \sum_{k \in \mathbb{Z}^3} |k|^{2s} |\widehat{\mathbf{u}}_k|^2.$$

The seminorm in physical space is defined as

$$\|\mathbf{u}\|_{\dot{H}^s}^2 = \sum_{|\alpha|=k} \int_{\Omega} |\partial^\alpha \mathbf{u}|^2 d\mathbf{x}.$$

B Proof of Relation (2.25)

In this Appendix we prove (2.25) which shows that the parameter σ determines the relative 'size' of the Gevrey space in which we search for optimal initial conditions.

Proof. Let $\sigma_1, \sigma_2 \in \mathbb{R}$ be defined such that $0 < \sigma_1 < \sigma_2$. Therefore, from (2.22)

$$\begin{aligned}
 \|\mathbf{v}\|_{G^{\sigma_2}} &= \left[\int_{\mathbb{T}^3} (1 + |D|^2)^s e^{2\sigma_2|D|} \mathbf{v} \cdot \mathbf{v} \, d\mathbf{x} \right]^{1/2} \\
 &\geq \left[\int_{\mathbb{T}^3} (1 + |D|^2)^s e^{2\sigma_1|D|} \mathbf{v} \cdot \mathbf{v} \, d\mathbf{x} \right]^{1/2} = \|\mathbf{v}\|_{G^{\sigma_1}} \\
 &> \left[\int_{\mathbb{T}^3} (1 + |D|^2)^s e^0 \mathbf{v} \cdot \mathbf{v} \, d\mathbf{x} \right]^{1/2} = \|\mathbf{v}\|_{G^0} \\
 &= \left[\int_{\mathbb{T}^3} (1 + |D|^2)^s \mathbf{v} \cdot \mathbf{v} \, d\mathbf{x} \right]^{1/2} = \|\mathbf{v}\|_{H^s}.
 \end{aligned}$$

Thus,

$$\|\mathbf{v}\|_{G^{\sigma_2}} > \|\mathbf{v}\|_{G^{\sigma_1}} > \|\mathbf{v}\|_{G^0} = \|\mathbf{v}\|_{H^s}.$$

It follows from this that if $\|\mathbf{v}\|_{G^{\sigma_2}} < \infty$ then $\|\mathbf{v}\|_{G^{\sigma_1}} < \infty$ and so if $v \in G^{\sigma_2}$ then $v \in G^{\sigma_1}$. However, the converse is not true. The same argument shows that if $v \in G^{\sigma_1}$ then $v \in G^0$, which is equivalent to $v \in H^s$. Therefore,

$$G^{\sigma_2} \subset G^{\sigma_1} \subset G^0 = H^s.$$

□

C Rescaling the Euler equations

~~In this Appendix we derive the criteria that must be satisfied by a rescaled velocity field to be valid solution of the Euler equations.~~

Given a solution $\mathbf{u}$ of the Euler equations (1.2), we construct a new scaled solution with

$$\tau = \lambda t, \quad \text{and} \quad \mathbf{y} = \beta \mathbf{x}.$$

where $\lambda, \beta > 0$ are constants. Our new solution $\mathbf{v}$ is defined as

$$\mathbf{v} = \mathbf{v}(\mathbf{y}, \tau) = \gamma \mathbf{u}(\mathbf{y}/\beta, \tau/\lambda).$$

Therefore

$$\partial_t \mathbf{u} = \frac{1}{\gamma} \frac{\partial \mathbf{v}}{\partial \tau} \frac{d\tau}{dt} = \frac{\lambda}{\gamma} \partial_\tau \mathbf{v}.$$

and

$$\nabla \mathbf{u} = \frac{1}{\gamma} \nabla \mathbf{v} \frac{d\mathbf{y}}{d\mathbf{x}} = \frac{\beta}{\gamma} \nabla \mathbf{v}.$$

Substituting $\mathbf{v}$ into equation (1.2a)

$$\frac{\lambda}{\gamma} \partial_\tau \mathbf{v} + \frac{\beta}{\gamma^2} (\mathbf{v} \cdot \nabla) \mathbf{v} + \nabla q = 0,$$

where q is defined as whatever pressure is required to satisfy the incompressibility condition. This can be rearranged to the form,

$$\frac{\lambda\gamma}{\beta}\partial_\tau\mathbf{v} + (\mathbf{v} \cdot \nabla)\mathbf{v} + \nabla q' = 0.$$

Therefore, our function $\mathbf{v}$ can only be a solution to the Euler equations if

$$\frac{\lambda\gamma}{\beta} = 1.$$

For the Taylor Green Vortex with $L = 2\pi$ and $V_0 = 1$, let

$$\gamma = \lambda = \beta = 1.$$

In order to rescale this to the domain with $L = 1$ we must set $\beta = 1/2\pi$. Additionally, we want our rescaled solution to have the same time scale as the original solution and so we must set $\lambda = 1$. Therefore, for our scaled solution $\mathbf{v}$ to be a solution of the Euler equations we must set $\gamma = V_0 = 1/2\pi$.

D Rescaling the Euler-Voigt equations

In this Appendix ~~we derive the criteria that must be satisfied by a rescaled velocity field to be valid solution of the Euler-Voigt equations.~~

Given a solution $\mathbf{u}_\alpha$ of the Euler-Voigt equations (1.4), we construct a new scaled solution with

$$\tau = \lambda t, \quad \kappa = \beta \alpha, \quad \mathbf{y} = \beta \mathbf{x}.$$

where $\lambda, \beta > 0$ are constants. The variables $\mathbf{x}$ and α are both scaled by β because they are both spatial and we want to keep the relative size of the filter to the domain size fixed. Our new solution $\mathbf{v}$ is defined as

$$\mathbf{v} = \mathbf{v}(\mathbf{y}, \tau) = \gamma \mathbf{u}_\alpha(\mathbf{y}/\beta, \tau/\lambda).$$

Therefore

$$\partial_t \mathbf{u}_\alpha = \frac{1}{\gamma} \frac{\partial \mathbf{v}}{\partial \tau} \frac{d\tau}{dt} = \frac{\lambda}{\gamma} \partial_\tau \mathbf{v}.$$

and

$$\nabla \mathbf{u}_\alpha = \frac{1}{\gamma} \nabla \mathbf{v} \frac{d\mathbf{y}}{d\mathbf{x}} = \frac{\beta}{\gamma} \nabla \mathbf{v}.$$

and

$$\partial_t \Delta \mathbf{u}_\alpha = \partial_t \frac{\beta^2}{\gamma} \Delta \mathbf{v} = \frac{\lambda \beta^2}{\gamma} \partial_\tau \Delta \mathbf{v}.$$

Substituting $\mathbf{v}$ into equation (1.4a)

$$-\alpha^2 \frac{\lambda \beta^2}{\gamma} \partial_\tau \Delta \mathbf{v} + \frac{\lambda}{\gamma} \partial_\tau \mathbf{v} + \frac{\beta}{\gamma^2} (\mathbf{v} \cdot \nabla) \mathbf{v} + \nabla q = 0,$$

where q is defined as whatever pressure is required to satisfy the incompressibility condition. This can be rearranged to the form,

$$-\kappa^2 \partial_t \Delta \mathbf{v} + \partial_\tau \mathbf{v} + \frac{\beta}{\lambda \gamma} (\mathbf{v} \cdot \nabla) \mathbf{v} + \nabla q' = 0.$$

Therefore, our function $\mathbf{v}$ can only be a solution to the Euler-Voigt equations if

$$\frac{\beta}{\lambda \gamma} = 1.$$

For the Taylor Green Vortex with $L = 2\pi$ and $V_0 = 1$, let $\gamma = \lambda = \beta = 1$. In order to rescale this to the domain with $L = 1$ we must set $\beta = 1/2\pi$. Additionally, we want our rescaled solution to have the same time scale as the original solution and so we must set $\lambda = 1$. Therefore, for our scaled solution $\mathbf{v}$ to be a solution of the Euler-Voigt equations we must set $\gamma = V_0 = 1/2\pi$.

E Evaluation of $\|\boldsymbol{\eta}\|_{\dot{H}^1}$ for the General Taylor Green Vortex

In this Appendix we derive the $\dot{H}^1$ seminorm for the General Taylor Green Vortex. The $\dot{H}^1$ seminorm is required for properly scaling the initial conditions of the Euler and Euler-Voigt equations.

The general form of the 3D Taylor Green Vortex on the 3D torus Ω with characteristic length L is given in (2.29). To simplify the notation we drop the subscript *TGV* and substitute in

$$x' = 2\pi x/L, \quad y' = 2\pi y/L, \quad z' = 2\pi z/L.$$

The formula for $\|\boldsymbol{\eta}\|_{\dot{H}^1}$ is

$$\|\boldsymbol{\eta}\|_{\dot{H}^1} = \|\nabla \boldsymbol{\eta}\|_{L^2} = \left[\sum_{i,j=1}^3 \int_{\Omega} (\partial_i \boldsymbol{\eta}_j)^2 d\mathbf{x} \right]^{1/2},$$

where ∂_i is the partial derivative with respect to the i -th component and $\boldsymbol{\eta}_j$ is the j -th component of $\boldsymbol{\eta}$. Because the 3rd component of $\boldsymbol{\eta}$ is zero, the partial derivative

is also zero. This leaves us with six integrals to evaluate.

$$\begin{aligned} \sum_{i,j=1}^3 \int_{\Omega} (\partial_i \boldsymbol{\eta}_j)^2 d\mathbf{x} &= \int_{\Omega} (\partial_1 \boldsymbol{\eta}_1)^2 d\mathbf{x} + \int_{\Omega} (\partial_2 \boldsymbol{\eta}_1)^2 d\mathbf{x} + \int_{\Omega} (\partial_3 \boldsymbol{\eta}_1)^2 d\mathbf{x} \\ &\quad + \int_{\Omega} (\partial_1 \boldsymbol{\eta}_2)^2 d\mathbf{x} + \int_{\Omega} (\partial_2 \boldsymbol{\eta}_2)^2 d\mathbf{x} + \int_{\Omega} (\partial_3 \boldsymbol{\eta}_2)^2 d\mathbf{x}. \end{aligned}$$

Evaluating each partial derivative

$$\begin{aligned} \sum_{i,j=1}^3 \int_{\Omega} (\partial_i \boldsymbol{\eta}_j)^2 d\mathbf{x} &= \int_{\Omega} (2\pi V_0/L \cos(x') \cos(y') \cos(z'))^2 d\mathbf{x} \\ &\quad + \int_{\Omega} (-2\pi V_0/L \sin(x') \sin(y') \cos(z'))^2 d\mathbf{x} \\ &\quad + \int_{\Omega} (-2\pi V_0/L \sin(x') \cos(y') \sin(z'))^2 d\mathbf{x} \\ &\quad + \int_{\Omega} (2\pi V_0/L \sin(x') \sin(y') \cos(z'))^2 d\mathbf{x} \\ &\quad + \int_{\Omega} (-2\pi V_0/L \cos(x') \cos(y') \cos(z'))^2 d\mathbf{x} \\ &\quad + \int_{\Omega} (2\pi V_0/L \cos(x') \sin(y') \sin(z'))^2 d\mathbf{x}, \end{aligned}$$

which simplifies to

$$\begin{aligned} \sum_{i,j=1}^3 \int_{\Omega} (\partial_i \boldsymbol{\eta}_j)^2 d\mathbf{x} &= 4\pi^2 V_0^2/L^2 \int_{\Omega} \cos^2(x') \cos^2(y') \cos^2(z') d\mathbf{x} \\ &\quad + 4\pi^2 V_0^2/L^2 \int_{\Omega} \sin^2(x') \sin^2(y') \cos^2(z') d\mathbf{x} \\ &\quad + 4\pi^2 V_0^2/L^2 \int_{\Omega} \sin^2(x') \cos^2(y') \sin^2(z') d\mathbf{x} \\ &\quad + 4\pi^2 V_0^2/L^2 \int_{\Omega} \sin^2(x') \sin^2(y') \cos^2(z') d\mathbf{x} \\ &\quad + 4\pi^2 V_0^2/L^2 \int_{\Omega} \cos^2(x') \cos^2(y') \cos^2(z') d\mathbf{x} \\ &\quad + 4\pi^2 V_0^2/L^2 \int_{\Omega} \cos^2(x') \sin^2(y') \sin^2(z') d\mathbf{x}. \end{aligned}$$

Because the integrand is separable, the integrals can be reduced to products of,

$$\int_0^L \cos^2(x') dx = \frac{L}{2}, \quad \text{and} \quad \int_0^L \sin^2(x') dx = \frac{L}{2}.$$

Therefore, each of the six integrals is equal to

$$\frac{4\pi^2 V_0^2}{L^2} \frac{L^3}{8}.$$

The sum of the six integrals is

$$\sum_{i,j=1}^3 \int_{\Omega} (\partial_i \boldsymbol{\eta}_j)^2 d\mathbf{x} = 6 \frac{4\pi^2 V_0^2}{L^2} \frac{L^3}{8} = 3\pi^2 V_0^2 L.$$

Substituting into the formula for $\|\cdot\|_{\dot{H}^1}$ gives

$$\|\boldsymbol{\eta}_{TGV}\|_{\dot{H}^1} = [3\pi^2 V_0^2 L]^{1/2} = \sqrt{3L}\pi V_0.$$

F Evaluation of $\|\nabla \times \boldsymbol{\eta}\|_{L^\infty}$ for the General Taylor Green Vortex

In this Appendix we ~~derive~~ the L^∞ norm of the vorticity for the General Taylor Green Vortex. The L^∞ norm of the vorticity is required for properly scaling the initial conditions of the Euler and Euler-Voigt equations.

The general form of the 3D Taylor Green Vortex on the 3D torus Ω with characteristic length L is given in (2.29). To simplify the notation we drop the subscript *TGV* and substitute in

$$x' = 2\pi x/L, \quad y' = 2\pi y/L, \quad z' = 2\pi z/L.$$

The vorticity of $\boldsymbol{\eta}$ is

$$\nabla \times \boldsymbol{\eta} = \begin{bmatrix} \partial_2 \eta_3 - \partial_3 \eta_2 \\ \partial_3 \eta_1 - \partial_1 \eta_3 \\ \partial_1 \eta_2 - \partial_2 \eta_1 \end{bmatrix}$$

where ∂_i is the partial derivative with respect to the i -th component and $\boldsymbol{\eta}_j$ is the j -th component of $\boldsymbol{\eta}$. Substituting in the partial derivatives

$$\nabla \times \boldsymbol{\eta} = \begin{bmatrix} 0 - (-(2\pi V_0/L) \cos(x') \sin(y') \sin(z')) \\ -(2\pi V_0/L) \sin(x') \cos(y') \sin(z') - 0 \\ (2\pi V_0/L) \sin(x') \sin(y') \cos(z') + (2\pi V_0) \sin(x') \sin(y') \cos(z'), \end{bmatrix}$$

which simplifies to

$$\nabla \times \boldsymbol{\eta}_{TG} = \begin{bmatrix} (2\pi V_0/L) \cos(x') \sin(y') \sin(z') \\ -(2\pi V_0/L) \sin(x') \cos(y') \sin(z') \\ (4\pi V_0/L) \sin(x') \sin(y') \cos(z') \end{bmatrix}.$$

The $\|\nabla \times \boldsymbol{\eta}\|_{L^\infty}$ norm is the maximum absolute value of the components of $\nabla \times \boldsymbol{\eta}$.

$$\begin{aligned} \|\nabla \times \boldsymbol{\eta}_{TG}\|_{L^\infty} &= \max(|(2\pi V_0/L) \cos(x') \sin(y') \sin(z')| \\ &\quad , |-(2\pi V_0/L) \sin(x') \cos(y') \sin(z')| \\ &\quad , |(4\pi V_0/L) \sin(x') \sin(y') \cos(z')|) \end{aligned}$$

The maximums are

$$\|\nabla \times \boldsymbol{\eta}_{TG}\|_{L^\infty} = \max((2\pi V_0/L), (2\pi V_0/L), (4\pi V_0/L))$$

Therefore,

$$\|\nabla \times \boldsymbol{\eta}_{TG}\|_{L^\infty} = 4\pi V_0/L.$$

G Linearization of the Euler-Voigt System

In this Appendix we derive the linearization of the Euler-Voigt equations. ~~The linearization of the Euler-Voigt equations~~ is used to construct the gradient of the Objective functional.

The Euler-Voigt equations are defined as,

$$\begin{cases} -\alpha^2 \partial_t \Delta \mathbf{u}_\alpha + \partial_t \mathbf{u}_\alpha + (\mathbf{u}_\alpha \cdot \nabla) \mathbf{u}_\alpha + \nabla p = 0 \\ \nabla \cdot \mathbf{u}_\alpha = 0 \\ \mathbf{u}_\alpha|_{t=0} = \boldsymbol{\eta}_\alpha. \end{cases}$$

Substituting in $\mathbf{u}_\alpha = \mathbf{u}_\alpha + \mathbf{u}'_\alpha$, $\boldsymbol{\eta}_\alpha = \boldsymbol{\eta}_\alpha + \boldsymbol{\eta}'_\alpha$, and $p = p + p'$,

$$\begin{cases} -\alpha^2 \partial_t \Delta (\mathbf{u}_\alpha + \mathbf{u}'_\alpha) + \partial_t (\mathbf{u}_\alpha + \mathbf{u}'_\alpha) + ((\mathbf{u}_\alpha + \mathbf{u}'_\alpha) \cdot \nabla) (\mathbf{u}_\alpha + \mathbf{u}'_\alpha) + \nabla (p + p') = 0 \\ \nabla \cdot (\mathbf{u}_\alpha + \mathbf{u}'_\alpha) = 0 \\ (\mathbf{u}_\alpha + \mathbf{u}'_\alpha)|_{t=0} = \boldsymbol{\eta}_\alpha + \boldsymbol{\eta}'_\alpha. \end{cases}$$

By expanding each term we obtain,

$$\begin{cases} -\alpha^2 \partial_t \Delta \mathbf{u}_\alpha - \alpha^2 \partial_t \Delta \mathbf{u}'_\alpha + \partial_t \mathbf{u}_\alpha + \partial_t \mathbf{u}'_\alpha + \mathbf{u}_\alpha \cdot \nabla \mathbf{u}_\alpha + \\ \mathbf{u}_\alpha \cdot \nabla \mathbf{u}'_\alpha + \mathbf{u}'_\alpha \cdot \nabla \mathbf{u}_\alpha + \mathbf{u}'_\alpha \cdot \nabla \mathbf{u}'_\alpha + \nabla p + \nabla p' = 0 \\ \nabla \cdot (\mathbf{u}_\alpha + \mathbf{u}'_\alpha) = 0 \\ (\mathbf{u}_\alpha + \mathbf{u}'_\alpha)|_{t=0} = \boldsymbol{\eta}_\alpha + \boldsymbol{\eta}'_\alpha. \end{cases}$$

Substituting in the Euler-Voigt equations we simplify to

$$\begin{cases} -\alpha^2 \partial_t \Delta \mathbf{u}'_\alpha + \partial_t \mathbf{u}'_\alpha + \mathbf{u}_\alpha \cdot \nabla \mathbf{u}'_\alpha + \mathbf{u}'_\alpha \cdot \nabla \mathbf{u}_\alpha + \mathbf{u}'_\alpha \cdot \nabla \mathbf{u}'_\alpha + \nabla p' = 0 \\ \nabla \cdot \mathbf{u}'_\alpha = 0 \\ \mathbf{u}'_\alpha|_{t=0} = \boldsymbol{\eta}'_\alpha. \end{cases}$$

Linearizing the system by removing non-linear terms, we derive the linearized Euler-Voigt system

$$\begin{cases} -\alpha^2 \partial_t \Delta \mathbf{u}'_\alpha + \partial_t \mathbf{u}'_\alpha + \mathbf{u}_\alpha \cdot \nabla \mathbf{u}'_\alpha + \mathbf{u}'_\alpha \cdot \nabla \mathbf{u}_\alpha + \nabla p' = 0 \\ \nabla \cdot \mathbf{u}'_\alpha = 0 \\ \mathbf{u}'_\alpha|_{t=0} = \boldsymbol{\eta}'_\alpha. \end{cases}$$

This can be rearranged to the form

$$\begin{cases} \partial_t \mathbf{u}'_\alpha + (I - \alpha^2 \Delta)^{-1} (\mathbf{u}_\alpha \cdot \nabla \mathbf{u}'_\alpha + \mathbf{u}'_\alpha \cdot \nabla \mathbf{u}_\alpha + \nabla p') = 0 \\ \nabla \cdot \mathbf{u}'_\alpha = 0 \\ \mathbf{u}'_\alpha|_{t=0} = \boldsymbol{\eta}'_\alpha. \end{cases}$$

H Derivation of the Adjoint System

In this Appendix we derive the adjoint system (3.38) by ~~integrating the duality-pairing relation (3.39) in space and time.~~

For this derivation we will exclude the α subscript on the u_α , u'_α , and u_α^* terms for the sake of simplicity. Additionally, we substitute the integration with the following notation

$$\int_0^T \int_{\mathbb{T}^3} u \, dt = \int_D u \, dt$$

We derive the adjoint system using the following duality pairing relation,

$$(L[u'], u^*) := \int_D L[u'] u^* \, dx \, dt = (u', L[u^*]) + \int_a^b u' u^*|_{t=T} - \int_a^b u' u^*|_{t=0} \, dx = 0.$$

Substituting in the equation from the Euler-Voigt perturbation system,

$$(L[u'], u^*) := \int_D (\partial_t u' + (I - \alpha^2 \Delta)^{-1} (u \cdot \nabla u' + u' \cdot \nabla u + \nabla p')) u^* + (\nabla \cdot u') p^* \, dx \, dt.$$

This integral can be split into five parts. Solving the first part with integration by parts, with $u = u^*$ and $v' = \partial_t u'$

$$\int_D u^* \partial_t u' \, dx \, dt = \int_{\mathbb{T}^3} u^* u'|_{t=0}^{t=T} \, dx - \int_D u' \partial_t u^* \, dx \, dt.$$

Starting with the second part and expressing it in Einstein Summation notation,

$$[(I - \alpha^2 \Delta)^{-1} u \cdot \nabla u'] \cdot u^* dx dt = \left[(I - \alpha^2 \Delta)^{-1} u_j \frac{\partial u'_i}{\partial x_j} \right] u_i^* dx dt.$$

Using the derivative product rule and the fact that $\nabla \cdot u = 0$,

$$\frac{\partial}{\partial x_j} (u_j u'_i) = \frac{\partial u_j}{\partial x_j} u'_i + u_j \frac{\partial u'_i}{\partial x_j} = u_j \frac{\partial u'_i}{\partial x_j}.$$

Substituting this into the integral,

$$[(I - \alpha^2 \Delta)^{-1} u \cdot \nabla u'] \cdot u^* dx dt = \left[(I - \alpha^2 \Delta)^{-1} \frac{\partial}{\partial x_j} (u_j u'_i) \right] u_i^* dx dt.$$

Using the periodic boundaries of the system, we can now apply integration by parts with $u = [(I - \alpha^2 \Delta)^{-1} u_i^*]$ and $v' = \frac{\partial}{\partial x_j} (u_j u'_i)$,

$$[(I - \alpha^2 \Delta)^{-1} u \cdot \nabla u'] \cdot u^* dx dt = -u_j u'_i \frac{\partial}{\partial x_j} [(I - \alpha^2 \Delta)^{-1} u_i^*] dx dt.$$

Converting back to vector notation,

$$[(I - \alpha^2 \Delta)^{-1} u \cdot \nabla u'] \cdot u^* dx dt = -u' \cdot [u \cdot \nabla [(I - \alpha^2 \Delta)^{-1} u^*]] dx dt.$$

Starting with the third part and expressing it in Einstein Summation notation,

$$[(I - \alpha^2 \Delta)^{-1} u' \cdot \nabla u] \cdot u^* dx dt = \left[(I - \alpha^2 \Delta)^{-1} u'_j \frac{\partial u_i}{\partial x_j} \right] u_i^* dx dt.$$

Applying integration by parts with $u = [(I - \alpha^2 \Delta)^{-1} u_i^*]$ and $v' = \frac{\partial}{\partial x_j} (u_j' u_i)$

$$[(I - \alpha^2 \Delta)^{-1} u' \cdot \nabla u] \cdot u^* dx dt = -u_j' u_i \frac{\partial}{\partial x_j} [(I - \alpha^2 \Delta)^{-1} u_i^*] dx dt.$$

Because the goal is to extract u' and $v \cdot w = v^T \cdot w^T$, we can apply a transpose to the above equation and so swap the indices to,

$$[(I - \alpha^2 \Delta)^{-1} u' \cdot \nabla u] \cdot u^* dx dt = -u_i' u_j \frac{\partial}{\partial x_i} [(I - \alpha^2 \Delta)^{-1} u_j^*] dx dt.$$

Converting back to vector notation,

$$[(I - \alpha^2 \Delta)^{-1} u' \cdot \nabla u] \cdot u^* dx dt = -u' \cdot \left[u \cdot [\nabla (I - \alpha^2 \Delta)^{-1} u^*]^T \right] dx dt.$$

Starting with the fourth part of the integral and expressing it in Einstein Summation notation,

$$\nabla p' \cdot u^* dx dt = \frac{\partial p'}{\partial x_i} u_i^* dx dt.$$

Applying integration by parts with $u = u_i^*$ and $v' = \partial x_i p'$,

$$\nabla p' \cdot u^* dx dt = -p' \frac{\partial u_i^*}{\partial x_i} dx dt.$$

Converting back to vector notation,

$$\nabla p' \cdot u^* dx dt = -p' (\nabla \cdot u^*) dx dt.$$

Starting with the final part of the integral and expressing it in Einstein Summation

notation,

$$(\nabla \cdot u') p^* dx dt = \frac{\partial u'_i}{\partial x_i} p^* dx dt.$$

Applying integration by parts with $u = p^*$ and $v' = \partial x_i u'_i$,

$$(\nabla \cdot u') p^* dx dt = -\frac{\partial p^*}{\partial x_i} u'_i dx dt.$$

Converting back to vector notation,

$$(\nabla \cdot u') p^* dx dt = -u' \cdot \nabla p^* dx dt.$$

Combining the integrals,

$$\int_{\mathbb{T}^3} u^* u' \Big|_{t=0}^{t=T} dx - u' \left[\partial_t u^* - [u \cdot \nabla [(I - \alpha^2 \Delta)^{-1} u^*]] - [u \cdot [\nabla (I - \alpha^2 \Delta)^{-1} u^*]^T] - \nabla p^* \right] - p'(\nabla \cdot u^*)$$

This gives us our adjoint system,

$$\mathbb{L}^* \begin{bmatrix} u_\alpha^* \\ p_\alpha^* \end{bmatrix} := \begin{bmatrix} -\partial_t u^* - (\nabla(I - \alpha^2)^{-1} u_\alpha^* + (\nabla(I - \alpha^2)^{-1} u_\alpha^*)^T) u_\alpha - \nabla p_\alpha^* \\ -\nabla \cdot u_\alpha^* \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

To derive the terminal condition of the adjoint system we compare the second term of the duality pairing to the objective functional.

$$\begin{aligned} \int_{\mathbb{T}^3} u'(x, T) \mathbf{u}^*(x, T) dx &= 2 \langle u(x, T), u'(x, T) \rangle_{\dot{H}^1} \\ &= 2 \int_{\mathbb{T}^3} |D| u(x, T) \mathbf{u}'(x, T) dx \end{aligned}$$

$$= \int_{\mathbb{T}^3} 2|D|^2 u(x, T) \mathbf{u}'(x, T) dx$$

Therefore,

$$\mathbf{u}^*(x, T) = 2|D|^2 u(x, T)$$

I Derivation of the Gâteaux Directional Differential of the Objective Functional

In this Appendix we derive the Gâteaux directional differential of the objective functional, $\Phi'_{\alpha,T}(\boldsymbol{\eta}_\alpha; \boldsymbol{\eta}'_\alpha)$ in terms of the $\dot{H}^1$ inner product and L^2 inner product. This differential is used to derive the gradient of the objective functional.

We begin by substituting the second definition of (3.32) into (3.34) to obtain



$$\Phi'_{\alpha,T}(\boldsymbol{\eta}_\alpha; \boldsymbol{\eta}'_\alpha) = \lim_{\epsilon \rightarrow 0} \frac{1}{\epsilon} [\|\mathbf{u}_\alpha(T; \boldsymbol{\eta}_\alpha + \epsilon \boldsymbol{\eta}'_\alpha)\|_{\dot{H}^1}^2 - \|\mathbf{u}_\alpha(T; \boldsymbol{\eta}_\alpha)\|_{\dot{H}^1}^2].$$

Using the $\dot{H}^1$ seminorm as defined in Appedix A, and taking the limit $\epsilon \rightarrow 0$, we can express $\Phi'_{\alpha,T}$ as

$$\Phi'_{\alpha,T}(\boldsymbol{\eta}_\alpha; \boldsymbol{\eta}'_\alpha) = 2(2\pi)^3 \sum_{\mathbf{k} \in \mathbb{Z}^3} |\mathbf{k}|^2 |\hat{\mathbf{u}}_{\mathbf{k}} \hat{\mathbf{u}}'_{\mathbf{k}}|.$$

This result can also be expressed in the form of the $\dot{H}^1$ inner product

$$\Phi'_{\alpha,T}(\boldsymbol{\eta}_\alpha; \boldsymbol{\eta}'_\alpha) = 2 \langle \mathbf{u}_\alpha(\boldsymbol{x}, T), \mathbf{u}'_\alpha(\boldsymbol{x}, T) \rangle_{\dot{H}^1}.$$

Using the integral form of the $\dot{H}^s$ seminorm from Appendix A, the Gâteaux differential can be expressed in physical space as

$$\Phi'_{\alpha,T}(\boldsymbol{\eta}_\alpha; \boldsymbol{\eta}'_\alpha) = \int_{\mathbb{T}^3} 2|D|^2 \mathbf{u}_\alpha(\mathbf{x}, T) \cdot \mathbf{u}'_\alpha(\mathbf{x}, T) d\mathbf{x}.$$

We now make use of the ~~adjoint-system terminal condition~~ imposed in Appendix H

$$\mathbf{u}_\alpha^*(x, T) = 2|D|^2 \mathbf{u}_\alpha(\mathbf{x}, T).$$

Therefore,

$$\Phi'_{\alpha,T}(\boldsymbol{\eta}_\alpha; \boldsymbol{\eta}'_\alpha) = \int_{\mathbb{T}^3} \mathbf{u}_\alpha^*(\mathbf{x}, T) \cdot \mathbf{u}'_\alpha(\mathbf{x}, T) d\mathbf{x}.$$

Now using the duality-pairing **defined** in Appendix H

$$\Phi'_{\alpha,T}(\boldsymbol{\eta}_\alpha; \boldsymbol{\eta}'_\alpha) = \int_{\mathbb{T}^3} \mathbf{u}_\alpha^*(\mathbf{x}, 0) \cdot \boldsymbol{\eta}'_\alpha(\mathbf{x}) d\mathbf{x}.$$

~~Which~~ lets us express the value of $\Phi'_{\alpha,T}$ with the L^2 inner product

$$\Phi'_{\alpha,T}(\boldsymbol{\eta}_\alpha; \boldsymbol{\eta}'_\alpha) = \langle \mathbf{u}_\alpha^*(\mathbf{x}, 0), \boldsymbol{\eta}'_\alpha(\mathbf{x}) \rangle_{L^2}.$$

Therefore,

$$\Phi'_{\alpha,T}(\boldsymbol{\eta}_\alpha; \boldsymbol{\eta}'_\alpha) = 2 \langle \mathbf{u}_\alpha(\mathbf{x}, T), \mathbf{u}'_\alpha(\mathbf{x}, T) \rangle_{\dot{H}^1} = \langle \mathbf{u}_\alpha^*(\mathbf{x}, 0), \boldsymbol{\eta}'_\alpha(\mathbf{x}) \rangle_{L^2}.$$

Bibliography

- [1] J. T. Beale, T. Kato, and A. Majda. Remarks on the breakdown of smooth solutions for the 3-D Euler equations. *Communications in Mathematical Physics*, 94(1):61–66, March 1984.
- [2] Marc E. Brachet, Daniel I. Meiron, Steven A. Orszag, B. G. Nickel, Rudolf H. Morf, and Uriel Frisch. Small-scale structure of the Taylor–Green vortex. *Journal of Fluid Mechanics*, 130:411–452, May 1983.
- [3] Miguel D. Bustamante and Marc Brachet. Interplay between the Beale-Kato-Majda theorem and the analyticity-strip method to investigate numerically the incompressible Euler singularity problem. *Physical Review E*, 86(6):066302, December 2012. arXiv:1112.1571 [physics].
- [4] Claudio Canuto, M. Youssuff Hussaini, Alfio Quarteroni, and Thomas A. Zang. *Spectral Methods: Fundamentals in Single Domains*. Scientific Computation. Springer, Berlin, Heidelberg, 2006.
- [5] Yanping Cao, Evelyn M. Lunasin, and Edriss S. Titi. Global well-posedness of the three-dimensional viscous and inviscid simplified Bardina turbulence models. *Communications in Mathematical Sciences*, 4(4):823–848, December 2006.

-
- [6] Jiajie Chen and Thomas Y. Hou. Finite time blowup of 2D Boussinesq and 3D Euler equations with $C^{1,\alpha}$ velocity and boundary. *Communications in Mathematical Physics*, 383(3):1559–1667, May 2021. arXiv:1910.00173 [math.AP].
- [7] Jiajie Chen and Thomas Y. Hou. Stable nearly self-similar blowup of the 2D Boussinesq and 3D Euler equations with smooth data I: Analysis, May 2023. arXiv:2210.07191 [math.AP].
- [8] Hyuntae Cho, Yesom Park, and Myungjoo Kang. Solving incompressible Navier–Stokes equations on irregular domains and quadrees by monolithic approach. *Journal of Computational Physics*, 463:111304, August 2022.
- [9] Ionut Danaila and Bartosz Protas. Computation of Ground States of the Gross-Pitaevskii Functional via Riemannian Optimization. *SIAM Journal on Scientific Computing*, 39(6):B1102–B1129, January 2017. arXiv:1703.07693 [math].
- [10] Charles R. Doering. The 3D Navier-Stokes Problem. *Annual Review of Fluid Mechanics*, 41(Volume 41, 2009):109–128, January 2009.
- [11] Charles L Fefferman. Existence and Smoothness of the Navier-Stokes equation. <https://www.claymath.org/wp-content/uploads/2022/06/navierstokes.pdf>, 2022.
- [12] John J. F. Fournier and Robert A. Adams. *Sobolev Spaces*. 2 edition, June 2003.
- [13] J. D. Gibbon. The three-dimensional Euler equations: Where do we stand? *Physica D: Nonlinear Phenomena*, 237(14):1894–1904, August 2008.

- [14] Thomas Y. Hou. Potential Singularity of the 3D Euler Equations in the Interior Domain. *Foundations of Computational Mathematics*, 23(6):2203–2249, December 2023.
- [15] Thomas Y. Hou and De Huang. Potential singularity formation of incompressible axisymmetric Euler equations with degenerate viscosity coefficients, May 2022. arXiv:2102.06663 [math.AP].
- [16] Tosio Kato. Nonstationary flows of viscous and ideal fluids in R^3 . *Journal of Functional Analysis*, 9(3):296–305, March 1972.
- [17] Adam Larios, Mark R. Petersen, Edriss S. Titi, and Beth Wingate. A computational investigation of the finite-time blow-up of the 3D incompressible Euler equations based on the Voigt regularization. *Theoretical and Computational Fluid Dynamics*, 32(1):23–34, February 2018.
- [18] Adam Larios and Edriss S. Titi. On the Higher-Order Global Regularity of the Inviscid Voigt-Regularization of Three-Dimensional Hydrodynamic Models, February 2010. arXiv:0910.3354 [math].
- [19] Jean Leray. Sur le mouvement d’un liquide visqueux emplissant l’espace. *Acta Mathematica*, 63(1):193–248, December 1934.
- [20] David G. Luenberger. *Optimization by Vector Space Methods*. 1969.
- [21] Guo Luo and Thomas Y. Hou. Potentially Singular Solutions of the 3D Incompressible Euler Equations. *Proceedings of the National Academy of Sciences*, 111(36):12968–12973, September 2014. arXiv:1310.0497 [physics.flu-dyn].

-
- [22] H. K. Moffatt. Singularities in Fluid Mechanics, July 2019. arXiv:1907.06400 [physics].
- [23] Giuseppe Di Molfetta, Giorgio Krstulovic, and Marc Brachet. Self-truncation and scaling in Euler-Voigt-alpha and related fluid models. *Physical Review E*, 92(1):013020, July 2015. arXiv:1502.05544 [physics].
- [24] Jorge Nocedal and Stephen J. Wright. *Numerical Optimization*. Second edition edition, 2000.
- [25] Bartosz Protas. Systematic Search For Extreme and Singular Behavior in Some Fundamental Models of Fluid Mechanics. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 380(2225):20210035, June 2022. arXiv:2112.14917 [math].
- [26] James C. Robinson, Jose L. Rodrigo, and Witold Sadowski. *The Three-Dimensional Navier-Stokes Equations*. 2016.
- [27] Xinyu Zhao and Bartosz Protas. Systematic Search for Singularities in 3D Euler Flows. *Journal of Nonlinear Science*, 33(6):120, October 2023.