

Systematic Search for Singularities in 3D Euler-Voigt Flows

Noah Bensler¹, Bartosz Protas¹, Xinyu Zhao^{1,2}

¹Department of Mathematics & Statistics
McMaster University, Hamilton, Ontario, Canada

²Department of Mathematical Sciences, New Jersey Institute of Technology,
Newark, NJ 07103, USA

Funded by NSERC (Canada), Computing Time Provided by COMPUTE CANADA

April 23, 2026

Agenda

Introduction

- Singularities & Euler-Voigt Flows
- Finite-time Blow-up Criteria
- Initial Conditions

Optimization Problem for Euler-Voigt

- Formulation of the Optimization Problem
- Riemannian Conjugate Gradient Method
- Numerical Methods

Results

- Pre-Optimization Results
- Optimization Results
- Conclusion

Introduction

- Navier-Stokes system ($\Omega = [0, L]^3$)

$$\begin{cases} \partial_t \mathbf{u} + (\mathbf{u} \cdot \nabla) \mathbf{u} + \nabla p - \nu \Delta \mathbf{u} = \mathbf{0}, & \text{in } \Omega \times (0, T] \\ \nabla \cdot \mathbf{u} = 0, & \text{in } \Omega \times (0, T] \\ \mathbf{u} = \boldsymbol{\eta} & \text{in } \Omega \text{ at } t = 0 \\ \text{Periodic Boundary Conditions} & \text{on } \partial\Omega \times (0, T] \end{cases}$$

- The Big Question:

Given a smooth initial condition $\boldsymbol{\eta}$, does the Navier-Stokes system always admit smooth solutions $\mathbf{u}(t)$ for arbitrarily long times t ?

- One of the Clay Institute “Millennium Problems” (\$ 1M prize!)

<https://www.claymath.org/millennium/navier-stokes-equation/>

- Navier-Stokes system ($\Omega = [0, L]^3$)

$$\begin{cases} \partial_t \mathbf{u} + (\mathbf{u} \cdot \nabla) \mathbf{u} + \nabla p - \nu \Delta \mathbf{u} = \mathbf{0}, & \text{in } \Omega \times (0, T] \\ \nabla \cdot \mathbf{u} = 0, & \text{in } \Omega \times (0, T] \\ \mathbf{u} = \boldsymbol{\eta} & \text{in } \Omega \text{ at } t = 0 \\ \text{Periodic Boundary Conditions} & \text{on } \partial\Omega \times (0, T] \end{cases}$$

- **The Big Question:**

Given a smooth initial condition $\boldsymbol{\eta}$, does the Navier-Stokes system always admit smooth solutions $\mathbf{u}(t)$ for arbitrarily long times t ?

- One of the Clay Institute “Millennium Problems” (\$ 1M prize!)

<https://www.claymath.org/millennium/navier-stokes-equation/>

- Navier-Stokes system ($\Omega = [0, L]^3$)

$$\begin{cases} \partial_t \mathbf{u} + (\mathbf{u} \cdot \nabla) \mathbf{u} + \nabla p - \nu \Delta \mathbf{u} = \mathbf{0}, & \text{in } \Omega \times (0, T] \\ \nabla \cdot \mathbf{u} = 0, & \text{in } \Omega \times (0, T] \\ \mathbf{u} = \boldsymbol{\eta} & \text{in } \Omega \text{ at } t = 0 \\ \text{Periodic Boundary Conditions} & \text{on } \partial\Omega \times (0, T] \end{cases}$$

- **The Big Question:**

Given a smooth initial condition $\boldsymbol{\eta}$, does the Navier-Stokes system always admit smooth solutions $\mathbf{u}(t)$ for arbitrarily long times t ?

- One of the Clay Institute “Millennium Problems” (\$ 1M prize!)

<https://www.claymath.org/millennium/navier-stokes-equation/>

- Euler system ($\Omega = [0, L]^3$)

$$\begin{cases} \partial_t \mathbf{u} + (\mathbf{u} \cdot \nabla) \mathbf{u} + \nabla p = \mathbf{0}, & \text{in } \Omega \times (0, T] \\ \nabla \cdot \mathbf{u} = 0, & \text{in } \Omega \times (0, T] \\ \mathbf{u} = \boldsymbol{\eta} & \text{in } \Omega \text{ at } t = 0 \\ \text{Periodic Boundary Conditions} & \text{on } \partial\Omega \times (0, T] \end{cases}$$

- The Big Question:

Given a smooth initial condition $\boldsymbol{\eta}$, does the Euler system always admit smooth solutions $\mathbf{u}(t)$ for arbitrarily long times t ?

- There is numerical evidence for singularity formation in Euler (J.D. Gibbon 2008).

- Euler system ($\Omega = [0, L]^3$)

$$\begin{cases} \partial_t \mathbf{u} + (\mathbf{u} \cdot \nabla) \mathbf{u} + \nabla p = \mathbf{0}, & \text{in } \Omega \times (0, T] \\ \nabla \cdot \mathbf{u} = 0, & \text{in } \Omega \times (0, T] \\ \mathbf{u} = \boldsymbol{\eta} & \text{in } \Omega \text{ at } t = 0 \\ \text{Periodic Boundary Conditions} & \text{on } \partial\Omega \times (0, T] \end{cases}$$

- **The Big Question:**

Given a smooth initial condition $\boldsymbol{\eta}$, does the Euler system always admit smooth solutions $\mathbf{u}(t)$ for arbitrarily long times t ?

- There is numerical evidence for singularity formation in Euler (J.D. Gibbon 2008).

- Euler system ($\Omega = [0, L]^3$)

$$\begin{cases} \partial_t \mathbf{u} + (\mathbf{u} \cdot \nabla) \mathbf{u} + \nabla p = \mathbf{0}, & \text{in } \Omega \times (0, T] \\ \nabla \cdot \mathbf{u} = 0, & \text{in } \Omega \times (0, T] \\ \mathbf{u} = \boldsymbol{\eta} & \text{in } \Omega \text{ at } t = 0 \\ \text{Periodic Boundary Conditions} & \text{on } \partial\Omega \times (0, T] \end{cases}$$

- **The Big Question:**

Given a smooth initial condition $\boldsymbol{\eta}$, does the Euler system always admit smooth solutions $\mathbf{u}(t)$ for arbitrarily long times t ?

- There is numerical evidence for singularity formation in Euler (J.D. Gibbon 2008).

► Euler-Voigt system ($\Omega = [0, L]^3$)

$$\begin{cases} -\alpha^2 \partial_t \Delta \mathbf{u}_\alpha + \partial_t \mathbf{u}_\alpha + (\mathbf{u}_\alpha \cdot \nabla) \mathbf{u}_\alpha + \nabla p = \mathbf{0}, & \text{in } \Omega \times (0, T] \\ \nabla \cdot \mathbf{u}_\alpha = 0, & \text{in } \Omega \times (0, T] \\ \mathbf{u}_\alpha = \boldsymbol{\eta}_\alpha & \text{in } \Omega \text{ at } t = 0 \\ \text{Periodic Boundary Conditions} & \text{on } \partial\Omega \times (0, T] \end{cases}$$

- $\alpha > 0$ is the regularization parameter with units of length. Acts as a 'spectral filter'. The Euler-Voigt system is globally well-posed given smooth initial conditions (Y. Cao et al. 2006).
- As $\alpha \rightarrow 0$ the solutions of Euler-Voigt converge to the solutions of Euler and so its behaviour can be used to determine if the Euler equations will develop a finite-time singularity.

- Euler-Voigt system ($\Omega = [0, L]^3$)

$$\begin{cases} -\alpha^2 \partial_t \Delta \mathbf{u}_\alpha + \partial_t \mathbf{u}_\alpha + (\mathbf{u}_\alpha \cdot \nabla) \mathbf{u}_\alpha + \nabla p = \mathbf{0}, & \text{in } \Omega \times (0, T] \\ \nabla \cdot \mathbf{u}_\alpha = 0, & \text{in } \Omega \times (0, T] \\ \mathbf{u}_\alpha = \boldsymbol{\eta}_\alpha & \text{in } \Omega \text{ at } t = 0 \\ \text{Periodic Boundary Conditions} & \text{on } \partial\Omega \times (0, T] \end{cases}$$

- $\alpha > 0$ is the regularization parameter with units of length. Acts as a 'spectral filter'. The Euler-Voigt system is globally well-posed given smooth initial conditions (Y. Cao et al. 2006).
- As $\alpha \rightarrow 0$ the solutions of Euler-Voigt converge to the solutions of Euler and so its behaviour can be used to determine if the Euler equations will develop a finite-time singularity.

- ▶ Euler-Voigt system ($\Omega = [0, L]^3$)

$$\begin{cases} -\alpha^2 \partial_t \Delta \mathbf{u}_\alpha + \partial_t \mathbf{u}_\alpha + (\mathbf{u}_\alpha \cdot \nabla) \mathbf{u}_\alpha + \nabla p = \mathbf{0}, & \text{in } \Omega \times (0, T] \\ \nabla \cdot \mathbf{u}_\alpha = 0, & \text{in } \Omega \times (0, T] \\ \mathbf{u}_\alpha = \boldsymbol{\eta}_\alpha & \text{in } \Omega \text{ at } t = 0 \\ \text{Periodic Boundary Conditions} & \text{on } \partial\Omega \times (0, T] \end{cases}$$

- ▶ $\alpha > 0$ is the regularization parameter with units of length. Acts as a 'spectral filter'. The Euler-Voigt system is globally well-posed given smooth initial conditions (Y. Cao et al. 2006).
- ▶ As $\alpha \rightarrow 0$ the solutions of Euler-Voigt converge to the solutions of Euler and so its behaviour can be used to determine if the Euler equations will develop a finite-time singularity.

- Theorem 5.1 (A. Larios et al. 2010).

Theorem

Given initial data $\eta, \eta_\alpha \in H^s(\mathbb{T}^3)$ for some $s \geq 3$, let

$$\mathbf{u}, \mathbf{u}_\alpha \in C([0, T], H^s(\mathbb{T}^3)) \cap C^1([0, T], H^{s-1}(\mathbb{T}^3))$$

be the corresponding solutions to Euler and Euler-Voigt, respectively, where $0 < T < T_{\max}$ and $T_{\max}$ is the maximal time for which a solution to Euler exists and is unique. For all $t \in [0, T]$, we have

$$\begin{aligned} \|\mathbf{u}(t) - \mathbf{u}_\alpha(t)\|_{L^2}^2 + \alpha^2 \|\nabla(\mathbf{u}(t) - \mathbf{u}_\alpha(t))\|_{L^2}^2 \leq \\ (\|\eta - \eta_\alpha\|_{L^2}^2 + \alpha^2 \|\nabla(\eta - \eta_\alpha)\|_{L^2}^2) e^{Ct} + C\alpha^2(e^{Ct} - 1). \end{aligned}$$

- Proven by Theorem 1 in (A. Larios et al. 2018).

Theorem

Let $\eta_\alpha \in H^s$, $s \geq 3$ with $\nabla \cdot \eta_\alpha = 0$ and let $\mathbf{u}_\alpha$ be the corresponding unique solution of Euler-Voigt. Suppose that there exists a time $T^* > 0$ such that

$$\limsup_{\alpha \rightarrow 0^+} \left(\alpha \sup_{t \in [0, T^*]} \|\nabla \mathbf{u}_\alpha(t)\|_{L^2} \right) > 0.$$

Then the Euler equations with the initial condition η_α must develop a singularity within the interval $[0, T^*]$.

- Modified from Theorem 5.2 (A. Larios et al. 2010).

Theorem

Given initial data $\boldsymbol{\eta}, \boldsymbol{\eta}_\alpha \in H^s(\Omega)$ for some $s \geq 3$ satisfying

$$\|\boldsymbol{\eta} - \boldsymbol{\eta}_\alpha\|_{L^2}^2 + \alpha^2 \|\nabla(\boldsymbol{\eta} - \boldsymbol{\eta}_\alpha)\|_{L^2}^2 \rightarrow 0, \quad \text{as } \alpha \rightarrow 0,$$

let $\mathbf{u}, \mathbf{u}_\alpha \in C([0, T], H^s(\Omega)) \cap C^1([0, T], H^{s-1}(\Omega))$ be the corresponding solutions to Euler and Euler-Voigt, respectively. If there exists a finite time $T^* > 0$ such that the solutions $\mathbf{u}_\alpha$ satisfies

$$\limsup_{\alpha \rightarrow 0} \left(\alpha^2 \sup_{t \in [0, T^*]} \|\nabla \mathbf{u}_\alpha(t)\|_{L^2}^2 \right) > 0,$$

then $\mathbf{u}$ develops a singularity in the interval $[0, T^*]$.

- We consider analytic initial conditions that belong to the following Gevrey class

$$G^\sigma := \left\{ \mathbf{v} \in H^s(\Omega) : \|\mathbf{v}\|_{G^\sigma} = \langle \mathbf{v}, \mathbf{v} \rangle_{G^\sigma}^{1/2} < \infty \right\}, \quad s \geq 3, \quad \sigma > 0.$$

- We select the following inner product

$$\begin{aligned} \forall \mathbf{v}, \mathbf{u} \in G^\sigma, \quad \langle \mathbf{v}, \mathbf{u} \rangle_{G^\sigma} &:= \sum_{\mathbf{j} \in \mathbb{Z}^3} (1 + |2\pi \mathbf{j}|^2)^s e^{4\pi\sigma|\mathbf{j}|} \widehat{\mathbf{v}}_{\mathbf{j}} \cdot \overline{\widehat{\mathbf{u}}_{\mathbf{j}}} \\ &= \int_{\Omega} (1 + |D|^2)^s e^{2\sigma|D|} \mathbf{v} \cdot \mathbf{u} \, d\mathbf{x}. \\ &= \left\langle (1 + |D|^2)^s e^{2\sigma|D|} \mathbf{v}, \mathbf{u} \right\rangle_{L^2} \end{aligned}$$

- The operators $|D|$ and $e^{\sigma|D|}$ are defined via

$$\left[\widehat{|D| \mathbf{v}} \right]_{\mathbf{j}} := 2\pi |\mathbf{j}| \widehat{\mathbf{v}}_{\mathbf{j}}, \quad \left[\widehat{e^{\sigma|D|} \mathbf{v}} \right]_{\mathbf{j}} := e^{2\pi\sigma|\mathbf{j}|} \widehat{\mathbf{v}}_{\mathbf{j}}.$$

- ▶ We consider analytic initial conditions that belong to the following Gevrey class

$$G^\sigma := \left\{ \mathbf{v} \in H^s(\Omega) : \|\mathbf{v}\|_{G^\sigma} = \langle \mathbf{v}, \mathbf{v} \rangle_{G^\sigma}^{1/2} < \infty \right\}, \quad s \geq 3, \quad \sigma > 0.$$

- ▶ We select the following inner product

$$\begin{aligned} \forall \mathbf{v}, \mathbf{u} \in G^\sigma, \quad \langle \mathbf{v}, \mathbf{u} \rangle_{G^\sigma} &:= \sum_{\mathbf{j} \in \mathbb{Z}^3} (1 + |2\pi \mathbf{j}|^2)^s e^{4\pi\sigma|\mathbf{j}|} \widehat{\mathbf{v}}_{\mathbf{j}} \cdot \overline{\widehat{\mathbf{u}}_{\mathbf{j}}} \\ &= \int_{\Omega} (1 + |D|^2)^s e^{2\sigma|D|} \mathbf{v} \cdot \mathbf{u} \, d\mathbf{x}. \\ &= \left\langle (1 + |D|^2)^s e^{2\sigma|D|} \mathbf{v}, \mathbf{u} \right\rangle_{L^2} \end{aligned}$$

- ▶ The operators $|D|$ and $e^{\sigma|D|}$ are defined via

$$\left[\widehat{|D| \mathbf{v}} \right]_{\mathbf{j}} := 2\pi |\mathbf{j}| \widehat{\mathbf{v}}_{\mathbf{j}}, \quad \left[\widehat{e^{\sigma|D|} \mathbf{v}} \right]_{\mathbf{j}} := e^{2\pi|\mathbf{j}|} \widehat{\mathbf{v}}_{\mathbf{j}}.$$

- ▶ We consider analytic initial conditions that belong to the following Gevrey class

$$G^\sigma := \left\{ \mathbf{v} \in H^s(\Omega) : \|\mathbf{v}\|_{G^\sigma} = \langle \mathbf{v}, \mathbf{v} \rangle_{G^\sigma}^{1/2} < \infty \right\}, \quad s \geq 3, \quad \sigma > 0.$$

- ▶ We select the following inner product

$$\begin{aligned} \forall \mathbf{v}, \mathbf{u} \in G^\sigma, \quad \langle \mathbf{v}, \mathbf{u} \rangle_{G^\sigma} &:= \sum_{\mathbf{j} \in \mathbb{Z}^3} (1 + |2\pi \mathbf{j}|^2)^s e^{4\pi\sigma|\mathbf{j}|} \hat{\mathbf{v}}_{\mathbf{j}} \cdot \overline{\hat{\mathbf{u}}_{\mathbf{j}}} \\ &= \int_{\Omega} (1 + |D|^2)^s e^{2\sigma|D|} \mathbf{v} \cdot \mathbf{u} \, d\mathbf{x}. \\ &= \left\langle (1 + |D|^2)^s e^{2\sigma|D|} \mathbf{v}, \mathbf{u} \right\rangle_{L^2} \end{aligned}$$

- ▶ The operators $|D|$ and $e^{\sigma|D|}$ are defined via

$$\left[\widehat{|D| \mathbf{v}} \right]_{\mathbf{j}} := 2\pi |\mathbf{j}| \hat{\mathbf{v}}_{\mathbf{j}}, \quad \left[\widehat{e^{\sigma|D|} \mathbf{v}} \right]_{\mathbf{j}} := e^{2\pi|\mathbf{j}|} \hat{\mathbf{v}}_{\mathbf{j}}.$$

- ▶ We assume zero mean velocity without loss of generality.

$$V := \{\mathbf{v} \in G^\sigma : \nabla \cdot \mathbf{v} = 0, \int_{\Omega} \mathbf{v} \, d\mathbf{x} = 0\}.$$

- ▶ We use a closed manifold $\mathcal{M}_1 \subset V$ defined as

$$\mathcal{M}_1 = \{\mathbf{v} \in V : \|\mathbf{v}\|_{\dot{H}^1} = \mathbf{C}\}.$$

- ▶ Euler ($\mathbf{u}$) and Euler-Voigt ($\mathbf{u}_\alpha$) time scaling property

$$\tilde{\mathbf{u}}_{(\alpha)}(\mathbf{x}, t) = \lambda \mathbf{u}_{(\alpha)}(\mathbf{x}, \lambda t), \quad (3)$$

- We assume zero mean velocity without loss of generality.

$$V := \{\mathbf{v} \in G^\sigma : \nabla \cdot \mathbf{v} = 0, \int_{\Omega} \mathbf{v} \, d\mathbf{x} = 0\}.$$

- We use a closed manifold $\mathcal{M}_1 \subset V$ defined as

$$\mathcal{M}_1 = \{\mathbf{v} \in V : \|\mathbf{v}\|_{\dot{H}^1} = \mathbf{C}\}.$$

- Euler ($\mathbf{u}$) and Euler-Voigt ($\mathbf{u}_\alpha$) time scaling property

$$\tilde{\mathbf{u}}_{(\alpha)}(\mathbf{x}, t) = \lambda \mathbf{u}_{(\alpha)}(\mathbf{x}, \lambda t), \quad (3)$$

- We assume zero mean velocity without loss of generality.

$$V := \{\mathbf{v} \in G^\sigma : \nabla \cdot \mathbf{v} = 0, \int_{\Omega} \mathbf{v} \, d\mathbf{x} = 0\}.$$

- We use a closed manifold $\mathcal{M}_1 \subset V$ defined as

$$\mathcal{M}_1 = \{\mathbf{v} \in V : \|\mathbf{v}\|_{\dot{H}^1} = \mathbf{C}\}.$$

- Euler ($\mathbf{u}$) and Euler-Voigt ($\mathbf{u}_\alpha$) time scaling property

$$\tilde{\mathbf{u}}_{(\alpha)}(\mathbf{x}, t) = \lambda \mathbf{u}_{(\alpha)}(\mathbf{x}, \lambda t), \quad (3)$$

- Taylor Green Vortex ($\Omega = [0, L]^3$)

$$\eta_{TGV} = \begin{bmatrix} V_0 \sin(2\pi x/L) \cos(2\pi y/L) \cos(2\pi z/L) \\ -V_0 \cos(2\pi x/L) \sin(2\pi y/L) \cos(2\pi z/L) \\ 0 \end{bmatrix}$$

- Characteristic time-scale

$$t_c = \frac{L}{V_0}$$

- Using ($\Omega = [0, 2\pi]^3$) and $V_0 = 1$, Bustamente et al. 2012 showed numerical evidence for finite-time singularity in Euler from η_{TGV} at $T^* \approx 4$. We use ($\Omega = [0, 1]^3$) and so scale down to $V_0 = 1/2\pi$.

- Taylor Green Vortex ($\Omega = [0, L]^3$)

$$\eta_{TGV} = \begin{bmatrix} V_0 \sin(2\pi x/L) \cos(2\pi y/L) \cos(2\pi z/L) \\ -V_0 \cos(2\pi x/L) \sin(2\pi y/L) \cos(2\pi z/L) \\ 0 \end{bmatrix}$$

- Characteristic time-scale

$$t_c = \frac{L}{V_0}$$

- Using ($\Omega = [0, 2\pi]^3$) and $V_0 = 1$, Bustamente et al. 2012 showed numerical evidence for finite-time singularity in Euler from η_{TGV} at $T^* \approx 4$. We use ($\Omega = [0, 1]^3$) and so scale down to $V_0 = 1/2\pi$.

- Taylor Green Vortex ($\Omega = [0, L]^3$)

$$\eta_{TGV} = \begin{bmatrix} V_0 \sin(2\pi x/L) \cos(2\pi y/L) \cos(2\pi z/L) \\ -V_0 \cos(2\pi x/L) \sin(2\pi y/L) \cos(2\pi z/L) \\ 0 \end{bmatrix}$$

- Characteristic time-scale

$$t_c = \frac{L}{V_0}$$

- Using ($\Omega = [0, 2\pi]^3$) and $V_0 = 1$, Bustamente et al. 2012 showed numerical evidence for finite-time singularity in Euler from η_{TGV} at $T^* \approx 4$. We use ($\Omega = [0, 1]^3$) and so scale down to $V_0 = 1/2\pi$.

- Calculate C analytically

$$C = \|\boldsymbol{\eta}\|_{\dot{H}^1} = \|\nabla \boldsymbol{\eta}\|_{L^2} = \left[\sum_{i,j=1}^3 \int_{\Omega} (\partial_i \eta_j)^2 dx \right]^{1/2}$$

- Calculating for scaled $\boldsymbol{\eta}_{TGV}$

$$\|\boldsymbol{\eta}_{TGV}\|_{\dot{H}^1} = \sqrt{3}/2$$

- Create the manifold

$$\mathcal{M}_1 = \{\mathbf{v} \in V : \|\mathbf{v}\|_{\dot{H}^1} = \sqrt{3}/2\}.$$

- Calculate C analytically

$$C = \|\boldsymbol{\eta}\|_{\dot{H}^1} = \|\nabla \boldsymbol{\eta}\|_{L^2} = \left[\sum_{i,j=1}^3 \int_{\Omega} (\partial_i \eta_j)^2 dx \right]^{1/2}$$

- Calculating for scaled $\boldsymbol{\eta}_{TGV}$

$$\|\boldsymbol{\eta}_{TGV}\|_{\dot{H}^1} = \sqrt{3}/2$$

- Create the manifold

$$\mathcal{M}_1 = \{\mathbf{v} \in V : \|\mathbf{v}\|_{\dot{H}^1} = \sqrt{3}/2\}.$$

- Calculate C analytically

$$C = \|\boldsymbol{\eta}\|_{\dot{H}^1} = \|\nabla \boldsymbol{\eta}\|_{L^2} = \left[\sum_{i,j=1}^3 \int_{\Omega} (\partial_i \eta_j)^2 dx \right]^{1/2}$$

- Calculating for scaled $\boldsymbol{\eta}_{TGV}$

$$\|\boldsymbol{\eta}_{TGV}\|_{\dot{H}^1} = \sqrt{3}/2$$

- Create the manifold

$$\mathcal{M}_1 = \{\mathbf{v} \in V : \|\mathbf{v}\|_{\dot{H}^1} = \sqrt{3}/2\}.$$

- Beale-Kato-Majda (BKM) criterion (Beale et al. 1984): A Smooth solution u of the Euler system develops a singularity at $t = t_0$ if and only if

$$\lim_{t \rightarrow t_0} \int_0^t \|\omega\|_{L^\infty} d\tau = \infty, \quad \omega = \nabla \times u$$




Figure: (Bustamente et al. 2012 Fig. 1b) Maximum of vorticity $\|\omega(\cdot, t)\|_\infty$. Results from runs performed at different resolutions are displayed together: 512^3 (brown triangles), 1024^3 (blue squares), 2048^3 (green diamonds), and 4096^3 (red circles).




Figure: Maximum of vorticity $\|\omega(\cdot, t)\|_\infty$. Results from runs performed at different resolutions are displayed together: 128^3 (orange plus), 256^3 (cyan star), 512^3 (brown triangles), 1024^3 (blue squares).

Optimization Problem for Euler-Voigt

- Define our objective function $\Phi_{\alpha,T} : V \rightarrow \mathbb{R}^+$

$$\Phi_{\alpha,T}(\eta) := \|\nabla \mathbf{u}_\alpha(T; \eta_\alpha)\|_{L^2}^2 = \|\mathbf{u}_\alpha(T; \eta_\alpha)\|_{\dot{H}^1}^2$$

- Given $\alpha, T \in \mathbb{R}^+$, find

$$\tilde{\eta}_{\alpha,T} = \operatorname{argmax}_{\eta_\alpha \in \mathcal{M}_1} \Phi_{\alpha,T}(\eta_\alpha).$$

- If $\|\eta - \tilde{\eta}_{\alpha,T}\|_{L^2}^2 + \alpha^2 \|\nabla(\eta - \tilde{\eta}_{\alpha,T})\|_{L^2}^2 \rightarrow 0$ and $\alpha^2 \Phi_{\alpha,T}(\tilde{\eta}_{\alpha,T}) > 0$ as $\alpha \rightarrow 0$ then a singularity forms in Euler the interval $[0, T]$ from initial condition η .

- Define our objective function $\Phi_{\alpha,T} : V \rightarrow \mathbb{R}^+$

$$\Phi_{\alpha,T}(\eta) := \|\nabla \mathbf{u}_\alpha(T; \eta_\alpha)\|_{L^2}^2 = \|\mathbf{u}_\alpha(T; \eta_\alpha)\|_{\dot{H}^1}^2$$

- Given $\alpha, T \in \mathbb{R}^+$, find

$$\tilde{\eta}_{\alpha,T} = \operatorname{argmax}_{\eta_\alpha \in \mathcal{M}_1} \Phi_{\alpha,T}(\eta_\alpha).$$

- If $\|\eta - \tilde{\eta}_{\alpha,T}\|_{L^2}^2 + \alpha^2 \|\nabla(\eta - \tilde{\eta}_{\alpha,T})\|_{L^2}^2 \rightarrow 0$ and $\alpha^2 \Phi_{\alpha,T}(\tilde{\eta}_{\alpha,T}) > 0$ as $\alpha \rightarrow 0$ then a singularity forms in Euler the interval $[0, T]$ from initial condition η .

- Define our objective function $\Phi_{\alpha,T} : V \rightarrow \mathbb{R}^+$

$$\Phi_{\alpha,T}(\eta) := \|\nabla \mathbf{u}_\alpha(T; \eta_\alpha)\|_{L^2}^2 = \|\mathbf{u}_\alpha(T; \eta_\alpha)\|_{\dot{H}^1}^2$$

- Given $\alpha, T \in \mathbb{R}^+$, find

$$\tilde{\eta}_{\alpha,T} = \operatorname{argmax}_{\eta_\alpha \in \mathcal{M}_1} \Phi_{\alpha,T}(\eta_\alpha).$$

- If $\|\eta - \tilde{\eta}_{\alpha,T}\|_{L^2}^2 + \alpha^2 \|\nabla(\eta - \tilde{\eta}_{\alpha,T})\|_{L^2}^2 \rightarrow 0$ and $\alpha^2 \Phi_{\alpha,T}(\tilde{\eta}_{\alpha,T}) > 0$ as $\alpha \rightarrow 0$ then a singularity forms in Euler the interval $[0, T]$ from initial condition η .

- ▶ The first order finite difference equation for the Gâteaux differential $\Phi'_{\alpha,T}(\eta_\alpha; \eta'_\alpha) : V \times V \rightarrow \mathbb{R}$,

$$\Phi'_{\alpha,T}(\eta_\alpha; \eta'_\alpha) := \lim_{\epsilon \rightarrow 0} \frac{1}{\epsilon} [\Phi_{\alpha,T}(\eta_\alpha + \epsilon \eta'_\alpha) - \Phi_{\alpha,T}(\eta_\alpha)] .$$

- ▶ Substituting perturbations into Euler-Voigt

$$\mathbf{u}_\alpha \rightarrow \mathbf{u}_\alpha + \mathbf{u}'_\alpha, \eta_\alpha \rightarrow \eta_\alpha + \eta'_\alpha, \text{ and } p \rightarrow p + p'$$

- ▶ Linearization of Euler-Voigt

$$\mathcal{L} \begin{bmatrix} \mathbf{u}'_\alpha \\ p' \end{bmatrix} := \begin{bmatrix} \partial_t \mathbf{u}'_\alpha + (I - \alpha^2 \Delta)^{-1} (\mathbf{u}_\alpha \cdot \nabla \mathbf{u}'_\alpha + \mathbf{u}'_\alpha \cdot \nabla \mathbf{u}_\alpha + \nabla p') = 0 \\ \nabla \cdot \mathbf{u}'_\alpha \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\mathbf{u}'_\alpha(x, 0) = \eta'_\alpha$$

- ▶ The first order finite difference equation for the Gâteaux differential $\Phi'_{\alpha,T}(\boldsymbol{\eta}_\alpha; \boldsymbol{\eta}'_\alpha) : V \times V \rightarrow \mathbb{R}$,

$$\Phi'_{\alpha,T}(\boldsymbol{\eta}_\alpha; \boldsymbol{\eta}'_\alpha) := \lim_{\epsilon \rightarrow 0} \frac{1}{\epsilon} [\Phi_{\alpha,T}(\boldsymbol{\eta}_\alpha + \epsilon \boldsymbol{\eta}'_\alpha) - \Phi_{\alpha,T}(\boldsymbol{\eta}_\alpha)] .$$

- ▶ Substituting perturbations into Euler-Voigt

$$\mathbf{u}_\alpha \rightarrow \mathbf{u}_\alpha + \mathbf{u}'_\alpha, \boldsymbol{\eta}_\alpha \rightarrow \boldsymbol{\eta}_\alpha + \boldsymbol{\eta}'_\alpha, \text{ and } p \rightarrow p + p'$$

- ▶ Linearization of Euler-Voigt

$$\mathcal{L} \begin{bmatrix} \mathbf{u}'_\alpha \\ p' \end{bmatrix} := \begin{bmatrix} \partial_t \mathbf{u}'_\alpha + (I - \alpha^2 \Delta)^{-1} (\mathbf{u}_\alpha \cdot \nabla \mathbf{u}'_\alpha + \mathbf{u}'_\alpha \cdot \nabla \mathbf{u}_\alpha + \nabla p') = 0 \\ \nabla \cdot \mathbf{u}'_\alpha \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\mathbf{u}'_\alpha(x, 0) = \boldsymbol{\eta}'_\alpha$$

- ▶ The first order finite difference equation for the Gâteaux differential $\Phi'_{\alpha,T}(\boldsymbol{\eta}_\alpha; \boldsymbol{\eta}'_\alpha) : V \times V \rightarrow \mathbb{R}$,

$$\Phi'_{\alpha,T}(\boldsymbol{\eta}_\alpha; \boldsymbol{\eta}'_\alpha) := \lim_{\epsilon \rightarrow 0} \frac{1}{\epsilon} [\Phi_{\alpha,T}(\boldsymbol{\eta}_\alpha + \epsilon \boldsymbol{\eta}'_\alpha) - \Phi_{\alpha,T}(\boldsymbol{\eta}_\alpha)] .$$

- ▶ Substituting perturbations into Euler-Voigt

$$\mathbf{u}_\alpha \rightarrow \mathbf{u}_\alpha + \mathbf{u}'_\alpha, \boldsymbol{\eta}_\alpha \rightarrow \boldsymbol{\eta}_\alpha + \boldsymbol{\eta}'_\alpha, \text{ and } p \rightarrow p + p'$$

- ▶ Linearization of Euler-Voigt

$$\mathcal{L} \begin{bmatrix} \mathbf{u}'_\alpha \\ p' \end{bmatrix} := \begin{bmatrix} \partial_t \mathbf{u}'_\alpha + (I - \alpha^2 \Delta)^{-1} (\mathbf{u}_\alpha \cdot \nabla \mathbf{u}'_\alpha + \mathbf{u}'_\alpha \cdot \nabla \mathbf{u}_\alpha + \nabla p') = 0 \\ \nabla \cdot \mathbf{u}'_\alpha \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\mathbf{u}'_\alpha(x, 0) = \boldsymbol{\eta}'_\alpha$$

- Integration by parts on the following duality-pairing relation

$$\begin{aligned}
 \left(\mathcal{L} \begin{bmatrix} \mathbf{u}'_\alpha \\ \rho' \end{bmatrix}, \begin{bmatrix} \mathbf{u}^*_\alpha \\ \rho^* \end{bmatrix} \right) &:= \int_0^T \int_{\mathbb{T}^3} \mathcal{L} \begin{bmatrix} \mathbf{u}'_\alpha \\ \rho' \end{bmatrix}, \begin{bmatrix} \mathbf{u}^*_\alpha \\ \rho^* \end{bmatrix} d\mathbf{x} dt \\
 &= \left(\begin{bmatrix} \mathbf{u}'_\alpha \\ \rho' \end{bmatrix}, \mathcal{L}^* \begin{bmatrix} \mathbf{u}^*_\alpha \\ \rho^* \end{bmatrix} \right) + \int_{\mathbb{T}^3} \mathbf{u}'_\alpha(\mathbf{x}, T) \cdot \mathbf{u}^*_\alpha(\mathbf{x}, T) d\mathbf{x} \\
 &\quad - \int_{\mathbb{T}^3} \boldsymbol{\eta}'(\mathbf{x}) \cdot \mathbf{u}^*_\alpha(\mathbf{x}, 0) d\mathbf{x} \\
 &= 0.
 \end{aligned}$$

- Adjoint System

$$\mathcal{L}^* \begin{bmatrix} \mathbf{u}^*_\alpha \\ \rho^*_\alpha \end{bmatrix} := \begin{bmatrix} -\partial_t \mathbf{u}^*_\alpha - (\nabla(I - \alpha^2)^{-1} \mathbf{u}^*_\alpha + (\nabla(I - \alpha^2)^{-1} \mathbf{u}^*_\alpha)^\top) \mathbf{u}_\alpha - \nabla \rho^*_\alpha \\ -\nabla \cdot \mathbf{u}^*_\alpha \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\mathbf{u}^*_\alpha(\mathbf{x}, T) = 2|D|^2 \mathbf{u}_\alpha(\mathbf{x}, T)$$

- Integration by parts on the following duality-pairing relation

$$\begin{aligned}
 \left(\mathcal{L} \begin{bmatrix} \mathbf{u}'_\alpha \\ p' \end{bmatrix}, \begin{bmatrix} \mathbf{u}^*_\alpha \\ p^* \end{bmatrix} \right) &:= \int_0^T \int_{\mathbb{T}^3} \mathcal{L} \begin{bmatrix} \mathbf{u}'_\alpha \\ p' \end{bmatrix}, \begin{bmatrix} \mathbf{u}^*_\alpha \\ p^* \end{bmatrix} d\mathbf{x} dt \\
 &= \left(\begin{bmatrix} \mathbf{u}'_\alpha \\ p' \end{bmatrix}, \mathcal{L}^* \begin{bmatrix} \mathbf{u}^*_\alpha \\ p^* \end{bmatrix} \right) + \int_{\mathbb{T}^3} \mathbf{u}'_\alpha(\mathbf{x}, T) \cdot \mathbf{u}^*_\alpha(\mathbf{x}, T) d\mathbf{x} \\
 &\quad - \int_{\mathbb{T}^3} \boldsymbol{\eta}'(\mathbf{x}) \cdot \mathbf{u}^*_\alpha(\mathbf{x}, 0) d\mathbf{x} \\
 &= 0.
 \end{aligned}$$

- Adjoint System

$$\mathcal{L}^* \begin{bmatrix} \mathbf{u}^*_\alpha \\ p^*_\alpha \end{bmatrix} := \begin{bmatrix} -\partial_t \mathbf{u}^*_\alpha - (\nabla(I - \alpha^2)^{-1} \mathbf{u}^*_\alpha + (\nabla(I - \alpha^2)^{-1} \mathbf{u}^*_\alpha)^\top) \mathbf{u}_\alpha - \nabla p^*_\alpha \\ -\nabla \cdot \mathbf{u}^*_\alpha \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\mathbf{u}^*_\alpha(\mathbf{x}, T) = 2|D|^2 \mathbf{u}_\alpha(\mathbf{x}, T)$$

- ▶ Substituting $\Phi_{\alpha,T}$ into the Gâteaux differential,

$$\Phi'_{\alpha,T}(\eta; \eta') = 2 \langle \mathbf{u}_{\alpha}(\mathbf{x}, T), \mathbf{u}'_{\alpha}(\mathbf{x}, T) \rangle_{\dot{H}^1} = 2 \langle \mathbf{u}_{\alpha}^*(\mathbf{x}, 0), \mathbf{u}'_{\alpha}(\mathbf{x}, 0) \rangle_{L^2}^2$$

- ▶ From the Riesz representation theorem (Luenberger 1969)

$$\Phi'_{\alpha,T}(\eta : \eta') = \langle \nabla \Phi_{\alpha,T}(\eta_{\alpha}), \eta'_{\alpha} \rangle_{G^{\sigma}}$$

- ▶ The gradient of the objective functional is

$$\nabla \Phi_{\alpha,T}(\eta_{\alpha}) = (1 + |D|^2)^{-3} e^{-2\sigma|D|} \mathbf{u}_{\alpha}^*(x, 0)$$

- ▶ Substituting $\Phi_{\alpha,T}$ into the Gâteaux differential,

$$\Phi'_{\alpha,T}(\eta; \eta') = 2 \langle \mathbf{u}_{\alpha}(\mathbf{x}, T), \mathbf{u}'_{\alpha}(\mathbf{x}, T) \rangle_{\dot{H}^1} = 2 \langle \mathbf{u}_{\alpha}^*(\mathbf{x}, 0), \mathbf{u}'_{\alpha}(\mathbf{x}, 0) \rangle_{L^2}^2$$

- ▶ From the Riesz representation theorem (Luenberger 1969)

$$\Phi'_{\alpha,T}(\eta : \eta') = \langle \nabla \Phi_{\alpha,T}(\eta_{\alpha}), \eta'_{\alpha} \rangle_{G^{\sigma}}$$

- ▶ The gradient of the objective functional is

$$\nabla \Phi_{\alpha,T}(\eta_{\alpha}) = (1 + |D|^2)^{-3} e^{-2\sigma|D|} \mathbf{u}_{\alpha}^*(x, 0)$$

- ▶ Substituting $\Phi_{\alpha,T}$ into the Gâteaux differential,

$$\Phi'_{\alpha,T}(\eta; \eta') = 2 \langle \mathbf{u}_{\alpha}(\mathbf{x}, T), \mathbf{u}'_{\alpha}(\mathbf{x}, T) \rangle_{\dot{H}^1} = 2 \langle \mathbf{u}_{\alpha}^*(\mathbf{x}, 0), \mathbf{u}'_{\alpha}(\mathbf{x}, 0) \rangle_{L^2}^2$$

- ▶ From the Riesz representation theorem (Luenberger 1969)

$$\Phi'_{\alpha,T}(\eta : \eta') = \langle \nabla \Phi_{\alpha,T}(\eta_{\alpha}), \eta'_{\alpha} \rangle_{G^{\sigma}}$$

- ▶ The gradient of the objective functional is

$$\nabla \Phi_{\alpha,T}(\eta_{\alpha}) = (1 + |D|^2)^{-3} e^{-2\sigma|D|} \mathbf{u}_{\alpha}^*(x, 0)$$

- ▶ Project the gradient $\nabla\Phi_T(\eta_\alpha)$ onto the tangent space to $\mathcal{M}_1$ at η_α .
- ▶ Perform vector transport operation to construct a Riemannian conjugate ascent direction.
- ▶ Retract back to $\mathcal{M}_1$.

- ▶ Project the gradient $\nabla\Phi_T(\eta_\alpha)$ onto the tangent space to $\mathcal{M}_1$ at η_α .
- ▶ Perform vector transport operation to construct a Riemannian conjugate ascent direction.
- ▶ Retract back to $\mathcal{M}_1$.

- ▶ Project the gradient $\nabla\Phi_T(\eta_\alpha)$ onto the tangent space to $\mathcal{M}_1$ at η_α .
- ▶ Perform vector transport operation to construct a Riemannian conjugate ascent direction.
- ▶ Retract back to $\mathcal{M}_1$.

- Tangent space to $\mathcal{M}_1$ at $\mathbf{v}$

$$\mathcal{T}_{\mathbf{v}}\mathcal{M}_1 = \{z \in V : \langle \mathbf{v}, z \rangle_{\dot{H}^1} = 0\}$$

- For any $z \in \mathcal{T}_{\mathbf{v}}\mathcal{M}_1$

$$\begin{aligned} \langle \mathbf{v}, z \rangle_{\dot{H}^1} &= \int_{\mathbb{T}^3} |D|\mathbf{v} \cdot |D|z \, d\mathbf{x} \\ &= \int_{\mathbb{T}^3} (1 + |D|^2)^3 e^{2\sigma|D|} \left((1 + |D|^2)^{-3} e^{-2\sigma|D|} |D|^2 \mathbf{v} \right) \cdot z \, d\mathbf{x} \\ &= \left\langle (1 + |D|^2)^{-3} e^{-2\sigma|D|} |D|^2 \mathbf{v}, z \right\rangle_{G^\sigma} = 0 \end{aligned}$$

- Characterized using the unit 'normal' vector

$$\mathbf{n}_{\mathbf{v}} = \frac{(1 + |D|^2)^{-3} e^{-2\sigma|D|} |D|^2 \mathbf{v}}{\|(1 + |D|^2)^{-3} e^{-2\sigma|D|} |D|^2 \mathbf{v}\|_{G^\sigma}}$$

- Tangent space to $\mathcal{M}_1$ at $\mathbf{v}$

$$\mathcal{T}_{\mathbf{v}}\mathcal{M}_1 = \{z \in V : \langle \mathbf{v}, z \rangle_{\dot{H}^1} = 0\}$$

- For any $z \in \mathcal{T}_{\mathbf{v}}\mathcal{M}_1$

$$\begin{aligned} \langle \mathbf{v}, z \rangle_{\dot{H}^1} &= \int_{\mathbb{T}^3} |D|\mathbf{v} \cdot |D|\mathbf{z} \, d\mathbf{x} \\ &= \int_{\mathbb{T}^3} (1 + |D|^2)^3 e^{2\sigma|D|} \left((1 + |D|^2)^{-3} e^{-2\sigma|D|} |D|^2 \mathbf{v} \right) \cdot \mathbf{z} \, d\mathbf{x} \\ &= \left\langle (1 + |D|^2)^{-3} e^{-2\sigma|D|} |D|^2 \mathbf{v}, \mathbf{z} \right\rangle_{G^\sigma} = 0 \end{aligned}$$

- Characterized using the unit 'normal' vector

$$\mathbf{n}_{\mathbf{v}} = \frac{(1 + |D|^2)^{-3} e^{-2\sigma|D|} |D|^2 \mathbf{v}}{\|(1 + |D|^2)^{-3} e^{-2\sigma|D|} |D|^2 \mathbf{v}\|_{G^\sigma}}$$

- Tangent space to $\mathcal{M}_1$ at $\mathbf{v}$

$$\mathcal{T}_{\mathbf{v}}\mathcal{M}_1 = \{z \in V : \langle \mathbf{v}, z \rangle_{\dot{H}^1} = 0\}$$

- For any $z \in \mathcal{T}_{\mathbf{v}}\mathcal{M}_1$

$$\begin{aligned} \langle \mathbf{v}, z \rangle_{\dot{H}^1} &= \int_{\mathbb{T}^3} |D|\mathbf{v} \cdot |D|\mathbf{z} \, d\mathbf{x} \\ &= \int_{\mathbb{T}^3} (1 + |D|^2)^3 e^{2\sigma|D|} \left((1 + |D|^2)^{-3} e^{-2\sigma|D|} |D|^2 \mathbf{v} \right) \cdot \mathbf{z} \, d\mathbf{x} \\ &= \left\langle (1 + |D|^2)^{-3} e^{-2\sigma|D|} |D|^2 \mathbf{v}, \mathbf{z} \right\rangle_{G^\sigma} = 0 \end{aligned}$$

- Characterized using the unit 'normal' vector

$$\mathbf{n}_{\mathbf{v}} = \frac{(1 + |D|^2)^{-3} e^{-2\sigma|D|} |D|^2 \mathbf{v}}{\|(1 + |D|^2)^{-3} e^{-2\sigma|D|} |D|^2 \mathbf{v}\|_{G^\sigma}}$$

- ▶ Projection operator $P_{\mathcal{T}_v\mathcal{M}_1} : V \rightarrow \mathcal{T}_v\mathcal{M}_1$

$$\forall \mathbf{w} \in V, \quad P_{\mathcal{T}_v\mathcal{M}_1} \mathbf{w} = \mathbf{w} - \langle \mathbf{n}_v, \mathbf{w} \rangle_{G^\sigma} \mathbf{n}_v$$

- ▶ Retraction operator $R : V \rightarrow \mathcal{M}_1$

$$R(\mathbf{v}) = \frac{\sqrt{3}}{2} \frac{\mathbf{v}}{\|\mathbf{v}\|_{\dot{H}^1}}, \quad \mathbf{v} \neq 0$$

- ▶ Vector field ξ transported along $\mathcal{M}_1$ in the direction of ϕ

$$\mathcal{TM}_1 \oplus \mathcal{TM}_1 \rightarrow \mathcal{TM}_1 : (\varphi, \xi) \mapsto \Gamma_\varphi(\xi) \in \mathcal{TM}_1,$$

where $\mathcal{TM}_1 = \cup_{\mathbf{v} \in \mathcal{M}_1} \mathcal{T}_v\mathcal{M}_1$ is the tangent bundle on $\mathcal{M}_1$ (absil et al. 2008). It is calculated via the differentiated retraction

$$\Gamma_\varphi(\xi) = \left. \frac{d}{dt} R(\varphi + t\xi) \right|_{t=0}.$$

- ▶ Projection operator $P_{\mathcal{T}_v\mathcal{M}_1} : V \rightarrow \mathcal{T}_v\mathcal{M}_1$

$$\forall \mathbf{w} \in V, \quad P_{\mathcal{T}_v\mathcal{M}_1} \mathbf{w} = \mathbf{w} - \langle \mathbf{n}_v, \mathbf{w} \rangle_{G^\sigma} \mathbf{n}_v$$

- ▶ Retraction operator $R : V \rightarrow \mathcal{M}_1$

$$R(\mathbf{v}) = \frac{\sqrt{3}}{2} \frac{\mathbf{v}}{\|\mathbf{v}\|_{\dot{H}^1}}, \quad \mathbf{v} \neq 0$$

- ▶ Vector field ξ transported along $\mathcal{M}_1$ in the direction of ϕ

$$\mathcal{TM}_1 \oplus \mathcal{TM}_1 \rightarrow \mathcal{TM}_1 : (\varphi, \xi) \mapsto \Gamma_\varphi(\xi) \in \mathcal{TM}_1,$$

where $\mathcal{TM}_1 = \cup_{\mathbf{v} \in \mathcal{M}_1} \mathcal{T}_v\mathcal{M}_1$ is the tangent bundle on $\mathcal{M}_1$ (absil et al. 2008). It is calculated via the differentiated retraction

$$\Gamma_\varphi(\xi) = \left. \frac{d}{dt} R(\varphi + t\xi) \right|_{t=0}.$$

- ▶ Projection operator $P_{\mathcal{T}_v\mathcal{M}_1} : V \rightarrow \mathcal{T}_v\mathcal{M}_1$

$$\forall \mathbf{w} \in V, \quad P_{\mathcal{T}_v\mathcal{M}_1} \mathbf{w} = \mathbf{w} - \langle \mathbf{n}_v, \mathbf{w} \rangle_{G^\sigma} \mathbf{n}_v$$

- ▶ Retraction operator $R : V \rightarrow \mathcal{M}_1$

$$R(\mathbf{v}) = \frac{\sqrt{3}}{2} \frac{\mathbf{v}}{\|\mathbf{v}\|_{\dot{H}^1}}, \quad \mathbf{v} \neq 0$$

- ▶ Vector field ξ transported along $\mathcal{M}_1$ in the direction of ϕ

$$\mathcal{TM}_1 \oplus \mathcal{TM}_1 \rightarrow \mathcal{TM}_1 : (\varphi, \xi) \mapsto \Gamma_\varphi(\xi) \in \mathcal{TM}_1,$$

where $\mathcal{TM}_1 = \cup_{\mathbf{v} \in \mathcal{M}_1} \mathcal{T}_v\mathcal{M}_1$ is the tangent bundle on $\mathcal{M}_1$ (absil et al. 2008). It is calculated via the differentiated retraction

$$\Gamma_\varphi(\xi) = \left. \frac{d}{dt} R(\varphi + t\xi) \right|_{t=0}.$$

- We can now construct the iterative relation

$$\boldsymbol{\eta}_\alpha^{(n+1)} = R \left[\boldsymbol{\eta}_\alpha^{(n)} + \tau_n \mathbf{d}^{(n)} \right]$$

- Let $\mathbf{G}^{(n)} := \nabla \Phi_{\alpha, T}(\boldsymbol{\eta}_\alpha^{(n)})$. Let $\mathcal{T}_n := \mathcal{T}_{\boldsymbol{\eta}_\alpha^{(n)}} \mathcal{M}_1$. $\mathbf{d}^{(n)}$ is computed as

$$\mathbf{d}^{(0)} = P_{\mathcal{T}_0} \mathbf{G}^{(0)}$$

$$\mathbf{d}^{(n)} = P_{\mathcal{T}_n} \mathbf{G}^{(n)} + \beta_n \Gamma_{\tau_{n-1} \mathbf{d}^{(n-1)}}(\mathbf{d}^{(n-1)}), \quad n \geq 1.$$

- 'Momentum' term is computed using the Polak-Ribière approach (Nocedal et al. 1999).

$$\beta_n = \frac{\left\langle P_{\mathcal{T}_n} \mathbf{G}^{(n)}, \left(P_{\mathcal{T}_n} \mathbf{G}^{(n)} - \Gamma_{\tau_{n-1} \mathbf{d}^{(n-1)}} P_{\mathcal{T}_{n-1}} \mathbf{G}^{(n-1)} \right) \right\rangle_{G^\sigma}}{\|P_{\mathcal{T}_{n-1}} \mathbf{G}^{(n-1)}\|_{G^\sigma}^2}$$

- Step size

$$\tau_n = \operatorname{argmax}_{\tau > 0} \Phi_T \left(R \left[\boldsymbol{\eta}_\alpha^{(n)} + \tau \mathbf{d}^{(n)} \right] \right)$$

- We can now construct the iterative relation

$$\boldsymbol{\eta}_\alpha^{(n+1)} = R \left[\boldsymbol{\eta}_\alpha^{(n)} + \tau_n \mathbf{d}^{(n)} \right]$$

- Let $\mathbf{G}^{(n)} := \nabla \Phi_{\alpha, T}(\boldsymbol{\eta}_\alpha^{(n)})$. Let $\mathcal{T}_n := \mathcal{T}_{\boldsymbol{\eta}_\alpha^{(n)}} \mathcal{M}_1$. $\mathbf{d}^{(n)}$ is computed as

$$\mathbf{d}^{(0)} = P_{\mathcal{T}_0} \mathbf{G}^{(0)}$$

$$\mathbf{d}^{(n)} = P_{\mathcal{T}_n} \mathbf{G}^{(n)} + \beta_n \Gamma_{\tau_{n-1} \mathbf{d}^{(n-1)}}(\mathbf{d}^{(n-1)}), \quad n \geq 1.$$

- 'Momentum' term is computed using the Polak-Ribière approach (Nocedal et al. 1999).

$$\beta_n = \frac{\left\langle P_{\mathcal{T}_n} \mathbf{G}^{(n)}, \left(P_{\mathcal{T}_n} \mathbf{G}^{(n)} - \Gamma_{\tau_{n-1} \mathbf{d}^{(n-1)}} P_{\mathcal{T}_{n-1}} \mathbf{G}^{(n-1)} \right) \right\rangle_{G^\sigma}}{\|P_{\mathcal{T}_{n-1}} \mathbf{G}^{(n-1)}\|_{G^\sigma}^2}$$

- Step size

$$\tau_n = \operatorname{argmax}_{\tau > 0} \Phi_T \left(R \left[\boldsymbol{\eta}_\alpha^{(n)} + \tau \mathbf{d}^{(n)} \right] \right)$$

- We can now construct the iterative relation

$$\boldsymbol{\eta}_\alpha^{(n+1)} = R \left[\boldsymbol{\eta}_\alpha^{(n)} + \tau_n \mathbf{d}^{(n)} \right]$$

- Let $\mathbf{G}^{(n)} := \nabla \Phi_{\alpha, T}(\boldsymbol{\eta}_\alpha^{(n)})$. Let $\mathcal{T}_n := \mathcal{T}_{\boldsymbol{\eta}_\alpha^{(n)}} \mathcal{M}_1$. $\mathbf{d}^{(n)}$ is computed as

$$\mathbf{d}^{(0)} = P_{\mathcal{T}_0} \mathbf{G}^{(0)}$$

$$\mathbf{d}^{(n)} = P_{\mathcal{T}_n} \mathbf{G}^{(n)} + \beta_n \Gamma_{\tau_{n-1} \mathbf{d}^{(n-1)}}(\mathbf{d}^{(n-1)}), \quad n \geq 1.$$

- 'Momentum' term is computed using the Polak-Ribière approach (Nocedal et al. 1999).

$$\beta_n = \frac{\left\langle P_{\mathcal{T}_n} \mathbf{G}^{(n)}, \left(P_{\mathcal{T}_n} \mathbf{G}^{(n)} - \Gamma_{\tau_{n-1} \mathbf{d}^{(n-1)}} P_{\mathcal{T}_{n-1}} \mathbf{G}^{(n-1)} \right) \right\rangle_{G^\sigma}}{\|P_{\mathcal{T}_{n-1}} \mathbf{G}^{(n-1)}\|_{G^\sigma}^2}$$

- Step size

$$\tau_n = \operatorname{argmax}_{\tau > 0} \Phi_T \left(R \left[\boldsymbol{\eta}_\alpha^{(n)} + \tau \mathbf{d}^{(n)} \right] \right)$$

- We can now construct the iterative relation

$$\boldsymbol{\eta}_\alpha^{(n+1)} = R \left[\boldsymbol{\eta}_\alpha^{(n)} + \tau_n \mathbf{d}^{(n)} \right]$$

- Let $\mathbf{G}^{(n)} := \nabla \Phi_{\alpha, T}(\boldsymbol{\eta}_\alpha^{(n)})$. Let $\mathcal{T}_n := \mathcal{T}_{\boldsymbol{\eta}_\alpha^{(n)}} \mathcal{M}_1$. $\mathbf{d}^{(n)}$ is computed as

$$\mathbf{d}^{(0)} = P_{\mathcal{T}_0} \mathbf{G}^{(0)}$$

$$\mathbf{d}^{(n)} = P_{\mathcal{T}_n} \mathbf{G}^{(n)} + \beta_n \Gamma_{\tau_{n-1} \mathbf{d}^{(n-1)}}(\mathbf{d}^{(n-1)}), \quad n \geq 1.$$

- 'Momentum' term is computed using the Polak-Ribière approach (Nocedal et al. 1999).

$$\beta_n = \frac{\left\langle P_{\mathcal{T}_n} \mathbf{G}^{(n)}, \left(P_{\mathcal{T}_n} \mathbf{G}^{(n)} - \Gamma_{\tau_{n-1} \mathbf{d}^{(n-1)}} P_{\mathcal{T}_{n-1}} \mathbf{G}^{(n-1)} \right) \right\rangle_{G^\sigma}}{\|P_{\mathcal{T}_{n-1}} \mathbf{G}^{(n-1)}\|_{G^\sigma}^2}$$

- Step size

$$\tau_n = \operatorname{argmax}_{\tau > 0} \Phi_T \left(R \left[\boldsymbol{\eta}_\alpha^{(n)} + \tau \mathbf{d}^{(n)} \right] \right)$$




Figure: (Zhao et al. 2023, Fig. 1)




Figure: (Zhao et al. 2023, Fig. 1)




Figure: (Zhao et al. 2023, Fig. 1)




Figure: (Zhao et al. 2023, Fig. 1)

► Periodic algorithm restart ($\beta_n = 0$)



$$n = 20k, \quad k \in \mathbb{Z}^+$$

► The search direction $\mathbf{d}^{(n)}$ is not an ascent direction

$$\frac{\langle \mathbf{d}^{(n)}, P_{\mathcal{T}_n} \mathbf{G}^{(n)} \rangle_{G^\sigma}}{\|\mathbf{d}^{(n)}\|_{G^\sigma} \|P_{\mathcal{T}_n} \mathbf{G}^{(n)}\|_{G^\sigma}} < \epsilon, \quad 0 < \epsilon \ll 1.$$

► Iterations are stopped when the relative change of $\Phi_{\alpha, T}$ is below a specified tolerance

$$0 \leq \frac{\Phi_{\alpha, T}(\eta_\alpha^{(n+1)}) - \Phi_{\alpha, T}(\eta_\alpha^{(n)})}{\Phi_{\alpha, T}(\eta_\alpha^{(n)})} < tol.$$

- Periodic algorithm restart ($\beta_n = 0$)



$$n = 20k, \quad k \in \mathbb{Z}^+$$

- The search direction $\mathbf{d}^{(n)}$ is not an ascent direction

$$\frac{\langle \mathbf{d}^{(n)}, P_{\mathcal{T}_n} \mathbf{G}^{(n)} \rangle_{G^\sigma}}{\|\mathbf{d}^{(n)}\|_{G^\sigma} \|P_{\mathcal{T}_n} \mathbf{G}^{(n)}\|_{G^\sigma}} < \epsilon, \quad 0 < \epsilon \ll 1.$$

- Iterations are stopped when the relative change of $\Phi_{\alpha, T}$ is below a specified tolerance

$$0 \leq \frac{\Phi_{\alpha, T}(\eta_\alpha^{(n+1)}) - \Phi_{\alpha, T}(\eta_\alpha^{(n)})}{\Phi_{\alpha, T}(\eta_\alpha^{(n)})} < tol.$$

- Periodic algorithm restart ($\beta_n = 0$)



$$n = 20k, \quad k \in \mathbb{Z}^+$$

- The search direction $\mathbf{d}^{(n)}$ is not an ascent direction

$$\frac{\langle \mathbf{d}^{(n)}, P_{\mathcal{T}_n} \mathbf{G}^{(n)} \rangle_{G^\sigma}}{\|\mathbf{d}^{(n)}\|_{G^\sigma} \|P_{\mathcal{T}_n} \mathbf{G}^{(n)}\|_{G^\sigma}} < \epsilon, \quad 0 < \epsilon \ll 1.$$

- Iterations are stopped when the relative change of $\Phi_{\alpha, T}$ is below a specified tolerance

$$0 \leq \frac{\Phi_{\alpha, T}(\eta_\alpha^{(n+1)}) - \Phi_{\alpha, T}(\eta_\alpha^{(n)})}{\Phi_{\alpha, T}(\eta_\alpha^{(n)})} < tol.$$

- Periodic algorithm restart ($\beta_n = 0$)



$$n = 20k, \quad k \in \mathbb{Z}^+$$

- The search direction $\mathbf{d}^{(n)}$ is not an ascent direction

$$\frac{\left\langle \mathbf{d}^{(n)}, P_{\mathcal{T}_n} \mathbf{G}^{(n)} \right\rangle_{G^\sigma}}{\|\mathbf{d}^{(n)}\|_{G^\sigma} \|P_{\mathcal{T}_n} \mathbf{G}^{(n)}\|_{G^\sigma}} < \epsilon, \quad 0 < \epsilon \ll 1.$$

- Iterations are stopped when the relative change of $\Phi_{\alpha, T}$ is below a specified tolerance

$$0 \leq \frac{\Phi_{\alpha, T}(\boldsymbol{\eta}_\alpha^{(n+1)}) - \Phi_{\alpha, T}(\boldsymbol{\eta}_\alpha^{(n)})}{\Phi_{\alpha, T}(\boldsymbol{\eta}_\alpha^{(n)})} < tol.$$

- ▶ Discretized in space using a standard pseudospectral Fourier method
- ▶ Derivatives are evaluated in the Fourier space
- ▶ Nonlinear products are computed in the physical space on an uniform grid with N points in each direction.
- ▶ Explicit fourth-order Runge-Kutta
- ▶ 2/3 dealiasing
- ▶ Domain

resolution : $128^3, 256^3, 512^3, 1024^3$

α : $1/1024, 2/1024, 4/1024, 8/1024, 16/1024$

T : 3, 5

- ▶ Gevrey class

σ : 1E-5, 1E-4, 1E-3, 1E-2, 1E-1

s : 3

- ▶ Discretized in space using a standard pseudospectral Fourier method
- ▶ Derivatives are evaluated in the Fourier space
- ▶ Nonlinear products are computed in the physical space on an uniform grid with N points in each direction.
- ▶ Explicit fourth-order Runge-Kutta
- ▶ 2/3 dealiasing
- ▶ Domain

resolution : $128^3, 256^3, 512^3, 1024^3$

α : $1/1024, 2/1024, 4/1024, 8/1024, 16/1024$

T : 3, 5

- ▶ Gevrey class

σ : 1E-5, 1E-4, 1E-3, 1E-2, 1E-1

s : 3

- ▶ Discretized in space using a standard pseudospectral Fourier method
- ▶ Derivatives are evaluated in the Fourier space
- ▶ Nonlinear products are computed in the physical space on an uniform grid with N points in each direction.

- ▶ Explicit fourth-order Runge-Kutta
- ▶ 2/3 dealiasing
- ▶ Domain

resolution : $128^3, 256^3, 512^3, 1024^3$

$\alpha : 1/1024, 2/1024, 4/1024, 8/1024, 16/1024$

$T : 3, 5$

- ▶ Gevrey class

$\sigma : 1\text{E-}5, 1\text{E-}4, 1\text{E-}3, 1\text{E-}2, 1\text{E-}1$

$s : 3$

- ▶ Discretized in space using a standard pseudospectral Fourier method
- ▶ Derivatives are evaluated in the Fourier space
- ▶ Nonlinear products are computed in the physical space on an uniform grid with N points in each direction.
- ▶ Explicit fourth-order Runge-Kutta

▶ 2/3 dealiasing

▶ Domain

resolution : $128^3, 256^3, 512^3, 1024^3$

$\alpha : 1/1024, 2/1024, 4/1024, 8/1024, 16/1024$

$T : 3, 5$

▶ Gevrey class

$\sigma : 1\text{E-}5, 1\text{E-}4, 1\text{E-}3, 1\text{E-}2, 1\text{E-}1$

$s : 3$

- ▶ Discretized in space using a standard pseudospectral Fourier method
- ▶ Derivatives are evaluated in the Fourier space
- ▶ Nonlinear products are computed in the physical space on an uniform grid with N points in each direction.
- ▶ Explicit fourth-order Runge-Kutta
- ▶ 2/3 dealiasing

▶ Domain

resolution : $128^3, 256^3, 512^3, 1024^3$

$\alpha : 1/1024, 2/1024, 4/1024, 8/1024, 16/1024$

$T : 3, 5$

▶ Gevrey class

$\sigma : 1\text{E-}5, 1\text{E-}4, 1\text{E-}3, 1\text{E-}2, 1\text{E-}1$

$s : 3$

- ▶ Discretized in space using a standard pseudospectral Fourier method
- ▶ Derivatives are evaluated in the Fourier space
- ▶ Nonlinear products are computed in the physical space on an uniform grid with N points in each direction.
- ▶ Explicit fourth-order Runge-Kutta
- ▶ 2/3 dealiasing
- ▶ Domain

resolution : $128^3, 256^3, 512^3, 1024^3$

α : $1/1024, 2/1024, 4/1024, 8/1024, 16/1024$

T : 3, 5

- ▶ Gevrey class

σ : 1E-5, 1E-4, 1E-3, 1E-2, 1E-1

s : 3

- ▶ Discretized in space using a standard pseudospectral Fourier method
- ▶ Derivatives are evaluated in the Fourier space
- ▶ Nonlinear products are computed in the physical space on an uniform grid with N points in each direction.
- ▶ Explicit fourth-order Runge-Kutta
- ▶ 2/3 dealiasing
- ▶ Domain

resolution : $128^3, 256^3, 512^3, 1024^3$

α : $1/1024, 2/1024, 4/1024, 8/1024, 16/1024$

T : 3, 5

- ▶ Gevrey class

σ : 1E-5, 1E-4, 1E-3, 1E-2, 1E-1

s : 3

- Relative error to $\Phi_{\alpha,T}(\eta_{TGV})$ for $dt = 2^{-8}$. Resolution: 128

	1/1024	2/1024	4/1024	8/1024	16/1024
2^{-2}	Inf	Inf	Inf	0.0045	7.00E-05
2^{-3}	Inf	Inf	0.0572	0.0011	3.64E-06
2^{-4}	0.1662	0.1157	0.0332	4.85E-04	2.02E-07
2^{-5}	0.1093	0.0715	0.0181	2.33E-04	1.21E-08
2^{-6}	0.0593	0.0369	0.0085	1.02E-04	8.63E-10
2^{-7}	0.0228	0.0137	0.003	3.45E-05	9.44E-11
2^{-8}	0	0	0	0	0

- Relative error to $\Phi_{\alpha,T}(\eta_{TGV})$ for $dt = 2^{-8}$. Resolution: 256

	1/1024	2/1024	4/1024	8/1024	16/1024
2^{-2}	Inf	Inf	Inf	0.003	7.00E-05
2^{-3}	Inf	Inf	0.0112	1.82E-04	3.64E-06
2^{-4}	Inf	0.0652	0.0021	9.46E-06	2.01E-07
2^{-5}	0.1097	0.0348	7.13E-04	5.12E-07	1.17E-08
2^{-6}	0.0619	0.0171	3.08E-04	3.00E-08	7.01E-10
2^{-7}	0.025	0.0157	1.05E-04	1.96E-09	4.04E-11
2^{-8}	0	0	0	0	0

- Relative error to $\Phi_{\alpha,T}(\eta_{TGV})$ for $dt = 2^{-8}$. Resolution: 512

	1/1024	2/1024	4/1024	8/1024	16/1024
2^{-2}	Inf	Inf	Inf	0.003	7.00E-05
2^{-3}	Inf	Inf	0.009	1.82E-04	3.64E-06
2^{-4}	Inf	0.0174	7.61E-04	9.45E-06	2.01E-07
2^{-5}	Inf	0.0036	3.63E-05	5.10E-07	1.17E-08
2^{-6}	0.0306	7.85E-04	1.72E-06	2.90E-08	7.01E-10
2^{-7}	0.0115	2.42E-04	8.42E-08	1.63E-09	4.04E-11
2^{-8}	0	0	0	0	0

- Relative error to $\Phi_{\alpha,T}(\eta_{TGV})$ for $dt = 2^{-8}$. Resolution: 1024

	1/1024	2/1024	4/1024	8/1024	16/1024
2^{-2}	Inf	Inf	Inf	0.003	7.00E-05
2^{-3}	Inf	Inf	0.009	1.82E-04	3.64E-06
2^{-4}	Inf	0.015	7.61E-04	9.45E-06	2.01E-07
2^{-5}	Inf	0.0022	3.64E-05	5.10E-07	1.17E-08
2^{-6}	0.0039	2.64E-04	1.71E-06	2.90E-08	7.01E-10
2^{-7}	5.17E-04	4.92E-06	8.36E-08	1.63E-09	4.04E-11
2^{-8}	0	0	0	0	0

- Maximum relative error of α -Energy in $[0, T]$. Resolution: 128

	1/1024	2/1024	4/1024	8/1024	16/1024
2^{-2}	Inf	Inf	Inf	3.82E-04	8.73E-06
2^{-3}	Inf	Inf	1.94E-03	6.91E-05	3.72E-07
2^{-4}	3.38E-03	2.74E-03	1.11E-03	3.05E-05	1.76E-08
2^{-5}	2.38E-03	1.81E-03	6.49E-04	1.57E-05	9.59E-10
2^{-6}	1.51E-03	1.09E-03	3.56E-04	8.05E-06	7.55E-11
2^{-7}	8.67E-04	6.06E-04	1.87E-04	4.08E-06	1.47E-11
2^{-8}	4.69E-04	3.21E-04	9.59E-05	2.05E-06	6.00E-12

- Maximum relative error of α -Energy in $[0, T]$. Resolution: 256

	1/1024	2/1024	4/1024	8/1024	16/1024
2^{-2}	Inf	Inf	Inf	2.91E-04	8.73E-06
2^{-3}	Inf	Inf	4.34E-04	1.49E-05	3.72E-07
2^{-4}	Inf	8.36E-04	5.61E-05	6.85E-07	1.75E-08
2^{-5}	9.63E-04	4.35E-04	1.80E-05	3.40E-08	9.09E-10
2^{-6}	6.33E-04	2.48E-04	8.97E-06	1.89E-09	4.95E-11
2^{-7}	3.83E-04	1.37E-04	4.59E-06	1.38E-10	2.89E-12
2^{-8}	2.14E-04	7.23E-05	2.33E-06	2.37E-11	2.95E-12

- Maximum relative error of α -Energy in $[0, T]$. Resolution: 512

	1/1024	2/1024	4/1024	8/1024	16/1024
2^{-2}	Inf	Inf	Inf	2.91E-04	8.73E-06
2^{-3}	Inf	Inf	3.85E-04	1.49E-05	3.72E-07
2^{-4}	Inf	2.72E-04	2.58E-05	6.85E-07	1.75E-08
2^{-5}	Inf	2.89E-05	1.12E-06	3.38E-08	9.09E-10
2^{-6}	1.58E-04	8.18E-06	5.00E-08	1.82E-09	4.95E-11
2^{-7}	9.02E-05	3.74E-06	2.47E-09	1.03E-10	3.44E-12
2^{-8}	4.98E-05	1.91E-06	1.42E-10	6.31E-12	2.69E-12

- Maximum relative error of α -Energy in $[0, T]$. Resolution: 1024

	1/1024	2/1024	4/1024	8/1024	16/1024
2^{-2}	Inf	Inf	Inf	2.91E-04	8.73E-06
2^{-3}	Inf	Inf	3.85E-04	1.49E-05	3.72E-07
2^{-4}	Inf	2.53E-04	2.58E-05	6.85E-07	1.75E-08
2^{-5}	Inf	2.67E-05	1.12E-06	3.38E-08	9.09E-10
2^{-6}	1.46E-05	1.26E-06	4.99E-08	1.82E-09	4.95E-11
2^{-7}	2.53E-06	5.12E-08	2.45E-09	1.03E-10	3.25E-12
2^{-8}	1.06E-06	2.35E-09	1.30E-10	6.31E-12	2.78E-12

Results

- Results of $\alpha^2 \Phi_{\alpha, T}(\eta_{\alpha}^0)$ as $\alpha \rightarrow 0$ for each resolution at $T = 3, 5$.




Figure: Results for $T = 3$ (dotted line) and for $T = 5$ (solid line). Results from runs performed at different resolutions are displayed together: 128^3 (orange plus), 256^3 (cyan star), 512^3 (brown triangles), and 1024^3 (blue squares).

- ▶ Ansatz 4.2 (Larios et al. 2018)

$$\sup_{t \in [0, T^*]} \|\nabla \mathbf{u}_\alpha\|_{L^2} \approx \mathcal{O}(\alpha^p)$$

- ▶ Singularity formation criteria

$$\alpha^2 \Phi_{\alpha, T}(\eta_\alpha^0) > 0 \text{ as } \alpha \rightarrow 0$$

- ▶ Singularity formation occurs if $p \leq -1$.

- ▶ Ansatz 4.2 (Larios et al. 2018)

$$\sup_{t \in [0, T^*]} \|\nabla \mathbf{u}_\alpha\|_{L^2} \approx \mathcal{O}(\alpha^p)$$

- ▶ Singularity formation criteria

$$\alpha^2 \Phi_{\alpha, T}(\boldsymbol{\eta}_\alpha^0) > 0 \text{ as } \alpha \rightarrow 0$$

- ▶ Singularity formation occurs if $p \leq -1$.

- ▶ Ansatz 4.2 (Larios et al. 2018)

$$\sup_{t \in [0, T^*]} \|\nabla \mathbf{u}_\alpha\|_{L^2} \approx \mathcal{O}(\alpha^p)$$

- ▶ Singularity formation criteria

$$\alpha^2 \Phi_{\alpha, T}(\boldsymbol{\eta}_\alpha^0) > 0 \text{ as } \alpha \rightarrow 0$$

- ▶ Singularity formation occurs if $p \leq -1$.

- Results of $\alpha^2 \Phi_{\alpha, T}(\eta_{\alpha}^0)$ as $\alpha \rightarrow 0$. $T=3$.




Figure: Log-log plot of $\max_{t \in [0, T^*]} \|\nabla \mathbf{u}_{\alpha}\|_{L^2}^2$ vs α at $T = 0$ (blue) to $T = 5.0$ (red). $C\alpha^P$ (black)

- Results of $\alpha^2 \Phi_{\alpha, T}(\eta_{\alpha}^0)$ as $\alpha \rightarrow 0$. $T=3$.




Figure: Log-log plot of $\max_{t \in [0, T^*]} \|\nabla \mathbf{u}_{\alpha}\|_{L^2}^2$ vs α at $T = 0$ (blue) to $T = 5.0$ (red). $C\alpha^P$ (black)

- Results of $\alpha^2 \Phi_{\alpha, T}(\eta_{\alpha}^0)$ as $\alpha \rightarrow 0$. $T=5$.




Figure: Log-log plot of $\max_{t \in [0, T^*]} \|\nabla u_{\alpha}\|_{L^2}^2$ vs α at $T = 0$ (blue) to $T = 5.0$ (red). $C\alpha^P$ (black)

- Results of $\alpha^2 \Phi_{\alpha, T}(\eta_{\alpha}^0)$ as $\alpha \rightarrow 0$.




Figure: Log-log plot of $\max_{t \in [0, T^*]} \|\nabla u_{\alpha}\|_{L^2}^2$ vs α at $T = 0$ (blue) to $T = 5.0$ (red). $C\alpha^P$ (black)

- Results of $\alpha^2 \Phi_{\alpha, T}(\eta_{\alpha}^0)$ as $\alpha \rightarrow 0$.




Figure: Plot of the slope of $\max_{t \in [0, T^*]} \|\nabla \mathbf{u}_{\alpha}\|_{L^2}^2$ vs time.

- Therefore $p = -0.417$.
- Our pre-optimization results do not indicate that singularity formation will occur.

- Results of $\alpha^2 \Phi_{\alpha, T}(\eta_{\alpha}^0)$ as $\alpha \rightarrow 0$.




Figure: Plot of the slope of $\max_{t \in [0, T^*]} \|\nabla \mathbf{u}_{\alpha}\|_{L^2}^2$ vs time.

- Therefore $p = -0.417$.
- Our pre-optimization results do not indicate that singularity formation will occur.

- Results of $\alpha^2 \Phi_{\alpha, T}(\eta_{\alpha}^0)$ as $\alpha \rightarrow 0$.




Figure: Plot of the slope of $\max_{t \in [0, T^*]} \|\nabla \mathbf{u}_{\alpha}\|_{L^2}^2$ vs time.

- Therefore $p = -0.417$.
- Our pre-optimization results do not indicate that singularity formation will occur.

- Maximization of $\Phi_{\alpha,T}(\eta_{\alpha}^{(n)})$ vs n . Resolution: 128, $T=3$.




Figure: $\alpha : 16/1024$ (purple), $\alpha : 8/1024$ (red), $\alpha : 4/1024$ (green), $\alpha : 2/1024$ (orange), $\alpha : 1/1024$ (blue).

- Maximization of $\Phi_{\alpha,T}(\eta_{\alpha}^{(n)})$ vs n . Resolution=128, $T=5$.




Figure: $\alpha : 16/1024$ (purple), $\alpha : 8/1024$ (red), $\alpha : 4/1024$ (green), $\alpha : 2/1024$ (orange), $\alpha : 1/1024$ (blue).

- Results of $\alpha^2 \Phi_{\alpha, T}(\tilde{\eta}_{\alpha, T})$ as $\alpha \rightarrow 0$. Resolution=128.




Figure: Results for $T = 3$ (dotted line) and for $T = 5$ (solid line). Results from runs performed at 128^3 (orange plus).

- Results of $\alpha^2 \Phi_{\alpha, T}(\tilde{\eta}_{\alpha, T})$ as $\alpha \rightarrow 0$. Resolution=128, T=3.




Figure: Log-log plot of $\max_{t \in [0, T^*]} \|\nabla u_{\alpha}\|_{L^2}^2$ vs α at $T = 0$ (blue) to $T = 5.0$ (red). $C\alpha^p$ (black)

- Results of $\alpha^2 \Phi_{\alpha, T}(\eta_{\alpha}^0)$ as $\alpha \rightarrow 0$. Resolution=128, T=3.




Figure: Plot of the slope of $\max_{t \in [0, T^*]} \|\nabla \mathbf{u}_{\alpha}\|_{L^2}^2$ vs time.

- Therefore $p = -0.417$.
- Our optimization results do not indicate that singularity formation will occur.

- Results of $\alpha^2 \Phi_{\alpha, T}(\eta_{\alpha}^0)$ as $\alpha \rightarrow 0$. Resolution=128, T=3.




Figure: Plot of the slope of $\max_{t \in [0, T^*]} \|\nabla \mathbf{u}_{\alpha}\|_{L^2}^2$ vs time.

- Therefore $p = -0.417$.
- Our optimization results do not indicate that singularity formation will occur.

- Results of $\alpha^2 \Phi_{\alpha, T}(\eta_{\alpha}^0)$ as $\alpha \rightarrow 0$. Resolution=128, T=3.




Figure: Plot of the slope of $\max_{t \in [0, T^*]} \|\nabla \mathbf{u}_{\alpha}\|_{L^2}^2$ vs time.

- Therefore $p = -0.417$.
- Our optimization results do not indicate that singularity formation will occur.

- Results of $\alpha^2 \Phi_{\alpha, T}(\eta_{\alpha}^0)$ as $\alpha \rightarrow 0$. Resolution=128, T=5.




Figure: Plot of the slope of $\max_{t \in [0, T^*]} \|\nabla u_{\alpha}\|_{L^2}^2$ vs time.

- Therefore $p = -0.417$.
- Our optimization results do not indicate that singularity formation will occur.

- Results of $\alpha^2 \Phi_{\alpha, T}(\eta_{\alpha}^0)$ as $\alpha \rightarrow 0$. Resolution=128, T=5.




Figure: Plot of the slope of $\max_{t \in [0, T^*]} \|\nabla u_{\alpha}\|_{L^2}^2$ vs time.

- Therefore $p = -0.417$.
- Our optimization results do not indicate that singularity formation will occur.

- Results of $\alpha^2 \Phi_{\alpha, T}(\eta_{\alpha}^0)$ as $\alpha \rightarrow 0$. Resolution=128, T=5.




Figure: Plot of the slope of $\max_{t \in [0, T^*]} \|\nabla u_{\alpha}\|_{L^2}^2$ vs time.

- Therefore $p = -0.417$.
- Our optimization results do not indicate that singularity formation will occur.

- ▶ Our results indicate ...
- ▶ Repeat with higher resolutions.
- ▶ The optimization problem is non-convex. The TGV may not optimize to global maxima. We must expand to other initial conditions.

- ▶ Our results indicate ...
- ▶ Repeat with higher resolutions.
- ▶ The optimization problem is non-convex. The TGV may not optimize to global maxima. We must expand to other initial conditions.

- ▶ Our results indicate ...
- ▶ Repeat with higher resolutions.
- ▶ The optimization problem is non-convex. The TGV may not optimize to global maxima. We must expand to other initial conditions.

Thank You