

# Generalized Linear Mixed Models and Practical Applications in R

Sydney Benson, Davita Blyakher, Christina Knudson, Ph.D.

University of St. Thomas

Rose-Hulman Undergraduate Mathematics Conference

# Overview

Generalized linear mixed models can overcome limitations of other modeling techniques

- Response distribution
- Correlated data

Parallel computing can increase the speed of computation

# Linear Models

Assumptions:

- Independent responses
- Normally distributed responses
- Responses have equal variances

## Linear Model Issues

What if our responses are not normally distributed?

- Log odds of your favorite sports team winning a game (binomial)
- Log mean number of students per class at your university (Poisson or negative binomial)

## Linear Model Issues

What if our responses are not normally distributed?

- Log odds of your favorite sports team winning a game (binomial)
- Log mean number of students per class at your university (Poisson or negative binomial)

Use a generalized linear model (GLM)!

## Linear Model Issues

What if our responses are correlated?

- Repeated measurements on individuals
- Measurements on siblings, parents, relatives

## Linear Model Issues

What if our responses are correlated?

- Repeated measurements on individuals
- Measurements on siblings, parents, relatives

Use a linear mixed model (LMM)!

- Differences between “grouped” data are “fixed” effects
- Differences within “grouped” data are “random” effects

# Linear Mixed Models

What are "random" effects?

Random variables, usually normally distributed with a mean 0

Random effects are non-observable, but can be estimated by parameters

Variance component(s): variance(s) of random effects

Parameters for linear mixed models:

- Fixed effects
- Variance components



# Generalized Linear Mixed Models

What if responses are not normally distributed and correlated?

Use a generalized linear mixed model (GLMM)!

- Combination of GLM and LMM
- Include random effects to account for correlation
- Model log odds or log mean

## Salamander Example



Salamanders: single species from 2 different locations

- Rough Butt (R)
- White Side (W)

Do salamanders prefer to mate with others from same location?

## Salamander Example

Correlated: each salamander was mated with multiple others

Unmeasurable: salamanders have personalized tendencies to mate

Assumption: each salamander's tendency is independent

## Salamander Example

What affects the probability that a pair of salamanders will mate?

- Type of cross (RR, RW, WR, WW)
- The female's mating tendency
- The male's mating tendency

## Salamander Example

What affects the probability that a pair of salamanders will mate?

- Type of cross (RR, RW, WR, WW)
- The female's mating tendency
- The male's mating tendency

Fixed effect: type of cross

Random effect: salamander mating tendencies

## Salamander Example

Translation to statistical modeling:

- Response: whether or not the pair mated
- Fixed effects:  $\beta_{RR}$ ,  $\beta_{RW}$ ,  $\beta_{WR}$ ,  $\beta_{WW}$  (log odds of mating)
- Random effects: each salamander (independent, normal)
- Variance components:  $\sigma_F^2$  and  $\sigma_M^2$

## Using glmm

```
> sal <- glmm(Mate ~ 0 + Cross,  
  random = list( ~ 0 + Female, ~ 0 + Male ),  
  varcomps.names = c( "F" , "M" ),  
  data = salamander, m = 10^4,  
  family.glmm = bernoulli.glmm)
```

## Using glmm

Fixed Effects:

	Estimate	Std. Error	z value	Pr(> z )	
CrossR/R	1.4629	0.2720	5.378	7.53e-08	***
CrossR/W	0.3781	0.2527	1.496	0.134612	
CrossW/R	-1.7398	0.3157	-5.512	3.55e-08	***
CrossW/W	1.0345	0.2683	3.857	0.000115	***



## Using glmm

We can translate the log odds back to probabilities:

$$P(\text{mating}) = \frac{\exp(\hat{\beta}_{RW})}{1 + \exp(\hat{\beta}_{RW})}$$

Cross	RR	WW	RW	WR
Probability of mating	0.812	0.798	0.584	0.149

# Parallel Computing

1 person doing 12 calculations

**versus**

4 people doing 3 calculations each

Splitting the computation work of a function amongst the cores available in the computing device

Most computers have 4 or more cores

Leads to increased computation speed

# Parallel Computing

How can we do this in R?

The R package `parallel`

- Detects cores
- Divides likelihood approximation calculations amongst cores
- Sums results from each core
- Returns likelihood approximation

Thank you!

`bens0104@stthomas.edu`

`https://github.com/bensonsyd`

`https://www.linkedin.com/in/bensonsyd`

## References

Knudson C. (2015). *glmm: Generalized Linear Mixed Models via Monte Carlo Likelihood Approximation*. R package version 1.0.2, URL <http://CRAN.R-project.org/package=glmm>.

Knudson C. (2016). *Monte Carlo Likelihood Approximation for Generalized Linear Mixed Models*. Ph.D. Thesis, University of Minnesota.

Ripley B., Tierney L., Urbanek S. (2017). *Package 'parallel'* R package version 3.3.1, URL <http://stat.ethz.ch/R-manual/R-devel/library/parallel/doc/parallel.pdf>.

## Using glmm

### Notes:

- 0+Cross produces log odds for each group, without using reference group.
- 0+Female centers random effects for females at 0, likewise for males.
- Bigger m gives better estimates but takes more time.