## APPLICATIONS OF MATHEMATICS IN FINANCE

CORBIN APPLE, BRIDGET HYLAND, BEN STERLING

- 1. Introduction. It has been well established by The United States Environmental Protection Agency and other independent organizations that average temperature is increasing across the country. This study examines monthly temperature data from two stations: Rye Patch Dam, Nevada, and Salt Lake City International Airport, Utah. These stations were chosen because they are roughly at the same latitude,  $40.498^{\circ}N$  and  $40.790^{\circ}N$ , roughly at the same elevation, 1260.3m and 1287.8m, relatively close longitudes,  $118.316^{\circ}$ W and  $111.980^{\circ}$ W, for Rye Patch and Salt Lake City respectively. The goal of the first hypothesis is to examine whether the effects of climate change statistically differ between these two locations; if they do, we can infer the climate change is predominantly man-made.
  - 2. First Hypothesis. abc
  - 3. Hypothesis 1. def

```
library(ggpubr);
library(ggplot2);
library(latex2exp);
# Disable pdf generation:
pdf(NULL);
data_Salt_Lake = read.csv("GSOY/GSOM_Salt_Lake.csv"); #1287.8m elevation
data_rye_patch = read.csv("GSOY/GSOM_rye_patch.csv"); #1260.3m elevation
Salt_Lake_Temp_Data = data_Salt_Lake$TAVG; rye_patch_Temp_Data = data_rye_patch$TAVG;
  \# \ Calculate \ means: (must \ round \ to \ first \ decimal \ place \ because \ this \ is \ the \ precision \ of \ our \ data: Salt_Lake_mean = round(10 * mean(Salt_Lake_Temp_Data[1:258], na.rm=TRUE)) / 10; rye_patch_mean = round(10 * mean(rye_patch_Temp_Data[151:415], na.rm=TRUE)) / 10; 
Salt_Lake_Diff = Salt_Lake_Temp_Data[259:length(Salt_Lake_Temp_Data)] - Salt_Lake_mean; rye_patch_Diff = rye_patch_Temp_Data[416:length(rye_patch_Temp_Data)] - rye_patch_mean;
# Remove missing data (values of NA) here:
Salt_Lake_Diff = Salt_Lake_Diff[!is.na(Salt_Lake_Diff)];
rye_patch_Diff = rye_patch_Diff[!is.na(rye_patch_Diff)];
                                                                variance, and sample size for each dataset:
# Print mean, variance, and sample size for each dataset:
print("Salt_Lake");
print(paste("\U03BC", "=", mean(Salt_Lake_Diff, na.rm=TRUE)));
print(paste("\U03C3", "=", sqrt(var(Salt_Lake_Diff, na.rm=TRUE))));
print(paste("n.=",length(Salt_Lake_Diff)));
 # New line cat("\n");
print( rye-patch ');
print(paste("\U03BC", "=", mean(rye-patch_Diff, na.rm=TRUE)));
print(paste("\U03C3", "=", sqrt(var(rye_patch_Diff, na.rm=TRUE))));
print(paste("n_=",length(rye_patch_Diff)));
# New line:
# New line:
cat("\n");
# Perform the Shapiro Tests:
shapiro.test(Salt_Lake_Diff);
shapiro.test(rye_patch_Diff);
# Visualize with QQ plots:
png(file = "Salt_Lake_Diff_QQ_Plot.png");
qqplot(y=Salt_Lake_Diff, x=1:length(Salt_Lake_Diff), xlab="x_index", ylab="Data_value",
main=TeX("Salt_Lake_City_Temperature_Increases:_$p_\\approx_1.084_\\times_10^{-12}\$"));
abline(a = min(Salt_Lake_Diff), b = (max(Salt_Lake_Diff) - min(Salt_Lake_Diff)) / length(Salt_Lake_Diff));
dev.off();
\begin{array}{lll} png\left( file = "Rye\_Patch\_Diff\_QQ\_Plot.png" \right); \\ qqplot\left( y=rye\_patch\_Diff \right), & x=1:length\left( rye\_patch\_Diff \right), & xlab="x\_index" , & ylab="Data\_vamain=TeX("Rye\_Patch\_Temperature\_Increases: $$p_\\arrowvert = $$p_\arrowvert = 
 \mathbf{abline}(\texttt{a} = \mathbf{min}(\texttt{rye\_patch\_Diff}), \texttt{ b} = (\mathbf{max}(\texttt{rye\_patch\_Diff}) - \mathbf{min}(\texttt{rye\_patch\_Diff})) \texttt{ / length}(\texttt{rye\_patch\_Diff}));
#Perform two sided t-test with unequal variances:
t.test(rye_patch_Diff, Salt_Lake_Diff);
```