



Busy, Not Bountiful

Validating the Campus Apiary's Pollinator Index Against What's Actually Visiting the Garden Beds

Thandiwe Solberg · Osei Vandermeer — School of Continuous Improvement

Background

Sustainability reports biodiversity progress through the campus apiary's Pollinator Index, built entirely from six smart hives' weekly weight gain. Nobody has checked whether a heavier hive means a garden bed is doing any better by its wild visitors.

Aims


- Test whether the Pollinator Index predicts non-*Apis* richness in nearby beds
- Compare wild-pollinator richness near hives against matched far beds
- Establish whether the Index can carry the biodiversity claim made for it

Methods

- 6 smart hives · weekly weight logs · 14-week spring season
- 3 near ($\leq 50\text{m}$) + 3 far ($>150\text{m}$) beds · fortnightly 10-min timed transects · non-*Apis* species counts
- Observers blind to hive-weight data; Index formula unchanged from Sustainability's own

Results

The Pollinator Index rose 38% across the season. Near-hive richness showed no reliable relationship to weekly index value ($r = 0.07$) and sat persistently below far-bed richness throughout.



Near-hive richness held a median of 3 species (IQR 2–4) against 6 (IQR 5–7) at far beds every fortnight of the Sept–Dec 2025 season, while the Index it's meant to justify climbed 38% over the same period.

Discussion & conclusions

The Index, built from hive weight alone, predicts nothing about the pollinators it's cited to protect. The near/far gap was already there in week one, before the Index had moved. Next: a floral-resource survey before the Index anchors autumn's sustainability comms.

References

1. Ropars, L., Dajoz, I., Fontaine, C., Muratet, A., & Geslin, B. (2019). Wild pollinator activity negatively related to honey bee colony densities in an urban context. *PLOS ONE*, 14(9), e0222316. doi:10.1371/journal.pone.0222316
2. MacInnis, G., Normandin, É., & Zitter, C. D. (2023). Decline in wild bee species richness associated with honey bee (*Apis mellifera* L.) abundance in an urban ecosystem. *PeerJ*, 11, e14699. doi:10.7717/peerj.14699
3. Catania, P., & Vallone, M. (2020). Application of a precision apiculture system to monitor honey daily production. *Sensors*, 20(7), 2012. doi:10.3390/s20072012
4. Roy, H. E., Baxter, E., Saunders, A., & Pocock, M. J. O. (2016). Focal plant observations as a standardised method for pollinator monitoring: opportunities and limitations for mass participation citizen science. *PLOS ONE*, 11(3), e0150794. doi:10.1371/journal.pone.0150794
5. Beng, C., Okwuosa, S., & Solberg, T. (2026). Tapped, Not Filled: A Field Audit of Hydration-Station Activations Against the Living Dashboard's Bottles-Saved Tally. Slop University. doi:10.5555/slop.gylqjf

Insect handling limited to visual identification; no specimens collected. Hive inspections followed routine apiary welfare protocol.

Office of Research Outputs · slop.university



“The index climbed all season. The garden beds nearby didn’t.”

Near-hive richness held at three species while the Pollinator Index rose 38% over 14 weeks.