


PLASMA METABOLOMICS AND LIPIDOMICS REVEALS FED SAMPLING IS SUPERIOR TO FASTED FOR EARLY DETECTION OF DIABETES IN NILE RAT MODEL



Benton J. Anderson¹, Huishi Toh³, Katherine A. Overmyer², Peng Jiang⁴, Joshua J. Coon^{1,2}

¹University of Wisconsin-Madison, Madison, WI, USA; ²Morgridge Institute for Research, Madison, WI, USA; ³University of Santa Barbara, Santa Barbara, CA, USA; ⁴Cleveland State University, Cleveland, OH, USA

BACKGROUND

FASTED VERSUS FED PLASMA METABOLITES

Nile rat animal model for type 2 diabetes

Nile rats are an emerging animal model of type 2 diabetes. Diabetic disease progression and risk indicators are more similar to humans than other rodent models, such as increased blood pressure, increased triacylglycerols, and hyper insulinemia. Also, onset of diabetes can be induced readily and at a young age with a high fat diet.



Question #1: Is collecting blood during random feeding a viable alternative to fasting?

Because diabetes onset occurs rapidly and at a young age in Nile rats, it is critical to sample blood plasma early and often. Sampling fasted rats causes high stress and therefore confounds analysis. We aim to assess the repeated-measures variability of sampling blood under random feeding via bulk phenotypes (random-fed blood glucose) and by using liquid chromatography-mass spectrometry (LCMS) metabolomics and lipidomics to measure plasma metabolites.




Question #2: Do plasma metabolites show different predictive ability of diabetes in Fed versus Fasted?

To understand differences in plasma metabolites that correlate to diabetes between fasted and fed sampling, we constructed statistical models that predict diabetic status of Nile rats, then used feature importance metrics from these models to propose metabolites that best correlate with diabetes.

METHODS & STUDY DESIGN




FASTED VERSUS FED REPRODUCIBILITY




In 50% of Nile rats, the median metabolite variance is lower in non-fasted plasma. This goes against conventional wisdom that eating induces high variability in plasma metabolites.

Key stat:
60%
of plasma metabolites show better reproducibility when Nile rats are fed



MOLECULAR CORRELATIONS TO DIABETES



CONCLUSIONS

Fed sampling has superior repeatability versus fasted

- Lower variance from fed samples ensures better repeatability of plasma metabolite and lipid analysis
- Avoiding fasting will induce less stress on animals and minimize confounding effects

Fed sampling reveals more metabolites that show effects of diabetes

- Fasting minimizes the presence or differential expression of metabolites such as glucose or triacylglycerols in plasma, obscuring important metabolic changes that occur in diabetes

REFERENCES

(1) Refinetti, R. The Nile Grass Rat as a Laboratory Animal. Lab Anim 33, 54–57 (2004). <https://doi.org/10.1038/labana1004-54>

(2) Hutchins, P. D.; Russell, J. D.; Coon, J. J. LipiDex: An Integrated Software Package for High-Confidence Lipid Identification. Cell Systems, 2018, 6, 621–625.e5. <https://doi.org/10.1016/j.cels.2018.03.011>.

We thank the University of Wisconsin-Madison and the National Institutes of Health for support. Work in this project was supported by the National Institute of General Medical Sciences of the National Institutes of Health under Award P41GM108538. Benton Anderson acknowledges support of the NLM training grant 5T15LM007359.