



Ben Trengrove

Your test results

Well Man Check Premium

Summary

Congratulations on taking the Well Man Check Premium which puts you in control of your health data!



i-screen

Collection Date: 26 Mar 2026

Full Blood Count with differential

Your blood counts are all within the normal range and don't indicate anaemia or recent infection.

RDW 12.6 %

Red cell distribution width (RDW) is a calculation of the variation in the size of your RBCs. In some anaemias, such as pernicious anaemia (due to vitamin B12 deficiency), the amount of variation in RBC size causes an increase in the RDW.


[results history](#)Red Blood Cell Count 5.56 x 10¹²/L

Responsible for carrying oxygen around the body. A high count can increase the risk of heart attack or stroke, whilst a low count can mean your body isn't getting the oxygen it needs.

[results history](#)

Haemoglobin 166 g/L

A good measure of your blood's ability to carry oxygen throughout your body. Elevated haemoglobin can be an indicator of lung disease, whilst a low result indicates anaemia.

[results history](#)

Haematocrit 0.48 L/L

A measure of the percentage of red blood cells in the total blood volume. Elevated haematocrit can increase the risk of heart attack or stroke.

[results history](#)

MCV 86 fL

Mean corpuscular volume (MCV) is a measure of the

86

average size of the RBCs. The MCV is elevated when RBCs are larger than normal, eg in anaemia caused by vitamin B12 deficiency. When MCV is decreased, RBCs are smaller than normal as seen in iron deficiency anaemia.

[results history](#)



MCH 30 pg

Mean corpuscular haemoglobin (MCH) is a calculation of the average amount of oxygen-carrying haemoglobin inside a red blood cell. Large RBCs are large tend to have a higher MCH, while small red cells would have a lower value.

[results history](#)



White Blood Cell Count $7.0 \times 10^9/L$

Responsible for fighting infection. A high count can indicate recent infection and even stress, whilst a low count can result from vitamin deficiencies, liver disease and immune diseases.

[results history](#)



Basophils $0.0 \times 10^9/L$

Basophils are a type of white blood cell. Basophils can increase in cases of leukaemia, long-standing inflammation and hypersensitivity to food.


[results history](#)



Eosinophils $0.1 \times 10^9/L$

A type of white blood cell. Can increase in response to allergic disorders, inflammation of the skin and parasitic infections. They can also occur in response to some infections or to various bone marrow malignancies.


[results history](#)



Monocytes $0.9 \times 10^9/L$

A type of white blood cell. Can increase in response to infection as well as inflammatory disorders, and occasionally with some types of leukaemias. Decreased monocyte levels can indicate bone marrow injury or failure and some forms of leukaemia.

[results history](#)



Lymphocytes $2.9 \times 10^9/L$

A type of white blood cell. Can increase with bacterial or viral infection, leukaemia, lymphoma, radiation therapy or acute illness. Decreased lymphocyte levels are common in later life but can also indicate steroid medication, stress, lupus and HIV infection.

[results history](#)



Neutrophils $3.0 \times 10^9/L$

A type of white blood cell. Can increase in response to bacterial infection, inflammatory disease, steroid medication, or more rarely leukaemia. Decreased neutrophil levels may be the result of severe infection or other conditions.


[results history](#)



Platelet Count $258 \times 10^9/L$

Responsible for blood clotting and healing. A high count can indicate a risk of thrombosis, whilst a low count can lead to easy bruising.

[results history](#)



Iron Studies


Your iron study test results are all within the normal range suggesting neither iron deficiency nor overload.

*Note that ferritin levels may be falsely elevated where there is inflammation.

Iron 13 $\mu\text{mol/L}$

An essential trace element is necessary for forming healthy red blood cells and for some enzymes.

[results history](#)



Transferrin 2.8 g/L

A protein that binds iron and transports it around the body (also known as TIBC). High levels indicate iron deficiency.


[results history](#)



Transferrin Saturation 18 %

Low levels typically indicate iron deficiency, and high levels can indicate iron overload.

[results history](#)



Ferritin 305 $\mu\text{g/L}$

The ferritin concentration within the blood stream reflects the amount of iron stored in your body and is reduced in anaemia.

[results history](#)



Male Hormone Panel

Your sex hormones and regulatory hormones are all within the normal range and don't indicate an imbalance.

Testosterone 9.4 nmol/L

Testosterone is an anabolic hormone responsible for bone and muscle strength, as well as mood, energy and sexual function.

[results history](#)



Free Testosterone 229 pmol/L

Most testosterone is strongly bound to sex hormone binding globulin (SHBG). This test measures the proportion of unbound testosterone which is available to the body's tissues. This is usually a calculated value using a formula or an online tool such as <https://www.issam.ch/freetesto.htm>


[results history](#)



SHBG 22 nmol/L

Sex Hormone Binding Globulin (SHBG) is a protein that binds tightly to testosterone and oestradiol. Changes in SHBG levels can affect the amount of hormone that is available to be used by the body's tissues.

[results history](#)



FSH 8.8 U/L

Levels of follicle stimulating hormone in men rise with age, but can also indicate testicular damage and reduced sperm production. Low levels of FSH are detected when men are not producing sperm.

[results history](#)



LH 6.4 U/L

Luteinising hormone is responsible for stimulating testosterone production and sperm generation. Raised LH can signal that the testes are not producing enough testosterone and is relevant when evaluating hypogonadism.

[results history](#)



25-hydroxyvitamin D

Your vitamin D levels are within the normal range and don't indicate a deficiency. Typically we like to see vitamin D levels up above 75 nmol/L.

Vitamin D 99 nmol/L

Although called a vitamin, vitamin D (25-OHD) is actually a steroid hormone which is activated by sunshine on the skin. It is essential for bone strength as it helps the intestines absorb calcium.

[results history](#)



Cholesterol


Your lipid profile is outside of normal range as your LDL "bad" cholesterol levels are elevated, which is a risk factor for cardiovascular disease. As such follow up with your GP is recommended, whilst simple and sustainable lifestyle/dietary changes may assist in bringing these values back down toward normal levels.

Your triglycerides are elevated. The body uses triglycerides and cholesterol differently - triglycerides are a type of fuel, while cholesterol is needed for various metabolic processes such as making particular hormones and building cells. High triglycerides contribute to the development of atherosclerosis (the formation of fatty plaques on blood vessel walls). This may increase the risk of cardiovascular disease. As such follow up with your GP is recommended.

Total Cholesterol 6.6 mmol/l

High total cholesterol is a risk factor for cardiovascular disease.

[results history](#)



LDL 4.3 mmol/l

LDL (low density lipoprotein) cholesterol is often called 'bad cholesterol' because it contributes to plaque, a thick, hard deposit that can clog arteries and make them less flexible.

[results history](#)



HDL 0.95 mmol/l

HDL (high density lipoprotein) cholesterol is often called 'good cholesterol' and is protective against atherosclerosis.

[results history](#)



Triglycerides 3 mmol/l

The main storage form of fatty acids in the body. Elevated triglyceride levels may contribute to hardening of the arteries, and increase the risk of heart disease or stroke.

[results history](#)



Cholesterol Ratio 6.9 mmol/l

A calculation of your HDL cholesterol over your total cholesterol which is used to calculate your likelihood of developing heart disease.

[results history](#)




Blood Glucose

Your fasting glucose is within the normal range and doesn't indicate increased risk of diabetes.

Fasting glucose 5.2 mmol/L

If you have diabetes your body doesn't process glucose effectively.

[results history](#)



Liver Function (with AST)

Your AST and ALT levels are elevated. These enzymes are necessary for amino acid metabolism in the liver and will leak into the bloodstream as the liver becomes inflamed or damaged. Causes of raised AST/ALT may need to be further investigated so it is important to see your GP if you haven't already done so. There is evidence that weight lifting can cause significant increases AST and ALT levels (as well as creatine kinase levels) for up to 7 days post work out.

Bilirubin 6 umol/L

Removed from the body by the liver, and elevated levels may indicate liver disease.


[results history](#)



Alkaline phosphatase (ALP) 55 U/L

Alkaline phosphatase (ALP) is an enzyme located mainly in the liver and the bones. High levels can indicate liver disease.

[results history](#)



GGT 31 U/L

Gamma-glutamyl transferase (GGT) is a liver enzyme which can be used to diagnose alcohol abuse as it is typically raised in long term drinkers.

[results history](#)



AST 123 U/L

Aspartate aminotransferase (AST) is an enzyme created mainly by the liver and the heart. High levels can indicate damage to your liver caused by alcohol, drugs or hepatitis.


[results history](#)



ALT 152 U/L

Alanine aminotransferase (ALT) is an enzyme mainly produced by the liver. A good indicator of liver damage caused by alcohol, drugs or hepatitis.


[results history](#)



Total Protein 72 g/L

A measure of all of the proteins in the plasma portion of your blood. Proteins are important building blocks of all cells and tissues - they are important for body growth and health.


[results history](#)



Albumin 43 g/L

Albumin is a protein which keeps fluid from leaking out of blood vessels, nourishes tissues, and carries hormones, vitamins, drugs, and ions like calcium throughout the body. Albumin is made in the liver and is sensitive to liver damage.


[results history](#)



Globulin 29 g/L

Any of a group of simple proteins found in the blood.

[results history](#)



Inflammation

Your CRP levels are within the normal range and show no sign of inflammation or infection.

C-Reactive Protein 0.8 mg/L

A protein made by the liver and secreted into the blood. It is often the first evidence of inflammation - its concentration increases in the blood within a few hours after the start of inflammatory injury.

[results history](#)



Kidney Function

Your potassium levels are slightly elevated, which may be of no clinical

significance, however this should be closely monitored as small changes in potassium levels may manifest with severe symptoms that should not be ignored such as muscle weakness, fatigue or heart palpitations. In most situations these may be artefactually elevated due to the blood collection process, however a repeat sample would be reassuring.

Your urate levels are slightly above normal - this can cause crystals to form in the joints, which leads to the joint inflammation and pain characteristic of gout. Urate can also form crystals or kidney stones that can damage the kidneys.

Sodium 140 mmol/L

Helps regulate the water and electrolyte balance of your body, and is important in the function of your nerves and muscles. Too much sodium can indicate kidney disease.

[results history](#)



Potassium 5.3 mmol/L

Minor changes in serum potassium can have significant consequences. An abnormal concentration can alter the function of the nerves and muscles for example, the heart muscle may lose its ability to contract.

[results history](#)



Urea 6.3 mmol/L

A high concentration of this waste product can indicate dehydration or that your kidneys aren't working properly.

[results history](#)



Creatinine 99 umol/L

A waste molecule generated from muscle metabolism, and an accurate marker of kidney function.


[results history](#)



Urate 0.49 mmol/L

If too much urate is produced or not enough is excreted, it can accumulate and lead to gout - an inflammation that occurs in joints.

[results history](#)



eGFR 84 mL/min/1.73m2

The estimated glomerular filtration rate (eGFR) measures how well your kidneys filter the wastes from your blood and is the best overall measure of kidney function.

[results history](#)



Recommendations

Find out your Biological Age!

We can calculate your Biological Age using your existing samples!

If you contact us within the next 48hrs we c...

Consider a Kidney Check

Your kidney function may be at risk and I recommend either visiting your GP for further assessment or having your kid...

[Read more](#)

Manage your 'bad' cholesterol

Your lipid profile is abnormal and follow up with your GP for further assessment and to determine whether medication ...

[Read more](#)

Check in with your GP

As always, please visit your GP to discuss your results. Laboratory investigations are an important aspect of healthc...

[Read more](#)

[Read more](#)

Reference Documents

TRENGROVE, BENJAMIN 5.pdf

TRENGROVE, BENJAMIN 1.pdf

TRENGROVE, BENJAMIN 7.pdf

TRENGROVE, BENJAMIN 10.pdf

TRENGROVE, BENJAMIN 6.pdf

TRENGROVE, BENJAMIN 2.pdf

TRENGROVE, BENJAMIN 9.pdf

TRENGROVE, BENJAMIN 3.pdf

TRENGROVE, BENJAMIN 4.pdf

TRENGROVE, BENJAMIN 8.pdf

print results



Return to [dashboard.](#)