

VECTOR DERIVATIVES

Cartesian. $d\mathbf{l} = dx \hat{\mathbf{x}} + dy \hat{\mathbf{y}} + dz \hat{\mathbf{z}}; \quad d\tau = dx dy dz$

Gradient: $\nabla t = \frac{\partial t}{\partial x} \hat{\mathbf{x}} + \frac{\partial t}{\partial y} \hat{\mathbf{y}} + \frac{\partial t}{\partial z} \hat{\mathbf{z}}$

Divergence: $\nabla \cdot \mathbf{v} = \frac{\partial v_x}{\partial x} + \frac{\partial v_y}{\partial y} + \frac{\partial v_z}{\partial z}$

Curl: $\nabla \times \mathbf{v} = \left(\frac{\partial v_z}{\partial y} - \frac{\partial v_y}{\partial z} \right) \hat{\mathbf{x}} + \left(\frac{\partial v_x}{\partial z} - \frac{\partial v_z}{\partial x} \right) \hat{\mathbf{y}} + \left(\frac{\partial v_y}{\partial x} - \frac{\partial v_x}{\partial y} \right) \hat{\mathbf{z}}$

Laplacian: $\nabla^2 t = \frac{\partial^2 t}{\partial x^2} + \frac{\partial^2 t}{\partial y^2} + \frac{\partial^2 t}{\partial z^2}$

Spherical. $d\mathbf{l} = dr \hat{\mathbf{r}} + r d\theta \hat{\boldsymbol{\theta}} + r \sin \theta d\phi \hat{\boldsymbol{\phi}}; \quad d\tau = r^2 \sin \theta dr d\theta d\phi$

Gradient: $\nabla t = \frac{\partial t}{\partial r} \hat{\mathbf{r}} + \frac{1}{r} \frac{\partial t}{\partial \theta} \hat{\boldsymbol{\theta}} + \frac{1}{r \sin \theta} \frac{\partial t}{\partial \phi} \hat{\boldsymbol{\phi}}$

Divergence: $\nabla \cdot \mathbf{v} = \frac{1}{r^2} \frac{\partial}{\partial r} (r^2 v_r) + \frac{1}{r \sin \theta} \frac{\partial}{\partial \theta} (\sin \theta v_\theta) + \frac{1}{r \sin \theta} \frac{\partial v_\phi}{\partial \phi}$

Curl: $\nabla \times \mathbf{v} = \frac{1}{r \sin \theta} \left[\frac{\partial}{\partial \theta} (\sin \theta v_\phi) - \frac{\partial v_\theta}{\partial \phi} \right] \hat{\mathbf{r}} + \frac{1}{r} \left[\frac{1}{\sin \theta} \frac{\partial v_r}{\partial \phi} - \frac{\partial}{\partial r} (r v_\phi) \right] \hat{\boldsymbol{\theta}} + \frac{1}{r} \left[\frac{\partial}{\partial r} (r v_\theta) - \frac{\partial v_r}{\partial \theta} \right] \hat{\boldsymbol{\phi}}$

Laplacian: $\nabla^2 t = \frac{1}{r^2} \frac{\partial}{\partial r} \left(r^2 \frac{\partial t}{\partial r} \right) + \frac{1}{r^2 \sin \theta} \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial t}{\partial \theta} \right) + \frac{1}{r^2 \sin^2 \theta} \frac{\partial^2 t}{\partial \phi^2}$

Cylindrical. $d\mathbf{l} = ds \hat{\mathbf{s}} + s d\phi \hat{\boldsymbol{\phi}} + dz \hat{\mathbf{z}}; \quad d\tau = s ds d\phi dz$

Gradient: $\nabla t = \frac{\partial t}{\partial s} \hat{\mathbf{s}} + \frac{1}{s} \frac{\partial t}{\partial \phi} \hat{\boldsymbol{\phi}} + \frac{\partial t}{\partial z} \hat{\mathbf{z}}$

Divergence: $\nabla \cdot \mathbf{v} = \frac{1}{s} \frac{\partial}{\partial s} (s v_s) + \frac{1}{s} \frac{\partial v_\phi}{\partial \phi} + \frac{\partial v_z}{\partial z}$

Curl: $\nabla \times \mathbf{v} = \left[\frac{1}{s} \frac{\partial v_z}{\partial \phi} - \frac{\partial v_\phi}{\partial z} \right] \hat{\mathbf{s}} + \left[\frac{\partial v_s}{\partial z} - \frac{\partial v_z}{\partial s} \right] \hat{\boldsymbol{\phi}} + \frac{1}{s} \left[\frac{\partial}{\partial s} (s v_\phi) - \frac{\partial v_s}{\partial \phi} \right] \hat{\mathbf{z}}$

Laplacian: $\nabla^2 t = \frac{1}{s} \frac{\partial}{\partial s} \left(s \frac{\partial t}{\partial s} \right) + \frac{1}{s^2} \frac{\partial^2 t}{\partial \phi^2} + \frac{\partial^2 t}{\partial z^2}$

Satisfying the normalization condition (9.139), we find

$$\langle \theta, \phi | l, l \rangle = Y_{l,l}(\theta, \phi) = \frac{(-1)^l}{2^l l!} \sqrt{\frac{(2l+1)!}{4\pi}} e^{il\phi} \sin^l \theta \quad (9.146)$$

• We now apply the lowering operator to determine the remaining spherical harmonics. From Chapter 3 we know that

$$\hat{L}_- |l, m\rangle = \sqrt{l(l+1) - m(m-1)} \hbar |l, m-1\rangle \quad (9.147)$$

Combining (9.146) and (9.147), we find (see Problem 9.18) for $m \geq 0$

$$Y_{l,m}(\theta, \phi) = \frac{(-1)^l}{2^l l!} \sqrt{\frac{(2l+1)(l+m)!}{4\pi(l-m)!}} e^{im\phi} \frac{1}{\sin^m \theta} \frac{d^{l-m}}{d(\cos \theta)^{l-m}} \sin^{2l} \theta \quad (9.148)$$

The choice of the phase factor $(-1)^l$ is taken to ensure that $Y_{l,0}(\theta, \phi)$, which is independent of ϕ , has a real positive value for $\theta = 0$. In fact,

$$Y_{l,0}(\theta, \phi) = \sqrt{\frac{2l+1}{4\pi}} P_l(\cos \theta) \quad (9.149)$$

where $P_l(\cos \theta)$ is the standard **Legendre polynomial**. The spherical harmonics for $m < 0$ are given by

$$Y_{l,-m}(\theta, \phi) = (-1)^m [Y_{l,m}(\theta, \phi)]^* \quad (9.150)$$

It is useful to list the spherical harmonics with $l = 0, 1$, and 2 :

$$Y_{0,0}(\theta, \phi) = \sqrt{\frac{1}{4\pi}} \quad (9.151)$$

$$Y_{1,\pm 1}(\theta, \phi) = \mp \sqrt{\frac{3}{8\pi}} e^{\pm i\phi} \sin \theta \quad (9.152a)$$

$$Y_{1,0}(\theta, \phi) = \sqrt{\frac{3}{4\pi}} \cos \theta \quad (9.152b)$$

$$Y_{2,\pm 2}(\theta, \phi) = \sqrt{\frac{15}{32\pi}} e^{\pm 2i\phi} \sin^2 \theta \quad (9.153a)$$

$$Y_{2,\pm 1}(\theta, \phi) = \mp \sqrt{\frac{15}{8\pi}} e^{\pm i\phi} \sin \theta \cos \theta \quad (9.153b)$$

$$Y_{2,0}(\theta, \phi) = \sqrt{\frac{5}{16\pi}} (3 \cos^2 \theta - 1) \quad (9.153c)$$

Figure 9.11 shows plots of $|Y_{l,m}(\theta, \phi)|^2$ as a function of θ and ϕ . Since the spherical harmonics depend on ϕ through $e^{im\phi}$, these plots are all independent of ϕ . The $l = 0$ state, often called an s state, is spherically symmetric. Thus if a

Table 4.1: Some associated Legendre functions, $P_l^m(\cos \theta)$.

$P_1^1 = \sin \theta$	$P_3^3 = 15 \sin \theta (1 - \cos^2 \theta)$
$P_1^0 = \cos \theta$	$P_3^2 = 15 \sin^2 \theta \cos \theta$
$P_2^2 = 3 \sin^2 \theta$	$P_3^1 = \frac{3}{2} \sin \theta (5 \cos^2 \theta - 1)$
$P_2^1 = 3 \sin \theta \cos \theta$	$P_3^0 = \frac{1}{2} (5 \cos^3 \theta - 3 \cos \theta)$
$P_2^0 = \frac{1}{2} (3 \cos^2 \theta - 1)$	

Notice that l must be a nonnegative integer for the Rodrigues formula to make any sense; moreover, if $|m| > l$, then Equation 4.27 says $P_l^m = 0$. For any given l , then, there are $(2l + 1)$ possible values of m :

$$l = 0, 1, 2, \dots; m = -l, -l + 1, \dots, -1, 0, 1, \dots, l - 1, l. \quad [4.29]$$

But wait! Equation 4.25 is a second-order differential equation: It should have *two* linearly independent solutions, for *any old* values of l and m . Where are all the *other* solutions? *Answer: They exist*, of course, as mathematical solutions to the equation, but they are *physically unacceptable* because they blow up at $\theta = 0$ and/or $\theta = \pi$, and do not yield normalizable wave functions (see Problem 4.4).

Now, the volume element in spherical coordinates⁷ is

$$d^3\mathbf{r} = r^2 \sin \theta \, dr \, d\theta \, d\phi, \quad [4.30]$$

so the normalization condition (Equation 4.6) becomes

$$\int |\psi|^2 r^2 \sin \theta \, dr \, d\theta \, d\phi = \int |R|^2 r^2 \, dr \int |Y|^2 \sin \theta \, d\theta \, d\phi = 1.$$

It is convenient to normalize R and Y individually:

$$\int_0^\infty |R|^2 r^2 \, dr = 1 \quad \text{and} \quad \int_0^{2\pi} \int_0^\pi |Y|^2 \sin \theta \, d\theta \, d\phi = 1. \quad [4.31]$$

The normalized angular wave functions⁸ are called **spherical harmonics**:

⁷See, for instance, Boas, (footnote 2), Chapter 5, Section 4.

⁸The normalization factor is derived in Problem 4.47. The ϵ factor is chosen for consistency with the notation we will be using in the theory of angular momentum; it is reasonably standard, though some older books use other conventions. Notice that

$$Y_l^{-m} = (-1)^m Y_l^m.$$