

1 Some Algebra

Definition: A group is a set G with a binary operation $\circ : G \times G \rightarrow G$ such that,

1. $\forall x, y, z \in G : x \circ (y \circ z) = (x \circ y) \circ z$
2. $\exists e \in G : \forall x \in G : x \circ e = e \circ x = x$
3. $\forall x \in G : \exists x^{-1} \in G : x \circ x^{-1} = x^{-1} \circ x = e$

A group is abelian if $\forall x, y \in G : x \circ y = y \circ x$.

Definition: A ring is a set R with binary operations $+, \cdot$ such that $(R, +)$ is an abelian group such that,

1. $\forall x, y, z \in R : x \cdot (y \cdot z) = (x \cdot y) \cdot z$
2. $\forall x, y, z \in R : x \cdot (y + z) = x \cdot y + x \cdot z$ and $(x + y) \cdot z = x \cdot z + y \cdot z$
3. $\exists 1 \in R : 1 \cdot x = x \cdot 1 = x$

Definition: A field is a commutative division ring meaning that multiplication is commutative and all nonzero elements have inverses.

Definition: An R -module over a ring R is an abelian group M equipped with a bilinear map $R \times M \rightarrow M$ written $(r, m) \mapsto r \cdot m$ such that,

1. $r \cdot (m_1 + m_2) = r \cdot m_1 + r \cdot m_2$
2. $(r_1 + r_2) \cdot m = r_1 \cdot m + r_2 \cdot m$
3. $(r_1 r_2) \cdot m = r_1 \cdot (r_2 \cdot m)$
4. $1_R \cdot m = m$

If R is a field then we call M a k -vectorspace. An R -linear map of R -modules is a group homomorphism $\phi : M_1 \rightarrow M_2$ such that $\phi(r \cdot m) = r \cdot \phi(m)$.

Definition: A k -algebra over a field k is a k -vectorspace V with a bilinear operation $B : V \times V \rightarrow V$. A morphism of k -algebras is a k -linear map $\Phi : V_1 \rightarrow V_2$ preserving the bilinear operation, i.e. $B_2(\Phi(v_1), \Phi(v_2)) = \Phi(B_1(v_1, v_2))$.

Definition: Given a k -algebra V and a basis $\{\mathbf{e}_i\}$ of V the structure constants are coefficients $C_{ijk} \in k$ defined by,

$$B(\mathbf{e}_i, \mathbf{e}_j) = \sum_{k=1}^{\dim V} C_{ijk} \mathbf{e}_k$$

Example 1.1. The complex numbers \mathbb{C} are a 2-dimensional (associative unital commutative division) \mathbb{R} -algebra. We can write $z = x + iy$ and the product is defined,

$$z_1 \cdot z_2 = x_1x_2 - y_1y_2 + i(x_1y_2 + x_2y_1)$$

Thus the structure constants in the basis $\{1, i\}$ are,

$$C_{..1} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \quad C_{..i} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$

Remark 1. Willaim Rowan Hamilton famously searched for a 3-dimensional version of the complex numbers. However he soon discovered the following.

Proposition 1.2. There does not exist a 3-dimensional associative unital \mathbb{R} -algebra with two independent elements which square to -1 .

Proof. Suppose we have $i^2 = j^2 = -1$. By associativity,

$$i \cdot (i \cdot j) = i^2 \cdot j = -j \quad \text{and} \quad (i \cdot j) \cdot j = i \cdot j^2 = -i$$

Now expand,

$$i \cdot j = a + ib + jc$$

However,

$$i \cdot (i \cdot j) = ia - b + (a + ib + jc)c = ac - b + i(a + bc) + jc^2 = -j$$

meaning that $c^2 = -1$ which is impossible. \square

Remark 2. However, all was not lost. He realized that if he added a fourth dimension then the construction would work as long as he gave up the requirement of commutativity leading to the Quaternions.

Example 1.3. The Quaternions \mathbb{H} are a 4-dimensional (associative unital division) \mathbb{R} -algebra,

$$\mathbb{H} = \{w + \mathbf{i}x + \mathbf{j}y + \mathbf{k}z \mid w, x, y, z \in \mathbb{R}\}$$

with biliar multiplication defined by,

Proposition 1.4. We can write any quaternion in vector and scalar part $q = s + \vec{v}$ where $\vec{v} = \mathbf{i}x + \mathbf{j}y + \mathbf{k}z$. Furthermore, define the conjugate $\bar{q} = s - \vec{v}$. Now multiplication becomes,

$$q \cdot q' = ss' - \vec{v} \cdot \vec{v}' + s\vec{v}' + s'\vec{v} + \vec{v} \times \vec{v}'$$

where $\vec{v} \cdot \vec{v}'$ is the dot product of vectors and $\vec{v} \times \vec{v}'$ is the cross product of vectors. Furthermore, we define the norm,

$$|q|^2 = q\bar{q} = w^2 + x^2 + y^2 + z^2$$

which is multiplicative,

$$|qq'|^2 = |q|^2 \cdot |q'|^2$$

Proof. \square

Remark 3. HISTROY OF PRODUCTS

2 Lie Groups and Lie Algebras

Definition: A Lie Group is a group object in the category of smooth manifolds.

Definition: A Lie Algebra is a k -algebra \mathfrak{g} over a field k with its bilinear bracket written as $[\bullet, \bullet] : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$ and satisfying,

1. $[x, x] = 0$
2. $[x, [y, z]] + [y, [z, x]] + [z, [x, y]] = 0$

Example 2.1. The \mathbb{R} -vectorspace $V = \mathbb{R}^3$ with the cross product \times is a Lie algebra. In the standard basis it has structure coefficients ϵ_{ijk} .

Theorem 2.2. Every Lie group G has an associated Lie algebra $\mathfrak{g} = \text{Lie}(G) = T_e G$. Furthermore, if $f : G \rightarrow H$ is a Lie group homomorphism then $df : \mathfrak{g} \rightarrow \mathfrak{h}$ is a Lie algebra homomorphism.

Proof. The Lie bracket is the second-order term in the expansion of $g_1 g_2 g_1^{-1} g_2^{-1}$ about the identity. \square

Example 2.3. The standard example of a Lie group is $\text{GL}(n)$ the group of invertible real $n \times n$ matrices. In particular, for any real vectorspace V then $\text{Aut}(V) = \text{GL}(\dim V)$ is a Lie group. We can write elements of $\text{GL}(n)$ close to the identity as $M = I + \Omega$ for small matrix Ω . So $\mathfrak{gl}(n) = \text{End}(\mathbb{R}^n)$ all $n \times n$ matrices. Then, to second-order,

$$\begin{aligned} M_1 M_2 M_1^{-1} M_2^{-1} &= (I + \Omega_1)(I + \Omega_2)(I - \Omega_1 + \Omega_1^2)(I - \Omega_2 + \Omega_2^2) \\ &= I + \Omega_1 \Omega_2 - \Omega_2 \Omega_1 + O(\Omega^3) = I + [\Omega_1, \Omega_2] + O(\Omega^3) \end{aligned}$$

So the Lie bracket on $\mathfrak{gl}(n) = \text{Lie}(\text{GL}(n))$ is the standard commutator.

2.1 Main Theorems of Lie

Definition: We say a connected topological space is simply connected if every loop may be continuously contracted to a point.

Theorem 2.4. Every finite-dimensional real Lie algebra is $\text{Lie}(G)$ for some simply-connected Lie group G . Furthermore, if $\mathfrak{h} \subset \mathfrak{g}$ is a Lie subalgebra then there exists a unique connected Lie subgroup $H \subset G$ with $\text{Lie}(H) = \mathfrak{h}$.

Theorem 2.5. Let G and H be Lie groups with G simply-connected. Any Lie algebra homomorphism $\phi : \mathfrak{g} \rightarrow \mathfrak{h}$ lifts to a unique Lie group homomorphism $f : G \rightarrow H$ such that $\phi = df$.

2.2 Representations of Lie Groups and Algebras

Definition: A Lie group representation is a vectorspace V and a Lie group homomorphism $G \rightarrow \text{Aut}(V)$. And a representation of Lie Algebras is a Lie algebra homomorphism $\mathfrak{g} \rightarrow \mathfrak{gl}(V)$. That is a linear map $\rho : \mathfrak{g} \rightarrow \text{End}(V)$ which preserves the bracket i.e.

$$\rho([X, Y]) = \rho(X)\rho(Y) - \rho(Y)\rho(X)$$

Proposition 2.6. Every Lie group representation induces a Lie algebra representation. If G is simply connected then every representation of $\mathfrak{g} = \text{Lie}(G)$ gives rise to a unique Lie group representation of G .

Proof. Immediate from the correspondence between homomorphisms of Lie groups and their Lie algebras for simply connected domains. \square

3 Representation Theory of $\text{SU}(2)$ and $\text{SO}(3)$

Definition: The n -sphere is the n -dimensional manifold,

$$S^n = \{\vec{x} \in \mathbb{R}^{n+1} \mid |\vec{x}| = 1\}$$

Remark 4. One of the most important examples of a Lie group is the group $\text{SO}(n)$ of rotations of \mathbb{R}^n or equivalently orientation preserving isometries of S^{n-1} . This group can be constructed as the group of $n \times n$ determinant 1 (special) orthogonal matrices (hence the name). We want to understand this group in detail. First we will start with the simpler question of which spheres can be Lie groups.

DEFINE ISOMETRY GROUPS

Theorem 3.1. S^n can be given a Lie group structure only when $n = 1, 3$.

Definition: We give S^1 a Lie group structure by identifying $S^1 = \{z \in \mathbb{C} \mid |z| = 1\}$. Since $|z \cdot z'| = |z| \cdot |z'| = 1$ and $z\bar{z} = |z|^2 = 1$ this is indeed a group. By analogy, we define a Lie group structure on S^3 by identifying,

$$S^3 = \{q \in \mathbb{H} \mid |q| = 1\}$$

which is a Lie group because $|q \cdot q'| = |q| \cdot |q'| = 1$ and $q\bar{q} = |q|^2 = 1$. From now on, let S^3 refer to this group.

Theorem 3.2. The group S^3 is isomorphic to the group $\text{SU}(2)$.

Proof. \square

Theorem 3.3. There is a 2-fold covering map $\pi : \text{SU}(2) \rightarrow \text{SO}(3)$ making $\text{SU}(2)$ the universal cover of $\text{SO}(3)$.

Proof. We will give two equivalent descriptions of this map in terms of quaternions and complex matrices. This map turns out to be extremely practically useful in physics and physics applications for parametrizing rotations. Therefore, it is worth understanding it in detail. \square

Remark 5. Topologically this is the covering map $\pi : S^3 \rightarrow \mathbb{RP}^3$.

4 The Hopf Fibration

Remark 6. There is a notion of a fibration and of a fibre bundle. In this case they are equivalent and the concept of a fibre bundle is easier to understand.

Definition: A fibre bundle is a morphism $\pi : E \rightarrow B$ with fibre F usually written as

$$F \hookrightarrow E \longrightarrow B$$

is a map such that there are local trivializations, i.e an open cover $U_\alpha \subset B$ with isomorphisms $\varphi_\alpha : \pi^{-1}(U_\alpha) \rightarrow U_\alpha \times F$ such that the following diagram commutes,

$$\begin{array}{ccc} \pi^{-1}(U_\alpha) & \xrightarrow{\varphi_\alpha} & U_\alpha \times F \\ & \searrow \pi & \swarrow \\ & U & \end{array}$$

Definition: A section of a bundle $\pi : E \rightarrow B$ is a morphism $s : B \rightarrow E$ such that $\pi \circ s = \text{id}_B$.

Example 4.1. The trivial bundle with fiber F over X is simply $\pi : X \times F \rightarrow X$ given by projection by the first factor. For example, the trivial bundle of \mathbb{R} over S^1 is a cylinder $\mathbb{R} \times S^1$.

Example 4.2. The Möbius bundle is $\mathbb{R} \hookrightarrow M \rightarrow S^1$ is given by a twist of \mathbb{R} over S^1 such that M is a Möbius strip. M is nontrivial in the sense that $M \not\cong \mathbb{R} \times S^1$ because M does not admit a nonvanishing section.

Remark 7. A fibre bundle $F \hookrightarrow E \rightarrow B$ is morally a way of building E up from fibers F parametrized by the space B along the map $\pi : E \rightarrow B$.

Proposition 4.3. The Hopf Fibration or Hopf Bundle is a fibre bundle $S^1 \hookrightarrow S^3 \rightarrow S^2$ of spheres with the structure map $\pi : S^3 \rightarrow S^2$ defined by $q \mapsto q \cdot \vec{v} \cdot q^{-1}$ for $q \in S^3 \subset \mathbb{H}$ and \vec{v} a fixed unit vector in \mathbb{H} i.e. $\vec{v} \in S^2 \subset \mathbb{R}^3 \subset \mathbb{H}$.

Proof. Suppose that $q_1 \vec{v} q_1^{-1} = q_2 \vec{v} q_2^{-1}$ then $(q_2^{-1} q_1) \vec{v} (q_1 q_2^{-1})^{-1} = \vec{v}$. Thus $(q_2^{-1} q_1)$ maps to a rotation fixing \vec{v} so it must have axis parallel to \vec{v} so $(q_2^{-1} q_1) = e^{a\vec{v}}$ for some $a \in S^1$. Thus $F = \pi^{-1}(\vec{u}) \cong S^1$. The map is a fibre bundle because for sufficiently small open balls $U \subset S^2$ we have $\pi^{-1}(U) \cong U \times S^1$ which is a torus. Thus the Hopf fibration builds S^3 out of interlocking tori. \square

Remark 8. The Hopf fibration is simply the covering map $\pi : S^3 \rightarrow \text{SO}(3)$ composed with the canonical action of $\text{SO}(3)$ on S^2 by rotation.

Remark 9. Steriographic projection.

5 Clifford Algebras

Definition: A quadratic form over a field K is a map $q : V \rightarrow K$ where V is a finite-dimensional K -vectorspace which satisfies,

1. $\forall a \in K, v \in V : q(av) = a^2 q(v)$
2. The map $B(u, v) = q(u + v) - q(u) - q(v)$ is bilinear.

If $\text{char}(K) \neq 2$ then we may define the associated bilinear form,

$$B(u, v) = \frac{1}{2}(q(u + v) - q(u) - q(v))$$

such that $q(v) = B(v, v)$. Thus, for $\text{char}(K) \neq 2$ the theory of quadratic forms is equivalent to that of bilinear forms.

Definition: Let V be a K -vectorspace and $Q : V \rightarrow K$ a quadratic form on V . A Clifford algebra $\text{Cl}(V, Q)$ is a unital associative algebra over K with a linear map $i : V \rightarrow \text{Cl}(V, Q)$ which satisfies,

$$\forall v \in V : i(v)^2 = Q(v) \cdot 1$$

and the following universal property. Let A be any unital associative K -algebra and $j : V \rightarrow A$ any linear map satisfying the condition,

$$\forall v \in V : j(v)^2 = Q(v) \cdot 1_A$$

then there exists a unique K -algebra homomorphism $f : \text{Cl}(V, Q) \rightarrow A$ such that,

$$\begin{array}{ccc} V & \xrightarrow{i} & \text{Cl}(V, Q) \\ & \searrow j & \downarrow f \\ & & A \end{array}$$

commutes i.e. $f \circ i = j$. Thus $\text{Cl}(V, Q)$ is the “freest” such algebra. It is clear that $T(V)/I_Q$ where I_Q is the two-sided ideal of $T(V)$ generated by

$$\{v \otimes v - Q(v) \mid v \in V\}$$

satisfies this universal property and thus $\text{Cl}(V, Q) = T(V)/I_Q$ up to unique isomorphism making this *the* Clifford algebra over V with respect to Q .

Theorem 5.1. There is a functor Cl from the category of K -vectorspaces endowed with quadratic forms to the category of unital associative K -algebras.

Proof. Let (V, Q_V) and (W, Q_W) be vectorspaces paired with quadratic forms and $h : (V, Q_V) \rightarrow (W, Q_W)$ a morphism, that is a K -linear map $h : V \rightarrow W$ such that $Q_W \circ h = Q_V$. Then consider the diagram,

$$\begin{array}{ccc}
V & \xrightarrow{i_V} & \text{Cl}(V, Q_V) \\
\downarrow h & & \downarrow f \\
W & \xrightarrow{i_W} & \text{Cl}(W, Q_W)
\end{array}$$

Because $(i_W \circ h(v))^2 = Q_W(h(v)) \cdot 1_W = Q_V(v) \cdot 1_W$ then the map $i_W \circ h(v)$ factors uniquely through $\text{Cl}(V, Q_V)$ to give a unique map $f : \text{Cl}(V, Q_V) \rightarrow \text{Cl}(W, Q_W)$ lifting h . By the uniqueness of this lift this procedure automatically satisfies functoriality. \square

Definition: Let A be a unital associative K -algebra. An A -representation is a K -vectorspace V with a K -algebra map $\rho : A \rightarrow \text{End}(V)$. Equivalently, a representation V is simply an A -module with K -vectorsapce structure induced by the inclusion $K \rightarrow A$ making V a K -module.

Proposition 5.2. Let $\text{char}(K) \neq 2$ and $Q : V \rightarrow K$ be a quadratic form with associated bilinear form B . Then the Clifford condition is equivalent to,

$$\forall u, v \in V : i(v)i(u) + i(u)i(v) = 2B(v, u) \cdot 1$$

Proof. First, note that $B(v, v) = Q(v)$ and thus setting $u = v$ gives,

$$i(v)^2 + i(v)^2 = 2Q(v) \cdot 1$$

which for $\text{char}(K) \neq 2$ implies the defining condition. Convseresely, if for each $v \in V$ we have,

$$i(v)^2 = Q(v) \cdot 1$$

then consider,

$$i(v+u)^2 = (i(v)+i(u))^2 = i(v)^2 + i(v)i(u) + i(u)i(v) + i(u)^2 = Q(v) \cdot 1 + i(v)i(u) + i(u)i(v) + Q(u) \cdot 1$$

However, we also have,

$$i(v+u)^2 = Q(v+u) \cdot 1$$

which implies that,

$$i(v)i(u) + i(u)i(v) = [Q(v+u) - Q(v) - Q(u)] \cdot 1 = 2B(v, u) \cdot 1$$

\square

Remark 10. From now on we will assume that $\text{char}(K) \neq 2$ and V is a vectorspace of dimension n over K with quadratic form $Q : V \rightarrow K$ and associated bilinear form $B : V \times V \rightarrow K$.

Proposition 5.3. Let $\{e_1, \dots, e_n\}$ be an orthogonal basis for V with respect to B i.e. $B(e_i, e_j) = \delta_{ij}Q(e_i)$. Then $\text{Cl}(V, Q)$ has a basis,

$$\{e_{i_1}e_{i_2}\cdots e_{i_k} \mid 1 \leq i_1 < i_2 < \cdots < i_k \leq n \text{ and } 1 \leq k \leq n\}$$

and thus,

$$\dim_K \text{Cl}(V, Q) = \sum_{k=1}^n \binom{n}{k} = 2^n$$

Proposition 5.4. Any vectorspace over K with a nondegenerate quadratic form $Q : V \rightarrow K$ is isomorphic to K^n with the stanard quadratic form,

$$Q_{p,q}(v) = v_1^2 + \cdots + v_p^2 - v_{p+1}^2 - \cdots - v_{p+q}^2$$

where p is the number of positive eigenvalues and q the number of negative eigenvalues. We then denote the Clifford algebra in this case,

$$\text{Cl}_{p+q}(K) = \text{Cl}(K^n, Q_{p,q}) = \text{Cl}(V, Q)$$

(EXAMPLES)

(PROPOSITIONS)

($\text{Cl}(1,3)(\mathbb{C}) = \text{C04}(\mathbb{C})$ is WICK ROTATION)

5.1 The Pin and Spin Groups

5.2 Representations of Clifford Algebras

6 General Theory of Transformations in Physics

Remark 11. Quantum states are not defined up to scaling which motivates the following definition.

Definition: Let \mathcal{H} be a quantum Hilbert space then the projective space *state space* or *ray space* is $\mathbb{P}(\mathcal{H})$ with the metric,

$$d(\underline{\psi}, \underline{\phi}) = \cos^{-1} \sqrt{\frac{\langle \psi, \phi \rangle \langle \phi, \psi \rangle}{\langle \psi, \psi \rangle \langle \psi, \phi \rangle}}$$

where we define the *ray product*

$$\underline{\psi} \cdot \underline{\phi} = \sqrt{\frac{\langle \psi, \phi \rangle \langle \phi, \psi \rangle}{\langle \psi, \psi \rangle \langle \psi, \phi \rangle}} = \frac{|\langle \psi, \phi \rangle|}{|\psi||\phi|}$$

which is the Fubini-Study metric on \mathbb{CP}^n .

A *symmetry transformation* is a ray transformation, that is a homeomorphism

$$T : \mathbb{P}(\mathcal{H}) \rightarrow \mathbb{P}(\mathcal{H}')$$

which is an isometry i.e. preserves the ray product. Explicitly,

$$T\underline{\Psi} \cdot T\underline{\Phi} = \underline{\Psi} \cdot \underline{\Phi}$$

Theorem 6.1 (Wigner). Any symmetry transformation $T : \mathbb{P}(\mathcal{H}) \rightarrow \mathbb{P}(\mathcal{H}')$ lifts to a \mathbb{R} -linear map $T : \mathcal{H} \rightarrow \mathcal{H}'$ which is either unitary (and thus \mathbb{C} -linear) or anti-unitary (and thus \mathbb{C} -antilinear).

Theorem 6.2. Let $T : \mathbb{P}(\mathcal{H}) \rightarrow \mathbb{P}(\mathcal{H}')$ be a ray transformation and $U, V : \mathcal{H} \rightarrow \mathcal{H}'$ are lifts of T i.e. linear maps compatible with T . Assume that $\dim \mathcal{H} \geq 2$ then $V = Ue^{i\alpha}$ for some $\alpha \in \mathbb{R}$.

Definition: The *symmetry group* of a Hilbert space \mathcal{H} is the group $\Gamma(\mathcal{H})$ of symmetry transformations of the ray space $\mathbb{P}(\mathcal{H}) \rightarrow \mathbb{P}(\mathcal{H})$. A *quantum representation* of a group G on \mathcal{H} is a projective representation $G \rightarrow \text{PAU}(\mathcal{H}) = \Gamma(\mathcal{H})$ by Wigner's theorem.

In general, a quantum field is an operator valued section of the associated bundle $E = P \times_{\rho} \mathbb{P}(V)$ to a projective representation $\rho : G \rightarrow \text{PGL}(V)$ where $P \rightarrow M$ is a principle G -bundle over M (usually the frame bundle or the bundle of spin frames).

7 The Wigner-Eckart Theorem

Lemma 7.1. Let G be a compact Lie group and $\rho : G \rightarrow \text{Aut}(V)$ a G -representation. There is a projection map $s : V \rightarrow V^G$ given by,

$$s(v) = \frac{1}{|G|} \int_{g \in G} \rho(g) \cdot v$$

Proof. First, this map is well-defined because for any $v \in V$ and $g \in G$ we have,

$$\rho(g) \cdot s(v) = \frac{1}{|G|} \int_{h \in G} \rho(g) \cdot \rho(h) \cdot v = \frac{1}{|G|} \int_{h \in G} \rho(gh) \cdot v = \frac{1}{|G|} \int_{h \in G} \rho(h) \cdot v = s(v)$$

Thus $s(v) \in V^G$. Consider the inclusion $\iota : V^G \rightarrow V$. Then,

$$s \circ \iota(v) = \frac{1}{|G|} \int_{g \in G} \rho(g) \cdot v = \frac{1}{|G|} \int_{g \in G} v = v$$

thus $p \circ \iota = \text{id}_{V^G}$. □

Lemma 7.2. Let G be a compact Lie group, V a Hilbert space, and $\rho : G \rightarrow \text{Aut}(V)$ a unitary representation. If V is finite-dimensional then for any G -subrep $W \subset V$ there exists G -equivariant projection $p : V \rightarrow W$ satisfying $p^\dagger = \iota_W$.

Proof. As inner product spaces, there automatically exists a projection $q : V \rightarrow W$ because V is finite-dimensional satisfying $q \circ \iota_W = \text{id}_W$ and $q^\dagger = \iota_W$ i.e. is orthogonal. Then apply the previous lemma to the G -rep $\text{Hom}(V, W)$ to get a morphism $p = s(q) \in \text{Hom}_G(V, W)$. It remains to show that $p \circ \iota_W = \text{id}_W$ and $p^\dagger = \iota_W$. Since ι_W is G -equivariant we have,

$$\begin{aligned} p \circ \iota_W &= \frac{1}{|G|} \int_{g \in G} \rho(g) \circ q \circ \rho(g)^{-1} \circ \iota_W = \frac{1}{|G|} \int_{g \in G} \rho(g) \circ q \circ \iota_W \circ \rho(g)^{-1} \\ &= \frac{1}{|G|} \int_{g \in G} \rho(g) \circ \text{id}_W \circ \rho(g)^{-1} = \text{id}_W \end{aligned}$$

Finally, because $\rho : G \rightarrow \text{Aut}(V)$ is a unitary rep,

$$\begin{aligned} p^\dagger &= \left(\frac{1}{|G|} \int_{g \in G} \rho(g) \circ q \circ \rho(g)^{-1} \right)^\dagger = \frac{1}{|G|} \int_{g \in G} (\rho(g)^{-1})^\dagger \circ q^\dagger \circ \rho(g)^\dagger \\ &= \frac{1}{|G|} \int_{g \in G} \rho(g) \circ q \circ \rho(g)^{-1} = p \end{aligned}$$

□

Lemma 7.3. Let G be a compact Lie group then any finite-dimensional unitary G -representation is decomposable as Hilbert space G -reps into sum of irreducible unitary G -reps.

Proof. Let $\rho : G \rightarrow \text{Aut}(V)$ be the lowest-dimension unitary G -rep which is not decomposable in the desired maner. In particular, V is not irreducible. Thus, there is a nontrivial submodule $W \subset V$ which then has a G -equivariant hermitian projection $p : V \rightarrow W$. Therefore, $\ker p$ and W are orthogonal since,

$$\langle v, w \rangle_W = \langle v, \iota_W w \rangle_W = \left\langle \iota_W^\dagger v, w \right\rangle_V = \langle p(v), w \rangle_V = 0$$

Then there is an exact sequence of $\mathbb{C}[G]$ -modules,

$$0 \longrightarrow \ker p \longrightarrow V \xrightleftharpoons[\iota_W]{p} W \longrightarrow 0$$

Since $p \circ \iota_W = \text{id}_W$ the sequence splits so we find that,

$$V = W \oplus \ker p$$

as G -representations and Hilbert spaces. Since the dimension of V is minimal, these factors must be decomposable in the proper way which implies that V is as well. □

Corollary 7.4. Let G be a compact Lie group. Then any finite-dimensional G -representation is semisimple.

Proof. Apply the unitarian trick: every finite-dimensional G -rep is unitary. To show this consider a finite-dimensional G -rep $\rho : G \rightarrow \text{Aut}(V)$ and any inner product $\langle \bullet, \bullet \rangle$ on V . Then define,

$$\langle v, u \rangle_G = \frac{1}{|G|} \int_{g \in G} \langle \rho(g) \cdot v, \rho(g) \cdot u \rangle$$

which is an inner products and $r\rho : G \rightarrow \text{Aut}(V)$ is unitary with respect to it. □

Definition: We say a connected compact Lie group G has the *Clebsch-Gordon property* if for every pair of finite-dimensional irreducible complex representations V_1 and V_2 then the tensor product representation $V_1 \otimes_{\mathbb{C}} V_2$ is decomposed into *distinct* irreducible G -representations i.e. $V_1 \otimes_{\mathbb{C}} V_2$ does not contain two isomorphic copies of the same irreducible representation.

Lemma 7.5. Let G be a connected compact Lie group and $\rho : G \rightarrow \text{Aut}(\mathcal{H})$ a unitary representation. Let $W_1, W_2 \subset \mathcal{H}$ be irreducible G -submodules. Then either $W_1 \cong W_2$ or $W_1 \perp W_2$.

Proof. There exists a G -invariant orthogonal projection map $p : \mathcal{H} \rightarrow V_2$. Then $p|_{W_1} : W_1 \rightarrow W_2$ gives a G -morphism. By Schur, either $W_1 \cong W_2$ or $p|_{W_1} = 0$. The latter implies that for each $w \in W_1$ we have $p(w) = 0$ i.e. $w \in W_2^\perp$ so $W_1 \perp W_2$. \square

Definition: Let V be a \mathbb{C} -vector space and $\bar{V} = V \otimes_\sigma \mathbb{C}$ where $\sigma : \mathbb{C} \rightarrow \mathbb{C}$ is complex conjugation. The canonical map $\sigma : V \rightarrow \bar{V}$ is then \mathbb{C} -antilinear (real) isomorphism since,

$$\sigma(\alpha v) = \alpha v \otimes 1 = v \otimes \bar{\alpha} = \bar{\alpha} \cdot (v \otimes 1) = \bar{\alpha} \sigma(v)$$

If V is a G -rep then $V \otimes_\sigma \mathbb{C}$ is canonically a G -rep via,

$$\bar{\rho}(g) = \sigma \circ \rho(g) \circ \sigma^{-1}$$

Lemma 7.6. Let V be a finite-dimensional G -rep then,

$$V \text{ is irreducible} \iff \bar{V} \text{ is irreducible} \iff V^* \text{ is irreducible}$$

Proof. If V^* is irreducible take a G -invariant subspace $W \subset V$ and consider,

$$W^\perp = \{\phi \in V^* \mid \forall w \in W : \phi(w) = 0\}$$

This is G -invariant because $\rho^*(g) \cdot \phi = \phi \circ \rho(g^{-1})$ and $\phi(\rho(g^{-1})w) = 0$ since W is G -invariant. Thus, either $W^\perp = (0)$ or $W^\perp = V^*$. In the first case $W = V$ otherwise there would be a nontrivial projection onto its complement sending it to zero, in the second case, $W = (0)$ otherwise the projection onto any vector in W would not vanish on W contradicting $W^\perp = V^*$. Therefore V is irreducible. Furthermore $V^{**} \cong V$ as G -modules so if V is irreducible then so is V^{**} which implies, by the above, that V^* is irreducible.

Now suppose that \bar{V} is irreducible take a G -invariant subspace $W \subset V$ and consider $W' = \sigma(W) \subset \bar{V}$. This is a subspace because it is additive and $\alpha \sigma(W) = \sigma(\bar{\alpha} W) = \sigma(W)$. Since \bar{V} is irreducible $W' = (0)$ or $W' = \bar{V}$. Since σ is an isomorphism this implies that $W = (0)$ or $W' = \bar{V}$ and thus V is irreducible. Furthermore, $\bar{\bar{V}} = V$ so if V is irreducible then so is $\bar{\bar{V}}$ which implies, by the above, that \bar{V} is irreducible. \square

Lemma 7.7. A sesquilinear form on V , equivalently a linear map $S : \bar{V} \otimes_{\mathbb{C}} V \rightarrow \mathbb{C}$ is the same as a linear map $\bar{V} \rightarrow V^*$. If V is a G -rep then a sesquilinear form on V is G -invariant iff the associated linear map $\bar{V} \rightarrow V^*$ is a G -morphism.

Proof. Consider the map $\phi_S : \bar{V} \rightarrow V^*$ via $\phi_S(v \otimes 1) = \langle v, - \rangle$ which is a linear functional on V and thus in V^* . Furthermore ϕ_S is \mathbb{C} -linear since,

$$\phi_S(v \otimes \alpha) = \phi_S(\bar{\alpha} v \otimes 1) = \langle \bar{\alpha} v, - \rangle_S = \alpha \langle v, - \rangle_S = \alpha \phi_S(v)$$

and it is clearly additive. Conversely, any \mathbb{C} -linear map $\phi : \bar{V} \rightarrow V^*$ defined a sesquilinear form via $\langle v, w \rangle_\phi = \phi(\sigma(v))(w)$. This is sesquilinear since,

$$\langle \alpha v + \beta u, w \rangle_\phi = \phi(\sigma(\alpha v + \beta u))(w) = \bar{\alpha}\phi(\sigma(v))(w) + \bar{\beta}\phi(\sigma(u))(w) = \bar{\alpha}\langle v, w \rangle_\phi + \bar{\beta}\langle u, w \rangle_\phi$$

Furthermore, if the sesquilinear map is G -invariant the,

$$\begin{aligned} \phi(\bar{\rho}(g)\bar{v}) &= \langle \rho(g)v, w \rangle = \langle \rho(g)v, \rho(g) \circ \rho(g^{-1})w \rangle = \langle v, \rho^{-1}(g)w \rangle \\ &= \phi(v)(\rho^{-1}(g)w) = [\rho^*(g) \cdot \phi(v)](w) \end{aligned}$$

and thus $\phi \circ \bar{\rho}(g) = \rho^*(g) \circ \phi$. Likewise if ϕ is a G -morphpism then,

$$\begin{aligned} \langle \rho(g) \cdot v, \rho(g) \cdot w \rangle &= \phi(\sigma \circ \rho(g) \cdot v)(\rho(g) \cdot w) = \phi(\bar{\rho}(g) \cdot \sigma(v))(\rho(g) \cdot w) \\ &= [\phi \circ \bar{\rho}(g)](v)(\rho(g) \cdot w) = [\rho^*(g) \circ \phi](\sigma(v))(\rho(g) \cdot w) \\ &= \phi(\sigma(v))(\rho(g^{-1}) \circ \rho(g) \cdot w) = \phi(\sigma(v))(w) = \langle v, w \rangle \end{aligned}$$

and thus $\langle -, - \rangle$ is G -invariant. \square

Lemma 7.8. Let V be a finite dimensional irreducible G -representation then, up to scale, there is a unique G -invariant sesquilinear form on V .

Proof. We know that G -invariant sesquilinear forms are the same as G -morphisms $\bar{V} \rightarrow V^*$ which, since V is irreducible, are irreducible. Therefore, by Schur's lemma, G -morphisms $\bar{V} \rightarrow V^*$ are unique up to scale. \square

Lemma 7.9. Let G be a connected compact Lie group and $\rho : G \rightarrow \text{Aut}(\mathcal{H})$ a unitary representation. Let $W_1, W_2 \subset \mathcal{H}$ be isomorphic irreducible G -submodules. Then, for any isomorphism $f : W_1 \rightarrow W_2$, there exists a constant c such that,

$$\forall w_1, w_2 \in W_1 : \langle w_1, f(w_2) \rangle = c \langle w_1, w_2 \rangle$$

Proof. The sesquilinear form $\langle w_1, f(w_2) \rangle$ is G -invariant because f is a G -morphpism meaning that,

$$\langle \rho(g) \cdot w_1, f(\rho(g) \cdot w_2) \rangle = \langle \rho(g) \cdot w_1, \rho(g) \cdot f(w_2) \rangle = \langle w_1, f(w_2) \rangle$$

where I have used the fact that $\langle -, - \rangle$ is G -invariant because $\rho : G \rightarrow \text{Aut}(V)$ is unitary. Therefore, by the lemma, the G -invariant sesquilinear forms $\langle -, - \rangle$ and $\langle -, f- \rangle$ are equal up to scale. Since $\langle -, - \rangle$ is a Hermitian inner product, $\forall v \neq 0 : \langle v, v \rangle > 0$, so it is nondegenerate (the associted G -morphpism $\bar{V} \rightarrow V^*$ is an isomorphism) which implies that any other sesquilinear form (equivalently isomorphism $\bar{V} \rightarrow V^*$), in particular $\langle -, f- \rangle$, must be a constant multiple of it. \square

Lemma 7.10. Let V be an irreducible G -rep and $f : V \rightarrow W$ a G -morphpism. Then either f is zero or induces an isomorphism onto its image making V a irreducible factor of W .

Proof. Since $\ker f \subset V$ is G -invariant we must have $\ker f = V$ or $\ker f = 0$. In the first case $f = 0$, in the second, $f : V \rightarrow W$ is injective and thus $f : V \rightarrow \text{Im}(f)$ is an isomorphism onto a G -submodule. \square

Theorem 7.11 (Wigner-Eckart). Let G be a connected compact Lie group with the Clebsch-Gordon property. Let \mathcal{H} be a Hilbert space and $\rho : G \rightarrow \text{Aut}(\mathcal{H})$ a unitary representation. Let $V_1, V_2 \subset \mathcal{H}$ be finite-dimensional irreducible G -subrepresentations. Let $\mathcal{O}_1, \mathcal{O}_2 \subset \text{End}(\mathcal{H})$ be finite-dimensional irreducible G -submodules and $f : \mathcal{O}_1 \rightarrow \mathcal{O}_2$ an isomorphism. Then there exists a constant $c_f \in \mathbb{C}$ such that either,

$$\forall v_1 \in V_1, v_2 \in V_2, T \in \mathcal{O}_1 : \langle v_1, f(T)v_2 \rangle = c \langle v_1, Tv_2 \rangle$$

or $\langle v_1, Tv_2 \rangle = 0$ for all $V_1 \in V_1, w_2 \in W_2$, and $T \in \mathcal{O}_1$.

Proof. Consider diagram of G -morphisms,

$$\begin{array}{ccc} \mathcal{O}_1 \otimes V_2 & \xrightarrow{e} & \mathcal{O}_1(V_2) \\ \downarrow f \otimes \text{id} & & \\ \mathcal{O}_2 \otimes V_2 & \xrightarrow{e} & \mathcal{O}_2(V_2) \end{array}$$

By the Clebsch-Gordon property, $\mathcal{O}_1 \otimes V_2$ and $\mathcal{O}_2 \otimes V_2$ decompose into irreducible G -reps with at most one factor of V_1 . Since the evaluation morphism is surjective, this implies that $\mathcal{O}_1(V_2)$ and $\mathcal{O}_2(V_2)$ can have at most one copy of V_1 in their decomposition. If none of its irreducible representations are isomorphic to V_1 then we have seen that the inner products $\langle v_1, Tv_2 \rangle$ vanish. Likewise, if V_1 does not appear in the decomposition of $\mathcal{O}_2(V_2)$ then all $\langle v_1, f(T)v_2 \rangle$ vanish. Therefore, we may assume that both $\mathcal{O}_1(V_2)$ and $\mathcal{O}_2(V_2)$ contain a copy of V_1 which we denote as W_1, W_2 respectively. Restricting the diagram, we find,

$$\begin{array}{ccc} W_1^\otimes \subset \mathcal{O}_1 \otimes V_2 & \xrightarrow{e} & W_1 \subset \mathcal{O}_1(V_2) \\ \downarrow f \otimes \text{id} & & \downarrow \tilde{f} \\ W_2^\otimes \subset \mathcal{O}_2 \otimes V_2 & \xrightarrow{e} & W_2 \subset \mathcal{O}_2(V_2) \end{array}$$

These arrows become isomorphisms since they are nontrivial G -morphisms which allows the construction of \tilde{f} . As we have shown, nonisomorphic irreducible G -submodules are perpendicular so to compute these inner products we need only to consider $Tv_2 \in W_1 \subset \mathcal{O}_1(V_2)$ and $T'v_2 \in W_2 \subset \mathcal{O}_2(V_2)$ since all other components product to zero with $v_1 \in V_1$. Now, consider,

$$f'(Tv_2) = e \circ (f \otimes \text{id}) \circ e^{-1}(Tv_2) = e \circ (f \otimes \text{id})(T \otimes v_2) = e(f(T) \otimes v_2) = f(T)v_2$$

Now choose an isomorphism $g : V_1 \rightarrow W_1$ then $f' \circ g : V_1 \rightarrow W_2$ is an isomorphism. Therefore, by the previous lemma, there exist constants, c_1 and c_2 such that,

$$\forall v_1, v'_2 \in V_1 : \langle v_1, g(v'_2) \rangle = c_1 \langle v_1, v'_2 \rangle \quad \langle v_1, f' \circ g(v'_2) \rangle = c_2 \langle v_1, v'_2 \rangle$$

In particular, choosing $v'_2 = g^{-1}(Tv_2)$ for any $Tv_2 \in W_2$ gives,

$$\langle v_1, Tv_2 \rangle = c_1 \langle v_1, g^{-1}(Tv_2) \rangle \quad \langle v_1, f(T)v_2 \rangle = c_2 \langle v_1, g^{-1}(Tv_2) \rangle$$

which, if $c_1 \neq 0$ taking $c_f = c_2/c_1$ and otherwise $\langle v_1, Tv_2 \rangle = 0$ proves the theorem. \square

Remark 12. In practice, if we have a G -representation on a set of operators this theorem reduces computing $\langle v_1, Tv_2 \rangle$ for each operator and vector in the representation, to computing a single constant as long as we know these inner products for any other set of operators in the same representation. In fact, we have shown that these inner products are up to scale the same as the overlap coefficients in the tensor product decomposition.

8 Proof of the Main Theorem

8.1 Setup

Consider the Hilbert space \mathcal{H} of our possibly interacting QFT which has unitary projective representation of the Poincare group or equivalently a unitary representation of its double cover. This representation gives a representation of the Lie algebra which has generators,

$$\hat{P}_i \quad \hat{K}_i \quad \hat{J}_i \quad (1a)$$

acting as operators on \mathcal{H} and satisfying the commutation relations,

$$[\hat{J}_i, \hat{J}_j] = i\epsilon_{ijk}\hat{J}_k \quad (2a)$$

We assume there is an (interacting) vacuum state $|\Omega\rangle$ is invariant under the action of the Poincaré group (AND UNDER CPT?). A particle is given by a multiplet of field operators $\hat{\phi}_i(x)$ in a finite-dimensional irreducible $\text{Spin}(1,3)$ -representation $S : \text{Spin}(1,3) \rightarrow W$. Explicitly, this multiplet transforms under a Lorentz transformation Λ (or more generally an element of $\text{Spin}(1,3)$) via,

$$U(\Lambda)\hat{\phi}_i(x)U(\Lambda)^{-1} = S_{ij}(\Lambda)\hat{\phi}_j(\Lambda^{-1}x) \quad (3a)$$

where $\Lambda \in \text{Spin}(1,3)$ acts on x via the covering projection $\pi : \text{Spin}(1,3) \rightarrow \text{SO}(1,3)$ which is thus acts as a matrix. Note that the representation W must be of the form (μ, ν) by the classification of projective $\text{SO}(1,3)$ -reps and cannot be unitary because W is finite dimensional. Furthermore, the Poincare group translations act via,

$$e^{i\hat{P}_\mu x^\mu} \hat{\phi}(y^\mu) e^{-i\hat{P}_\mu x^\mu} = \hat{\phi}(x^\mu + y^\mu)$$

8.2 The Källén-Lehmann Spectral Resolution

We must compute the two-point correlation functions,

$$\langle \Omega | \hat{\phi}_i(x) \hat{\phi}_j^\dagger(y) | \Omega \rangle \quad (4a)$$

$$\langle \Omega | \hat{\phi}_j^\dagger(y) \hat{\phi}_i(x) | \Omega \rangle \quad (4b)$$

We now use a resolution of unity. However, since $\hat{\phi}$ is a charged field, by selection rules, we need only consider single particle states in the resolution,

$$\langle \Omega | \hat{\phi}_i(x) \hat{\phi}_j^\dagger(y) | \Omega \rangle = \sum_{\lambda, s} \int_{p^2=m_\lambda^2} \frac{d^3p}{(2\pi)^3} \frac{1}{2p_0} \langle \Omega | \hat{\phi}_i(x) | \lambda_{p,s} \rangle \langle \lambda_{p,s} | \hat{\phi}_j^\dagger(y) | \Omega \rangle \quad (5a)$$

where $|\lambda_s\rangle$ is a W -multiplet of single particle rest states of mass m_λ and $|\lambda_{p,s}\rangle = U(B_p) |\lambda_s\rangle$ where B_p is the Lorentz transformation which performs a boost to momentum p . Since $\hat{P}_i |\lambda_s\rangle = 0$ and $[\hat{P}_i, U(B_p)] = p$, (WHY?) Therefore,

$$\langle \Omega | \hat{\phi}_i(x) | \lambda_{p,s} \rangle = \langle \Omega | e^{i\hat{P}_\mu x^\mu} \hat{\phi}_i(0) e^{-i\hat{P}_\mu x^\mu} | \lambda_{p,s} \rangle \quad (6a)$$

$$= \langle \Omega | \hat{\phi}_i(0) | \lambda_{p,s} \rangle e^{-ip \cdot x} \quad (6b)$$

using the fact that $e^{i\hat{P}_\mu x^\mu} |\lambda_{p,s}\rangle = e^{ip_\mu x^\mu} |\lambda_{p,s}\rangle$ and the vacuum is invariant under the Poincare group. Now,

$$\langle \Omega | \hat{\phi}_i(x) | \lambda_{p,s} \rangle = \langle \Omega | \hat{\phi}_i(0) U(B_p) | \lambda_s \rangle e^{-ip \cdot x} \quad (7a)$$

$$= \langle \Omega | \hat{\phi}_i(0) U(B_p) | \lambda_s \rangle e^{-ip \cdot x} \quad (7b)$$

Again, using the fact that $|\Omega\rangle$ is Poincaré invariant we find,

$$\langle \Omega | \hat{\phi}_i(x) | \lambda_{p,s} \rangle = \langle \Omega | U^\dagger(B_p) \hat{\phi}_i(0) U(B_p) | \lambda_s \rangle e^{-ip \cdot x} \quad (8a)$$

$$= S_{ik}(B_p^{-1}) \langle \Omega | \hat{\phi}_k(0) | \lambda_s \rangle e^{-ip \cdot x} \quad (8b)$$

Furthermore, since $\hat{\phi}_k$ and $|\lambda_s\rangle$ are both W -multiplets, by the Wigner-Ekart theorem we have,

$$\langle \Omega | \hat{\phi}_k(0) | \lambda_s \rangle = C_\lambda \delta_{ks} \quad (9)$$

for some constant $C_\lambda \in \mathbb{C}$. Thus,

$$\langle \Omega | \hat{\phi}_i(x) | \lambda_{p,s} \rangle = C_\lambda S_{is}(B_p^{-1}) e^{-ip \cdot x} \quad (10)$$

Notting that,

$$\langle \lambda_{p,s} | \hat{\phi}_j^\dagger(y) | \Omega \rangle = \overline{\langle \Omega | \hat{\phi}_j(y) | \lambda_{p,s} \rangle} = C_\lambda^* S_{js}^*(B_p^{-1}) e^{ip \cdot y} \quad (11)$$

Therefore, defining $Z_\lambda = |C_\lambda|^2 \geq 0$ we have,

$$\langle \Omega | \hat{\phi}_i(x) \hat{\phi}_j^\dagger(y) | \Omega \rangle = \sum_{\lambda} \int_{p^2=m_\lambda^2} \frac{d^3p}{(2\pi)^3} \frac{e^{-ip \cdot (x-y)}}{2p_0} Z_\lambda \sum_s S_{is}(B_p^{-1}) S_{js}^*(B_p^{-1}) \quad (12a)$$

$$= \sum_{\lambda} \int_{p^2=m_\lambda^2} \frac{d^3p}{(2\pi)^3} \frac{e^{-ip \cdot (x-y)}}{2p_0} Z_\lambda (S(B_p^{-1}) S^\dagger(B_p^{-1}))_{ij} \quad (12b)$$

A similar computation shows that,

$$\langle \Omega | \hat{\phi}_j^\dagger(y) \hat{\phi}_i(x) | \Omega \rangle = \sum_{\bar{\lambda}, s} \int_{p^2=m_{\bar{\lambda}}^2} \frac{d^3p}{(2\pi)^3} \frac{1}{2p_0} \langle \Omega | \hat{\phi}_j^\dagger(y) | \bar{\lambda}_{p,s} \rangle \langle \bar{\lambda}_{p,s} | \hat{\phi}_i(x) | \Omega \rangle \quad (13a)$$

$$= \sum_{\bar{\lambda}, s} \int_{p^2=m_{\bar{\lambda}}^2} \frac{d^3p}{(2\pi)^3} \frac{e^{-ip \cdot (y-x)}}{2p_0} \langle \Omega | S_{jk}^*(B_p^{-1}) \hat{\phi}_k^\dagger(0) | \bar{\lambda}_s \rangle \langle \bar{\lambda}_s | S_{i\ell}(B_p^{-1}) \hat{\phi}_\ell(0) | \Omega \rangle \quad (13b)$$

$$= \sum_{\bar{\lambda}} \int_{p^2=m_{\bar{\lambda}}^2} \frac{d^3p}{(2\pi)^3} \frac{e^{-ip \cdot (y-x)}}{2p_0} Z_{\bar{\lambda}} (S(B_p^{-1}) S^\dagger(B_p^{-1}))_{ij} \quad (13c)$$

8.3 The Representation Function

Let H_{m_λ} denote the hyperbola $\{p^2 = m_\lambda^4\} \subset \mathbb{R}^{1,3}$. Then we define a matrix of functions $f_{ij} : H_{m_\lambda} \rightarrow \mathbb{C}$ via,

$$f_{ij}(p) = (S(B_p) S^\dagger(B_p))_{ij} \quad (14)$$

I claim that $W = \text{Span}(f_{ij}) \subset \mathcal{C}^\infty(H_{m_\lambda})$ is a finite-dimensional Spin $(1, 3)$ -representation. Consider,

$$f(\Lambda p) = S(B_{\Lambda p}) S^\dagger(B_{\Lambda p}) \quad (15)$$

and the Lorentz transformation $T = B_p^{-1} \Lambda^{-1} B_{\Lambda p}$ which acts on the time Killing vector as follows,

$$B_p^{-1} \Lambda^{-1} B_{\Lambda p} m_\lambda e_t = B_p^{-1} \Lambda^{-1} (\Lambda p) = B_p^{-1} p = m_\lambda e_t \quad (16)$$

so T preserves the time direction. Therefore, T must be a rotation (see Lemma ??) so $T = R_\Lambda$. Therefore,

$$B_{\Lambda p} = \Lambda B_p R_\Lambda \quad (17)$$

This implies that,

$$f(\Lambda p) = S(\Lambda) S(B_p) S(R_\Lambda) S^\dagger(R_\Lambda) S^\dagger(B_p) S^\dagger(\Lambda) \quad (18a)$$

$$= S(\Lambda) S(B_p) S^\dagger(B_p) S^\dagger(\Lambda) = S(\Lambda) \cdot f(p) \cdot S^\dagger(\Lambda) \quad (18b)$$

Where I have used the fact that $\text{SO}(3) \subset \text{SO}(1, 3)$ is compact and thus we may choose our representation restricted to this compact subgroup $S|_{\text{SO}(3)}$ to be unitary. We have therefore, shown that V transforms in the $W \otimes \bar{W}$ representation which is finite and decomposes into irreducible $\text{SO}(1, 3)$ -reps as,

$$W \otimes \bar{W} = (\mu, \nu) \otimes (\nu, \mu) = \sum_{|\mu-\nu| \leq j_+, j_+ \leq \mu+\nu} (j_+, j_-) \quad (19)$$

However, V is a representation consisting of Lorentz covariant functions and therefore by general theory (CITE THIS) V decomposes as a sum of irreducible harmonic

functions on the hyperbola in representations (J, J) for $J \in \{0, \frac{1}{2}, 1, \frac{3}{2}, \dots\}$ so we in fact have only the following direct summands in the decomposition of V ,

$$V \hookrightarrow \bigoplus_{\mu-\nu \leq J \leq \mu+\nu} (J, J) \quad (20)$$

This implies the following harmonic decomposition,

$$f_{ij}(p^\mu) = \sum_{J=|\mu-\nu|}^{\mu+\nu} f_J \sum_{-J \leq m_+, m_- \leq J} \begin{pmatrix} i & j \\ J & m_+ & m_- \end{pmatrix} m_\lambda^{2J} \mathcal{Y}_{m_+, m_-}^J(p^\mu/m_\lambda) \quad (21)$$

where these symbols are the generalizations of the Clebsch-Gordon decomposition coefficients. The harmonics $\mathcal{Y}_{m_+, m_-}^J(x)$ are homogeneous polynomials of degree $2J$ (CITE THIS FACT). Therefore, we may analytically continue p off-shell i.e. over $H_{m_\lambda} \subset \mathbb{R}^{1,3} \subset \mathbb{C}^4$ to both arbitrary real and complex momenta by the definition,

$$f_{ij}(p) = \sum_{J=|\mu-\nu|}^{\mu+\nu} f_J \sum_{-J \leq m_+, m_- \leq J} \begin{pmatrix} i & j \\ J & m_+ & m_- \end{pmatrix} \mathcal{Y}_{m_+, m_-}^J(p) \quad (22)$$

Now, since \mathcal{Y}_{m_+, m_-}^J has degree $2J$ we have $\mathcal{Y}_{m_+, m_-}^J(-p) = (-1)^{2J} \mathcal{Y}_{m_+, m_-}^J(p)$ and furthermore, since $|\mu - \nu| \leq J \leq \mu + \nu$ indexes by one, each possible $2J$ has the same parity as $2(\mu + \nu) = 2s$ where s is the spin of the representation W . This follows from the fact that,

$$\hat{J}_+ + \hat{J}_- = \hat{J} \quad (23)$$

so when W restricts to $\text{SO}(3) \hookrightarrow \text{SO}(1, 3)$ it decomposes into irreps with spin $|\mu - \nu| \leq s \leq \mu + \nu$. Therefore, we again have that $(-1)^{2J} = (-1)^{2s} = (-1)^{2(\mu+\nu)}$ where $2s$ gives the parity for the possible spin states of the particle. Applying this discussion,

$$f_{ij}(-p) = \sum_{J=|\mu-\nu|}^{\mu+\nu} f_J \sum_{-J \leq m_+, m_- \leq J} \begin{pmatrix} i & j \\ J & m_+ & m_- \end{pmatrix} \mathcal{Y}_{m_+, m_-}^J(-p) \quad (24a)$$

$$= \sum_{J=|\mu-\nu|}^{\mu+\nu} f_J \sum_{-J \leq m_+, m_- \leq J} \begin{pmatrix} i & j \\ J & m_+ & m_- \end{pmatrix} (-1)^{2J} \mathcal{Y}_{m_+, m_-}^J(p) \quad (24b)$$

$$= (-1)^{2s} f_{ij}(p) \quad (24c)$$

8.4 Completing the Proof

The derived relations hold for $f_{ij}(p)$ off-shell so we need to modify the two-point correlation functions to take p^μ off-shell. Using the polynomial nature of the functions $f_{ij}(p)$,

$$\langle \Omega | \hat{\phi}_i(x) \hat{\phi}_j^\dagger(y) | \Omega \rangle = \sum_\lambda Z_\lambda f_{ij}(+i\partial_x) D_\lambda(x - y) \quad (25a)$$

$$\langle \Omega | \hat{\phi}_j^\dagger(y) \hat{\phi}_i(x) | \Omega \rangle = \sum_{\bar{\lambda}} Z_{\bar{\lambda}} f_{ij}(-i\partial_x) D_{\bar{\lambda}}(y - x) \quad (25b)$$

where,

$$D_\lambda(x-y) = \int_{p^2=m_\lambda^2} \frac{d^3p}{(2\pi)^3} \frac{e^{-ip \cdot (x-y)}}{2p_0} \quad (26)$$

When x and y are spacelike separated i.e. $(x-y)^2 < 0$ then $D(x-y) = D(y-x)$ (CITE PESKIN 28). In this case, applying our result of the previous section,

$$\langle \Omega | \hat{\phi}_j^\dagger(y) \hat{\phi}_i(x) | \Omega \rangle = \sum_\lambda Z_\lambda f_{ij}(-i\partial_x) D_\lambda(x-y) \quad (27a)$$

$$= (-1)^{2s} \sum_{\bar{\lambda}} Z_{\bar{\lambda}} f_{ij}(+i\partial_x) D_{\bar{\lambda}}(x-y) \quad (27b)$$

From relativistic causality arguments we know that these field operators must either commute or anti-commute. Since $Z_\lambda \geq 0$ the value of $(-1)^{2s}$ fixes the sign of this relation proving our theorem. In fact, we see slightly more. In order that these correlation functions actually commute or anti-commute, there must be a spectral symmetry $\lambda \leftrightarrow \bar{\lambda}$ reversing charge but preserving mass and spin. Thus, the spin-statistics relationship also establishes the existence and spectral symmetry of anti-particles.

9 Lemmas

Lemma 9.1. If a Lorentz transformation $\Lambda \in \text{SO}(1,3)$ preserves the temporal Killing vector e_t then Λ is a rotation.

Proof. We can write any four vector in the form $x^\mu = x^0 e_t + \vec{x}$. Then, $\Lambda x^\mu = x^0 e_t + \Lambda \vec{x}$. Furthermore $e_t \cdot \vec{x} = 0$ and Λ preserves the Minkowski inner product so $e_t \cdot \Lambda \vec{x}$ and thus \vec{x} is also a vector. Furthermore, Λ preserves $(x^0)^2 - \vec{x}^2$ so $\Lambda \vec{x}$ must be a vector with $|\Lambda \vec{x}|^2 = |\vec{x}|^2$ so Λ must be a rotation since $\det \Lambda = 1$. \square