

REU Notes

Benjamin Church

May 30, 2018

Contents

1 Introduction

2

1 Introduction

Suppose $f \in \mathbb{Z}[X_0, \dots, X_n]$ is an integral polynomial in n -variables. We want to find relations between the complex solutions to $f = 0$ and the solutions of f over $\mathbb{Z}, \mathbb{Q}, \mathbb{F}_{p^k}$. We want to find out how many solutions to the system,

$$\begin{aligned} f_1(x_0, \dots, x_n) &= 0 \\ &\vdots \\ f_r(x_0, \dots, x_n) &= 0 \end{aligned}$$

over $\mathbb{F}_q[x_0, \dots, x_n]$. How many solutions does this system have over \mathbb{F}_{q^k} .

Definition: $X(\mathbb{F}_{q^k})$ is the set of solutions to $\forall i : f_i = 0$ in $(\mathbb{F}_{q^k})^{n+1}$.

Weil defined the zeta function,

Definition:

$$\zeta_X(t) = \exp \left[\sum_{r=1}^{\infty} \frac{\#|X(\mathbb{F}_{q^r})|}{r} t^r \right]$$

and made the following conjectures which were later proven,

Theorem 1.1. *The following hold about ζ_X for a variety X ,*

1. $\zeta_X(t)$ is a rational function.
2. If f_i are homogenous polynomials define the projective solutions,

$$X'(\mathbb{F}_{q^k}) = \frac{X(\mathbb{F}_{q^k}) \setminus \{\vec{0}\}}{\mathbb{F}_{q^k}^\times}$$

and $\zeta_{X'}(t)$ satisfied a function equation if X' is smooth.

3. All roots and poles ζ_X in \mathbb{C} are q -Weil numbers e.g. if α is a root or pole then $|\alpha| = q^{i/2}$ for some $i \in \mathbb{Z}$.
4. If X is smooth and projective then

$$\zeta_X(t) = \prod_{i=0}^d p_i(t)^{(-1)^{i+1}}$$

where $d = \dim X$ and $p_i \in \mathbb{Z}[t]$. For almost all primes $\deg p_i = b_i(X(\mathbb{C}))$ and the roots of p_i have absolute value $q^{i/2}$ where b_i is the i^{th} Betti number.

Definition: Suppose $f(x_0, x_n)$ is a homogeneous polynomial in $\mathbb{F}_q[x_0, \dots, x_n]$ then,

$$X(\mathbb{F}_{q^k}) = \frac{\{\vec{x} \in (\mathbb{F}_{q^k})^{n+1} \mid f(\vec{x}) = 0\} \setminus \{\vec{0}\}}{\mathbb{F}_{q^k}^\times}$$

Definition: A variety X is supersingular if ζ_X has the property that every root and pole is of the form $\omega q^{i/2}$ where ω is a root of unity.

There are many conjectures about supersingular varieties.

Definition: A 2-dimensional hypersurface is unirational if there exist $x_i(t_1, t_2) \in \overline{\mathbb{F}_q}(t_1, t_2)$ such that $f(x_0(t_1, t_2), \dots, x_3(t_1, t_2)) = 0$.

Conjecture 1.2 (Shioda). *A supersingular 2-dimensional hypersurface is unirational*

Conjecture 1.3 (Tate). *Supersingular surfaces have a lot of curves in them!*