

# Physics GR8049 Quantum Field Theory III

## Assignment # 1

Benjamin Church

February 12, 2019

## 1 Unitarity Bound

### 1.1

Consider the conformal algebra with generators  $D$ ,  $K_\mu$ ,  $P_\mu$ , and  $M_{\mu\nu}$  satisfying the standard (Euclidean) commutation relations of which I only write down the relevant ones,

$$\begin{aligned}[D, P_\mu] &= P_\mu \\ [D, K_\mu] &= -K_\mu \\ [K_\mu, P_\nu] &= 2(\delta_{\mu\nu}D - M_{\mu\nu}) \\ [M_{\mu\nu}, P_\rho] &= \delta_{\nu\rho}P_\mu - \delta_{\mu\rho}P_\nu \\ [M_{\mu\nu}, K_\rho] &= \delta_{\nu\rho}K_\mu - \delta_{\mu\rho}K_\nu\end{aligned}$$

Let  $|\Delta\rangle = \mathcal{O}_\Delta(0)|\Omega\rangle$  be a spinless primary state corresponding to the spinless primary operator  $\mathcal{O}_\Delta(x)$  with scaling dimension  $\Delta$  i.e.  $M_{\mu\nu}|\Delta\rangle = 0$  and  $K_\mu|\Delta\rangle = 0$ . This state has scaling dimension  $\Delta$  i.e.  $D|\Delta\rangle = \Delta|\Delta\rangle$ . Then consider the combination,

$$\begin{aligned}\langle\Delta|K_\mu K_\nu P_\rho P_\sigma|\Delta\rangle &= \langle\Delta|K_\mu[K_\nu, P_\rho]P_\sigma|\Delta\rangle + \langle\Delta|K_\mu P_\rho K_\nu P_\sigma|\Delta\rangle \\ &= \langle\Delta|K_\mu 2(\delta_{\nu\rho}D - M_{\nu\rho})P_\sigma|\Delta\rangle + \langle\Delta|K_\mu P_\rho[K_\nu, P_\sigma]|\Delta\rangle + \langle\Delta|K_\mu P_\rho P_\sigma K_\nu|\Delta\rangle\end{aligned}$$

Where I have used the fact that  $K_\nu|\Delta\rangle = 0$  since  $|\Delta\rangle$  is primary. Then we have,

$$\begin{aligned}\langle\Delta|K_\mu K_\nu P_\rho P_\sigma|\Delta\rangle &= \langle\Delta|K_\mu 2(\delta_{\nu\rho}D - M_{\nu\rho})P_\sigma|\Delta\rangle + \langle\Delta|K_\mu P_\rho 2(\delta_{\mu\sigma}D - M_{\mu\sigma})|\Delta\rangle \\ &= \langle\Delta|K_\mu 2[\delta_{\nu\rho}D - M_{\nu\rho}, P_\sigma]|\Delta\rangle + \langle\Delta|K_\mu P_\sigma 2(\delta_{\nu\rho}D - M_{\nu\rho})|\Delta\rangle + \langle\Delta|K_\mu P_\rho 2\delta_{\mu\sigma}\Delta|\Delta\rangle \\ &= 2\langle\Delta|K_\mu(\delta_{\nu\rho}P_\sigma - (\delta_{\rho\sigma}P_\nu - \delta_{\nu\sigma}P_\rho))|\Delta\rangle + 2(\delta_{\mu\sigma} + \delta_{\nu\sigma})\Delta\langle\Delta|K_\mu P_\rho|\Delta\rangle\end{aligned}$$

Therefore, taking traces,

$$\begin{aligned}\langle\Delta|K_\mu K^\mu P_\rho P^\rho|\Delta\rangle &= 2\langle\Delta|K_\mu(\delta^{\mu\rho}P_\rho - (d\delta^{\mu\nu}P_\nu - \delta^{\mu\rho}P_\rho))|\Delta\rangle + 4\delta^{\mu\rho}\Delta\langle\Delta|K_\mu P_\rho|\Delta\rangle \\ &= 2(2 - d + 2\Delta)\langle\Delta|K^\mu P_\mu|\Delta\rangle\end{aligned}$$

However, in radial quantization we have the relation  $P_\mu^\dagger = K_\mu$  and thus,

$$|P_\mu P^\mu|\Delta\rangle|^2 = \langle\Delta|P_\mu^\dagger(P^\mu)^\dagger P_\rho P^\rho \text{ker } \Delta = \langle\Delta|K_\mu K^\mu P_\rho P^\rho \text{ker } \Delta \geq 0$$

Furthermore,

$$\langle\Delta|K^\mu P_\mu|\Delta\rangle = \delta^{\mu\rho}\langle\Delta|P_\rho^\dagger P_\mu|\Delta\rangle = \sum_\mu |P_\mu|\Delta\rangle|^2 \geq 0$$

Since both sides of the derived expression must be positive and  $\langle \Delta | K_\mu P_\rho | \Delta \rangle \geq 0$  we find that,

$$2 - d + 2\Delta \geq 0 \implies \Delta \geq \frac{d-2}{2}$$

giving the unitarity bound in arbitrary dimensions.

## 1.2

Using the scalar equations of motion, we find that, for a scalar of mass  $m$ , the primary scaling dimension must satisfy,

$$\Delta(\Delta - d) = m^2$$

which has solutions,

$$\Delta_{\pm} = \frac{d}{2} \pm \sqrt{m^2 + \left(\frac{d}{2}\right)^2}$$

Since the modes have asymptotic behavior  $r^{-\Delta}$  basic conditions on normalizability imply that  $\Delta > 0$ . The range in which normalization issue do not arise for either choice of  $\Delta$  occurs when both  $\Delta_+$  and  $\Delta_-$  are positive. Therefore, we must have,

$$\frac{d}{2} > \sqrt{m^2 + \left(\frac{d}{2}\right)^2} \implies m^2 < 0$$

However, the scaling dimension  $\Delta$  cannot become complex since this will violate the Hermiticity of  $D$  whose eigenvalues are  $\Delta$ . Therefore, we are limited by the minimum value of the quadratic  $\Delta(\Delta - d)$  over the reals which occurs at  $\Delta = \frac{d}{2}$  and gives,

$$m^2 \geq -\left(\frac{d}{2}\right)^2$$

Therefore, the consistent range of masses for two choices of boundary conditions is,

$$-\left(\frac{d}{2}\right)^2 \leq m^2 \leq 0$$

## 1.3

Suppose we choose to quantize both modes with “standard” and “alternate” boundary conditions simultaneously. Then we have a mode expansion containing both,

$$\phi_+(t, x) = \frac{e^{-i\Delta_+ t}}{(1 + r^2)^{\Delta_+/2}} \quad \text{and} \quad \phi_-(t, x) = \frac{e^{-i\Delta_- t}}{(1 + r^2)^{\Delta_-/2}}$$

However, we quantize with respect to the Klien-Gordon inner product,

$$(\phi_1, \phi_2) = -i \int \sqrt{-g} g^{tt} d^3x (\phi_1^* \partial_t \phi_2 - \phi_2 \partial_t \phi_1^*)$$

However, applying this inner product to the plus and minus modes we find,

$$\begin{aligned}
(\phi_+, \phi_-) &= -i \int \sqrt{-g} g^{tt} d^3x (\phi_+^* \partial_t \phi_- - \phi_- \partial_t \phi_+^*) \\
&= - \int \sqrt{-g} g^{tt} d^3x (1+r^2)^{-\frac{\Delta_+ + \Delta_-}{2}} (e^{i(\Delta_+ - \Delta_-)t} \Delta_- + e^{i(\Delta_+ - \Delta_-)t} \Delta_+) \\
&= -(\Delta_- + \Delta_+) e^{i(\Delta_+ - \Delta_-)t} \int \sqrt{-g} g^{tt} d^3x (1+r^2)^{-\frac{\Delta_+ + \Delta_-}{2}}
\end{aligned}$$

which is both nonzero and time-dependent. Since  $\phi_+$  and  $\phi_-$  are supposed to be energy modes with different energy eigenvalues, namely  $\Delta_+$  and  $\Delta_-$  respectively, if  $\hat{H}$  is Hermitian then they must be orthogonal. However, since they are not in fact orthogonal, this quantization prescription violates the Hermiticity of  $\hat{H}$  and thus the unitarity of the time evolution operator  $e^{-i\hat{H}t}$ . More directly, unitary time evolution implies that inner products are invariant under time translation and therefore cannot be time-dependent as we observed the inner product  $(\phi_+, \phi_-)$  to be.

## 2 Boundary Correlators of EAdS and Conformal Symmetry

### 2.1

The boundary two-point function has the expression,

$$\langle \mathcal{O}(\vec{u}) \mathcal{O}(\vec{u}') \rangle = \frac{c_\Delta}{|\vec{u} - \vec{u}'|^{2\Delta}}$$

where the constant  $c_\Delta$  is determined by matching the propagator in Minkowski space in the short distance limit to be,

$$c_\Delta = \frac{\Gamma(\Delta)}{2\pi^{d/2} \Gamma(\Delta - 1 - \frac{d}{2})}$$

We have show that there is a unitarity bound,

$$\Delta \geq \frac{d-2}{2}$$

When  $\Delta$  crosses this threshold, the argument of the denominator Gamma function becomes negative. Furthermore,  $\Gamma(x)$  as  $x$  crosses zero diverges to  $+\infty$  and then comes back from  $-\infty$  as  $x$  becomes negative. Thus  $\Gamma(x)$  crosses from being positive to negative as its argument does. This is more clearly summarized by taylor expanding,

$$\frac{1}{\Gamma(x)} = x + O(x^2)$$

Thus, we have, about  $\epsilon = 0$  where  $\epsilon = \Delta - \frac{d-2}{2}$  we have,

$$c_\Delta = \frac{\Gamma(\frac{d-2}{2})}{2\pi^{d/2}} \epsilon + O(\epsilon^2)$$

Thus as we lower  $\Delta$  past the unitarity bound  $\frac{d-2}{2}$  (i.e. take  $\epsilon$  negative) the coefficient  $c_\Delta$  changes sign (passing through zero) from positive to negative and, since all other terms remain well-defined in this transition, the two-point function must also become negative. Intreguingly, exactly at the unitarity bound  $\Delta = \frac{d-2}{2}$  the constant  $c_\Delta$  vanishes and thus,

$$\langle \mathcal{O}(\vec{u}) \mathcal{O}(\vec{u}') \rangle = 0$$

## 2.2

First, consider EAdS in Euclidean global coordinates,

$$X^0 = \sqrt{1+r^2} \cosh t \quad X^d = \sqrt{1+r^2} \sinh t \quad X^i = x^i$$

in which the metric becomes,

$$ds^2 = (1+r^2)dt^2 + \frac{dr^2}{1+r^2} + r^2 d\Omega^2$$

In Euclidean EAdS the length inner product becomes,

$$P = X^I Y_I = -X^0 Y^0 + X^I Y^I = -\sqrt{(1+r^2)(1+r'^2)} \cosh(t-t') + rr' \cos \eta$$

where  $\eta$  is the angle between  $x^i$  and  $y^i$  on the boundary sphere. We have shown that the two-point function in the large-distance limit becomes,

$$G_\Delta \rightarrow c_\Delta (-2p)^{-\Delta}$$

Therefore, setting  $r = r'$  and taking the limit  $r \rightarrow \infty$  gives,

$$P = r^2 (-\cosh(t-t') + \cos \eta)$$

Thus, define the boundary operator via,

$$\mathcal{O}(t, \Omega) = \lim_{r \rightarrow \infty} r^\Delta \phi(r, t, \Omega)$$

such that the boundary two-point function becomes,

$$\begin{aligned} \langle \mathcal{O}(t, \Omega) \mathcal{O}(t', \Omega') \rangle &= \lim_{r \rightarrow \infty} r^{2\Delta} \langle \phi(r, t, \Omega) \phi(r, t', \Omega') \rangle = \lim_{r \rightarrow \infty} \frac{r^{2\Delta} c_\Delta}{2^\Delta r^{2\Delta} (\cosh(t-t') - \cos \eta)^\Delta} \\ &= \frac{c_\Delta}{2^\Delta (\cosh(t-t') - \cos \eta)^\Delta} \end{aligned}$$

Second, consider EAdS in Euclidean hyperbolic ball coordinates,

$$X^I = \frac{2y^I}{1-|y|^2} \quad X^0 = \frac{1+|y|^2}{1-|y|^2}$$

where I restrict,

$$|y|^2 = \sum_{I=1}^{d+1} (y^I)^2 < 1$$

In these coordinates, the ambient coordinate dot product becomes,

$$P = X^I X_I = -\frac{1+|y|^2}{1-|y|^2} \frac{1+|y'|^2}{1-|y'|^2} + \frac{4y^I y'_I}{(1-|y|^2)(1-|y'|^2)}$$

In the case that  $|y| = |y'|$  (preparing to take the limits of both to the boundary). Then we have,

$$P = -\left(\frac{1+|y|^2}{1-|y|^2}\right)^2 + \frac{4y^I y'_I}{(1-|y'|^2)} = -\frac{1+2|y|^2-4y^I y'_I+|y|^4}{(1-|y|^2)^2} = -1 + \frac{4(y^I y'_I - |y|^2)}{(1-|y|^2)^2}$$

Thus, in the limit  $|y|^2 \rightarrow 1$  we find,

$$P = \frac{4(y^I y'_I - |y|^2)}{(1 - |y|^2)^2}$$

Thus, define the boundary operator via,

$$\mathcal{O}(y^I) = \lim_{|y| \rightarrow 1} (1 - |y|^2)^{-\Delta} \phi(y^I)$$

such that the boundary two-point function becomes,

$$\begin{aligned} \langle \mathcal{O}(y^I) \mathcal{O}(y'^I) \rangle &= \lim_{|y| \rightarrow 1} (1 - |y|^2)^{-2\Delta} \langle \phi(y^I) \phi(y'^I) \rangle = \lim_{|y| \rightarrow 1} (1 - |y|^2)^{-2\Delta} \frac{(1 - |y|^2)^{2\Delta} c_\Delta}{8^\Delta (1 - y^I y'^I)^\Delta} \\ &= \frac{c_\Delta}{8^\Delta (1 - y^I y'^I)^\Delta} \end{aligned}$$

### 2.3

In Euclidean Poincare coordinates, consider a change of coordinates  $z = \lambda \tilde{z}$  and  $\vec{u} = \lambda \vec{\tilde{u}}$  with  $\lambda > 0$  some constant. Then the bulk metric becomes,

$$ds^2 = \frac{d\vec{u}^2 + dz^2}{z^2} = \frac{d\vec{\tilde{u}}^2 + d\tilde{z}^2}{\tilde{z}^2}$$

and thus this scaling is an isometry. In these new coordinates, the boundary metric and operators are defined via,

$$\begin{aligned} d\tilde{\sigma}^2 &= \lim_{\tilde{z} \rightarrow 0} \tilde{z}^2 ds^2|_{\tilde{z}} = d\vec{\tilde{u}}^2 \\ \tilde{\mathcal{O}}(\vec{\tilde{u}}) &= \lim_{\tilde{z} \rightarrow 0} \tilde{z}^{-\Delta} \tilde{\phi}(\tilde{z}, \vec{\tilde{u}}) = \lim_{\tilde{z} \rightarrow 0} (z/\lambda)^{-\Delta} \phi(z, u) = \lambda^\Delta \mathcal{O}(\vec{u}) = \lambda^\Delta \mathcal{O}(\lambda \vec{\tilde{u}}) \end{aligned}$$

Now consider the two-point function expressed in terms of these new boundary operators in the new coordinates.

$$\langle \tilde{\mathcal{O}}(\vec{\tilde{u}}) \tilde{\mathcal{O}}(\vec{\tilde{u}}') \rangle = \langle \lambda^\Delta \mathcal{O}(\lambda \vec{\tilde{u}}) \lambda^\Delta \mathcal{O}(\lambda \vec{\tilde{u}}') \rangle = \lambda^{2\Delta} \langle \mathcal{O}(\lambda \vec{\tilde{u}}) \mathcal{O}(\lambda \vec{\tilde{u}}') \rangle = \frac{\lambda^{2\Delta} c_\Delta}{|\lambda^2 \vec{\tilde{v}} - \lambda^2 \vec{\tilde{v}}'|^2} = \frac{c_\Delta}{|\vec{\tilde{u}} - \vec{\tilde{u}}'|^2}$$

Therefore, the two-point function of the new boundary operators in the new boundary coordinates takes exactly the same form as the old operators in the old coordinates.

### 2.4

In Poincare coordinates, consider a coordinate transformation  $z = Z(\tilde{z}, \tilde{u})$  and  $u^i = U^i(\tilde{z}, \tilde{u})$  such that we preserve the metric,

$$ds^2 = \frac{(du^2 + dz^2)}{z^2} = \frac{(d\tilde{u}^2 + d\tilde{z}^2)}{\tilde{z}^2}$$

We care about the metric near the boundary so we Taylor expand about  $\tilde{z}0$ . Since such isometries must preserve the boundary, they map  $z = 0$  to  $\tilde{z} = 0$ . Therefore we may expand,

$$z = (\partial_{\tilde{z}} Z(\tilde{z}, \tilde{u})|_{\tilde{z}=0}) \tilde{z} + O(\tilde{z}^2)$$

Let  $\gamma(\tilde{u}) = \partial_{\tilde{z}} Z(\tilde{z}, \tilde{u})|_{\tilde{z}=0}$ . Then plugging in with  $d\tilde{z} = 0$  we find,

$$ds^2 = \frac{du^2 + d\gamma(\tilde{u})^2 \tilde{z}^2 + O(\tilde{z})}{\gamma(\tilde{u})^2 \tilde{z}^2 + O(\tilde{z}^3)} = \frac{du^2 + d\gamma(\tilde{u})^2 \tilde{z}^2}{\gamma(\tilde{u})^2 \tilde{z}^2} + O(\tilde{z}^{-1}) = \frac{du^2}{\gamma(\tilde{u})^2 \tilde{z}^2} + O(1)$$

Furthermore, using the fact that this transformation preserves the metric we find (recalling that we set  $d\tilde{z} = 0$ ),

$$ds^2 = \frac{d\tilde{u}^2}{\tilde{z}^2} = \frac{du^2}{\gamma(\tilde{u})^2 \tilde{z}^2} + O(1)$$

and thus,

$$\gamma(\tilde{u})^2 d\tilde{u}^2 = du^2 + O(\tilde{z}^2)$$

and therefore, in the limit  $z \rightarrow 0$  we find,

$$\gamma(\tilde{u})^2 d\tilde{u}^2 = du^2$$

or equivalently,

$$d\tilde{\sigma}^2 = \lim_{\tilde{z} \rightarrow 0} \tilde{z}^2 ds^2 = \lim_{\tilde{z} \rightarrow 0} d\tilde{u}^2 = \lim_{\tilde{z} \rightarrow 0} \left[ \frac{du^2}{\gamma(\tilde{u})^2} + O(\tilde{z}^2) \right] = \frac{du^2}{\gamma(\tilde{u})^2} = \frac{d\sigma^2}{\gamma(\tilde{u}^2)}$$

Therefore the conformal factor is

$$\rho(\tilde{u}) = \gamma(\tilde{u}) = (\partial_{\tilde{z}} Z(\tilde{z}, \tilde{u})|_{\tilde{z}=0})$$

Furthermore, the boundary operators can be written as,

$$\begin{aligned} \mathcal{O}(u) &= \lim_{z \rightarrow 0} z^{-\Delta} \phi(z, u) = \lim_{z \rightarrow 0} (z/\tilde{z})^{-\Delta} \tilde{z}^{-\Delta} \tilde{\phi}(\tilde{z}, \tilde{u}) = \lim_{z \rightarrow 0} (Z(\tilde{z}, \tilde{u})/\tilde{z})^{-\Delta} \lim_{z \rightarrow 0} \tilde{z}^{-\Delta} \tilde{\phi}(\tilde{z}, \tilde{u}) \\ &= (\partial_{\tilde{z}} Z(\tilde{z}, \tilde{u})|_{\tilde{z}=0})^{-\Delta} \tilde{\mathcal{O}}(\tilde{u}) = \gamma(\tilde{u})^{-\Delta} \tilde{\mathcal{O}}(\tilde{u}) \end{aligned}$$

Therefore,

$$\tilde{\mathcal{O}}(\tilde{u}) = \gamma(\tilde{u})^\Delta \mathcal{O}(U(\tilde{z}, \tilde{u}))$$

## 2.5

Consider the coordinate transformation from Euclidean Poincare coordinates to Euclidean global coordinates given by,

$$t = \log \sqrt{\vec{u}^2 + z^2} \xrightarrow{z \rightarrow 0} \log |u| \quad x^i = \frac{u^i}{z}$$

Now consider the tranformation between boundary operators,

$$\mathcal{O}_{\text{cyl}}(t, \Omega) = \lim_{r \rightarrow \infty} r^\Delta \phi(r, t, \Omega) = \lim_{z \rightarrow 0} \frac{|u|^\Delta}{z^\Delta} \phi(z, u) = |u|^\Delta \lim_{z \rightarrow 0} z^{-\Delta} \phi(z, u) = |u|^\Delta \mathcal{O}_{\text{pl}}(u)$$

Using the inverse transformation,

$$z = \frac{e^t}{\sqrt{1+r^2}} \quad \vec{u} = \frac{\vec{x}e^t}{\sqrt{1+r^2}} \xrightarrow{r \rightarrow \infty} \vec{\Omega}e^t$$

we can also go the opposite direction to find,

$$\mathcal{O}_{\text{pl}}(u) = \lim_{z \rightarrow 0} z^{-\Delta} \phi(z, u) = \lim_{r \rightarrow \infty} e^{-t\Delta} (1+r^2)^{\Delta/2} \phi(r, t, \Omega) = e^{-t\Delta} \lim_{r \rightarrow \infty} r^\Delta \phi(r, t, \Omega) = e^{-t\Delta} \mathcal{O}_{\text{cyl}}(t, \Omega)$$

Furthermore, these are compatible because  $e^t = |u|$ . Now consider the conformal two-point function,

$$\begin{aligned}\langle \mathcal{O}_{\text{cyl}}(t, \Omega) \mathcal{O}_{\text{cyl}}(t', \Omega') \rangle &= (|u||u'|)^\Delta \langle \mathcal{O}_{\text{pl}}(u) \mathcal{O}_{\text{pl}}(u') \rangle = \frac{c_\Delta (|u||u'|)^\Delta}{|\vec{u} - \vec{u}'|^{2\Delta}} \\ &= \frac{c_\Delta e^{(t+t')\Delta}}{|\vec{\Omega} e^t - \vec{\Omega}' e^{t'}|^{2\Delta}} = \frac{c_\Delta}{\left| \vec{\Omega} e^{\frac{1}{2}(t-t')\Delta} - \vec{\Omega}' e^{\frac{1}{2}(t'-t)\Delta} \right|^{2\Delta}}\end{aligned}$$

However,

$$\begin{aligned}\left| \vec{\Omega} e^{\frac{1}{2}(t-t')\Delta} - \vec{\Omega}' e^{\frac{1}{2}(t'-t)\Delta} \right|^2 &= |\vec{\Omega}|^2 e^{(t-t')\Delta} + |\vec{\Omega}'|^2 e^{(t'-t)\Delta} - 2\vec{\Omega} \cdot \vec{\Omega}' \\ &= (e^{(t-t')\Delta} + e^{(t'-t)\Delta}) - 2\cos\eta = 2(\cosh(t-t') - \cos\eta)\end{aligned}$$

Therefore,

$$\langle \mathcal{O}_{\text{cyl}}(t, \Omega) \mathcal{O}_{\text{cyl}}(t', \Omega') \rangle = \frac{c_\Delta}{2^\Delta (\cosh(t-t') - \cos\eta)^\Delta}$$

which is exactly the two-point function we computed earlier in Euclidean global coordinates. Likewise, we may check this result using the forward transformation,

$$\begin{aligned}\langle \mathcal{O}_{\text{pl}}(u) \mathcal{O}_{\text{pl}}(u') \rangle &= (|u||u'|)^{-\Delta} \langle \mathcal{O}_{\text{cyl}}(t, \Omega) \mathcal{O}_{\text{cyl}}(t', \Omega') \rangle = \frac{c_\Delta (|u||u'|)^{-\Delta}}{2^\Delta (\cosh(t-t') - \cos\eta)^\Delta} \\ &= \frac{c_\Delta (|u||u'|)^{-\Delta}}{2^\Delta (\cosh(\log|u| - \log|u'|) - (|u||u'|)^{-1} \vec{u} \cdot \vec{u}')^\Delta} \\ &= \frac{c_\Delta (|u||u'|)^{-\Delta}}{\left( \frac{|u|}{|u'|} + \frac{|u'|}{|u|} - 2(|u||u'|)^{-1} \vec{u} \cdot \vec{u}' \right)^\Delta} = \frac{c_\Delta}{(|u|^2 + |u'|^2 - 2\vec{u} \cdot \vec{u}')^\Delta} \\ &= \frac{c_\Delta}{|\vec{u} - \vec{u}'|^{2\Delta}}\end{aligned}$$

which is exactly the conformal two-point function in Poincare coordinates.

## 2.6

Consider the Eulclidean generators  $M_{ij}$ ,  $P_i$ ,  $K_i$  and  $D$  which are given in Poincare coordinates as,

$$\begin{aligned}M_{ij} &= -i(u^i \partial_{u^j} - u^j \partial_{u^i}) \\ D &= -i(u^i \partial_{u^i} + z \partial_z) \\ P_i &= -i \partial_{u^i} \\ K_i &= -i(u^2 + z^2) \partial_{u^i} + 2i u^i (u^j \partial_{u^j} + z \partial_z)\end{aligned}$$

Such transformations act on the conformal boundary operators under a transformation  $\delta\phi = i\epsilon G\phi$  via,

$$\delta\mathcal{O} = \lim_{z \rightarrow 0} z^{-\Delta} \delta\phi = i\epsilon \lim_{z \rightarrow 0} z^{-\Delta} G\phi = i\epsilon \left( G \lim_{z \rightarrow 0} z^{-\Delta} \phi - \lim_{z \rightarrow 0} [G, z^{-\Delta}] \phi \right) = \mathcal{G}\mathcal{O}$$

Therefore, first we need to compute the commutators,

$$\begin{aligned}[M_{ij}, z^{-\Delta}] &= 0 \\ [D, z^{-\Delta}] &= i\Delta z^{-\Delta} \\ [P_i, z^{-\Delta}] &= 0 \\ [K_i, z^{-\Delta}] &= -2iu^i\Delta z^{-\Delta}\end{aligned}$$

Therefore, we can write,

$$[G, z^{-\Delta}] = He^{-\Delta}$$

for some function  $H$  simplifying,

$$\delta\mathcal{O} = i\epsilon \left( G \lim_{z \rightarrow 0} z^{-\Delta} \phi - \lim_{z \rightarrow 0} H z^{-\Delta} \phi \right) = i\epsilon (G\mathcal{O} - H\mathcal{O})$$

Therefore, plugging in for the given generators,

$$\begin{aligned}M_{ij} &\implies \delta\mathcal{O} = \epsilon (u^i \partial_{u^j} - u^j \partial_{u^i}) \mathcal{O}(u) \\ D &\implies \delta\mathcal{O} = \epsilon (u^i \partial_{u^i} + \Delta) \mathcal{O} \\ P_i &\implies \delta\mathcal{O} = \epsilon \partial_{u^i} \mathcal{O} \\ K_i &\implies \delta\mathcal{O} = \epsilon (u^2 \partial_{u^i} - 2u^i u^j \partial_{u^j} - 2u^i \Delta) \mathcal{O}\end{aligned}$$