

# Mathematics GU4053 Algebraic Topology

## Assignment # 6

Benjamin Church

February 17, 2020

Note. My order of path concatenation follows Hatcher,

$$\gamma * \delta(x) = \begin{cases} \gamma(2x) & x \leq \frac{1}{2} \\ \delta(2x - 1) & x \geq \frac{1}{2} \end{cases}$$

### Problem 1.

Suppose the following diagram of abelian groups commutes,

$$\begin{array}{ccccccccc} A & \xrightarrow{a} & B & \xrightarrow{b} & C & \xrightarrow{c} & D & \xrightarrow{d} & E \\ \downarrow f & & \downarrow g & & \downarrow h & & \downarrow i & & \downarrow j \\ A' & \xrightarrow{a'} & B' & \xrightarrow{b'} & C' & \xrightarrow{c'} & D' & \xrightarrow{d'} & E' \end{array}$$

with exact rows and  $f, g, i$ , and  $j$  are isomorphisms. Suppose that  $h(x) = 0$  then  $c' \circ h(x) = 0$ . By commutativity,  $i \circ c(x) = 0$  but  $i$  is an injection so  $c(x) = 0$ . Thus,  $x \in \ker c = \text{Im}(b)$  so there exists  $y \in B$  such that  $b(y) = x$  but  $h(x) = 0$  so  $h \circ b(y) = b' \circ g(y) = 0$  so  $g(y) \in \ker b' = \text{Im}(a')$  so there exists  $z \in A'$  such that  $a'(z) = g(y)$ . But  $f$  is a surjection so there exists  $q \in A$  such that  $f(q) = z$ . Then,  $g \circ a(q) = a' \circ f(q) = a'(z) = g(y)$  but  $g$  is an injection so  $a(q) = y$ . Then  $b \circ a(q) = b(y) = x$ . However, the top row is exact so  $\ker b = \text{Im}(a)$  but  $a(q) \in \text{Im}(a)$  so  $a(q) \in \ker b$  so  $b \circ a(q) = x = 0$ . Thus,  $h$  is injective.

In this proof, we never used the maps  $d, j$ , and  $d'$  so only the first four groups in the sequences are needed. Also, I only used the fact that  $f$  is a surjection,  $g$  is an injection, and  $i$  is an injection.

### Problem 2.

Let  $p : (E, e_0) \rightarrow (B, b_0)$  be a pointed fibration. The fiber of  $p$  is the subspace  $F = p^{-1}(b_0)$ . Then, define the map  $\phi : F \rightarrow N_p$  by  $\phi(x) = (x, e_{b_0})$  where  $e_{b_0}$  is the constant loop at  $b_0$ . This map is well-defined because  $x \in F = p^{-1}(b_0)$  so  $p(x) = b_0 = e_{b_0}(0)$ . The projection  $\pi_1 : N_p \rightarrow E$  is given by  $\pi_1(x, \gamma) = x$ . Therefore,  $\pi_1 \circ \phi(x) = \pi_1(x, e_{b_0}) = x$  so  $\pi_1 \circ \phi = \text{id}_F$ . However,  $\phi \circ \pi_1(x, \gamma) = \phi(x) = (x, e_{b_0})$ . Define the homotopy  $H : N_p \times I \rightarrow N_p$  by  $H(x, \gamma, t) = (x, \gamma_t)$  where  $\gamma_t(s) = \gamma(1 - (1 - r)t)$ . Thus,  $\gamma_0(r) = \gamma(1) = b_0$  and  $\gamma_1(r) = \gamma(r)$ . Therefore,  $H(x, \gamma, 0) = (x, \gamma_0) = (x, e_{b_0}) = \phi \circ \pi_1(x, \gamma)$  and  $H(x, \gamma, 1) = (x, \gamma_1) = (x, \gamma)$ . Thus,  $H$  is a homotopy between  $\phi \circ \pi_1$  and  $\text{id}_{N_p}$  so  $\phi$  is a homotopy equivalence.

### Problem 3.

Let  $f : X \rightarrow Y$  be a map of pointed spaces. Consider the projection  $\pi_1 : N_f \rightarrow X$  given by  $\pi_1(x, \gamma) = x$ . Take any space  $Z$  and maps  $g : Z \rightarrow N_f$  and  $h : Z \times I \rightarrow X$  such that the following diagram commutes,

$$\begin{array}{ccccc}
 Z & \xrightarrow{g} & N_f & \xrightarrow{\pi_2} & Y^I \\
 \pi_Z \uparrow \downarrow \iota & \nearrow \tilde{h} & \downarrow \pi_1 & & \downarrow ev_0 \\
 Z \times I & \xrightarrow{h} & X & \xrightarrow{f} & Y
 \end{array}$$

There are maps  $h : Z \times I \rightarrow X$  and  $\pi_2 \circ g \circ \pi_Z : Z \times I \rightarrow Y^I$ . Therefore, by the universal property of the pullback, there exists a unique map  $\tilde{h} : Z \times I \rightarrow N_f$  which commutes with the diagram. Therefore,  $\pi_1 \circ \tilde{h} = h$ . Furthermore,  $\tilde{h} \circ \iota : Z \rightarrow N_f$  and  $\pi_1 \circ \tilde{h} \circ \iota = h \circ \iota = p_{i_1} \circ g$ . Also,  $\pi_2 \circ \tilde{h} \circ \iota = \pi_2 \circ g \circ \pi_Z \circ \iota = \pi_2 \circ g$ . However, by the universal property of the pullback,  $g$  is the unique map  $Z \rightarrow N_f$  satisfying this property under the projections. Therefore,  $\tilde{h} \circ \iota = g$ . Thus,  $\tilde{h}$  is a lift of  $h$  at  $g$  so  $\pi_1$  is a fibration.

The map  $\pi_1 : N_f \rightarrow X$  is a fibration. Thus, take,  $\phi : F \rightarrow N_\pi$ , the natural inclusion on the fiber  $F = \pi_1^{-1}(x_0)$  which is given by  $\phi(x_0, \gamma) = (x_0, \gamma, e_{x_0})$  where  $(x_0, \gamma) \in \pi_1^{-1}(x_0)$  so  $f(x_0) = \gamma(0) = y_0$ . However,  $Y^I$  is the space of based loops (with  $I$  based at 1) so  $\gamma(1) = y_0$ . Therefore,  $\gamma$  is a loop so  $F \cong \Omega Y$  by  $(x_0, \gamma, e_{x_0}) \mapsto \gamma$ . Thus,  $\phi$  can be viewed as a map  $\phi : \Omega Y \rightarrow N_\pi$ . However, as proven in problem (2),  $\phi : F \rightarrow N_\pi$  is a homotopy equivalence. Therefore,  $\phi : \Omega Y \rightarrow N_\pi$  is a homotopy equivalence.

### Problem 4.

Consider the covering map  $p : S^n \rightarrow \mathbb{RP}^n$  given by the quotient map on antipodal points. We know from covering space theory that for  $m \geq 2$ , the map  $p_* : \pi_m(S^n) \rightarrow \pi_m(\mathbb{RP}^n)$  is an isomorphism. However, since we have some fancy new long exact sequences it seems a shame not to use them!

The covering map  $p : S^n \rightarrow \mathbb{RP}^n$  is a fibration with fiber  $S^0$ . This fibration induces the long exact sequence,

$$\begin{array}{ccccccccccc}
 \cdots & \longrightarrow & \pi_4(S^0) & \longrightarrow & \pi_4(S^n) & \longrightarrow & \pi_4(\mathbb{RP}^n) & \longrightarrow & \pi_3(S^0) & \longrightarrow & \pi_3(S^n) & \longrightarrow & \pi_3(\mathbb{RP}^n) \\
 & & & & & & & & & & & & \downarrow \\
 & & & & & & & & & & \pi_2(S^0) & \longrightarrow & \pi_2(S^n) & \longrightarrow & \pi_2(\mathbb{RP}^n) & \longrightarrow & \pi_1(S^0) & \longrightarrow & \pi_1(S^n) & \longrightarrow & \pi_1(\mathbb{RP}^n)
 \end{array}$$

However,  $\pi_m(S^0) = 0$  for any  $m > 0$  because  $S^0$  is a disjoint union of points. Therefore, for each  $m \geq 2$ , we can pick out the exact sequence,

$$0 \longrightarrow \pi_m(S^n) \xrightarrow{f} \pi_m(\mathbb{RP}^m) \longrightarrow 0$$

Because this sequence is exact,  $\ker f = \text{Im}(0) = 0$  and  $\text{Im}(f) = \ker 0 = \pi_m(\mathbb{RP}^m)$  so  $f$  is an isomorphism. Therefore,  $\pi_m(S^n) \cong \pi_m(\mathbb{RP}^n)$  for  $m \geq 2$ .

## Problem 5.

For  $m, n \in \mathbb{Z}_{>1} \cup \{\infty\}$  let  $X = \mathbb{RP}^m \times S^n$  and  $Y = \mathbb{RP}^n \times S^m$ . Using the previous problem, for  $i \geq 2$ ,

$$\pi_i(X) = \pi_i(\mathbb{RP}^m) \times \pi_i(S^n) \cong \pi_i(S^m) \times \pi_i(S^n) \cong \pi_i(S^n) \times \pi_i(S^m) \cong \pi_i(\mathbb{RP})^n \times \pi_i(S)^m \cong \pi_i(\mathbb{RP}^n \times S^m) = \pi_i(Y)$$

For  $i = 0$  this statement is trivial because both spaces are connected. For  $i = 1$  we must check the formula explicitly,

$$\pi_1(\mathbb{RP}^m \times S^n) \cong (\mathbb{Z}/2\mathbb{Z}) \times 1 \cong \mathbb{Z}/2\mathbb{Z} \quad \text{and} \quad \pi_1(\mathbb{RP}^n \times S^m) \cong (\mathbb{Z}/2\mathbb{Z}) \times 1 \cong \mathbb{Z}$$

so  $\pi_1(\mathbb{RP}^m \times S^n) \cong \pi_1(\mathbb{RP}^n \times S^m)$ . I have used the formula  $\pi_1(S^n) = 1$  for  $n > 1$  and  $\pi_1(\mathbb{RP}^n) \cong \mathbb{Z}/2\mathbb{Z}$  for  $n > 1$  because  $S^n$  is a double cover of  $\mathbb{RP}^n$  which is the universal cover.

An alternative proof of this fact using covering spaces goes as follows. Because the product of covering maps is a covering map, the product of simply connected spaces is simply connected, and the universal cover is unique up to isomorphism, we know that  $\tilde{X} = S^m \times S^n$  and  $\tilde{Y} = S^n \times S^m$  because  $S^n$  is simply connected and the universal cover of  $\mathbb{RP}^m$  is  $S^m$ . Therefore,  $\tilde{X} \cong \tilde{Y}$ . However, for  $n \geq 2$  the covering map  $p : \tilde{X} \rightarrow X$  induces an isomorphism,  $p_* : \pi_i(\tilde{X}) \rightarrow \pi_i(X)$ . Therefore,

$$\pi_i(X) \cong \pi_i(\tilde{X}) \cong \pi_i(\tilde{Y}) \cong \pi_i(Y)$$

## Problem 6.

Consider the long exact sequence of abelian groups such that every third map  $\iota_n$  is injective,

$$\cdots \longrightarrow C_{n+1} \xrightarrow{f_{n+1}} A_n \xrightarrow{\iota_n} B_n \longrightarrow C_n \xrightarrow{f_n} A_{n-1} \xrightarrow{\iota_{n-1}} B_{n-1} \longrightarrow \cdots$$

Since  $\iota_n$  is injective,  $\ker \iota_n = 0 = \text{Im}(f_{n+1})$  so  $f_{n+1}$  is the zero map. Likewise,  $\iota_{n-1}$  is injective and the sequence is exact so  $\ker \iota_{n-1} = \text{Im}(f_n) = 0$  so  $f_n$  is the zero map. Therefore, the sequence,

$$0 \longrightarrow A_n \xrightarrow{\iota_n} B_n \longrightarrow C_n \longrightarrow 0$$

is short exact.

## Problem 7.

Suppose that the sequence of abelian groups,

$$0 \longrightarrow A \xrightarrow{f} B \xrightarrow{h} C \longrightarrow 0$$

$\quad \quad \quad \curvearrowleft \quad \quad \quad$   
 $\quad \quad \quad g \quad \quad \quad$

is short exact and the map  $g : B \rightarrow A$  satisfies  $g \circ f = \text{id}_A$ . For define the homomorphism  $F : B \rightarrow A \oplus C$  by  $F(x) = (g(x), h(x))$ . Because the kernel of the last zero map is  $C$ , the map  $h$  is surjective. Also,  $g$  is a left inverse so  $g$  is surjective. Thus,  $F$  is surjective. Furthermore, suppose that  $(g(x), h(x)) = 0$  then  $h(x) = 0$  so  $x \in \ker h = \text{Im}(f)$  so there exists  $y \in B$  such that  $f(y) = x$  but  $g \circ f(y) = y$  so  $g(x) = y = 0$ . Thus,  $y = 0$  so  $f(y) = x = 0$  so  $F$  is injective. Therefore,  $F$  is an isomorphism. Thus,  $B \cong A \oplus C$ .

## Problem 8.

Let  $(X, A)$  be a pointed pair. We showed in class that the following sequence induced by the inclusion  $\iota : A \rightarrow X$ ,

$$\cdots \longrightarrow \pi_2(X, A) \longrightarrow \pi_1(A) \xrightarrow{\iota_*} \pi_1(X) \longrightarrow \pi_1(X, A) \longrightarrow \pi_0(A) \xrightarrow{\iota_*} \pi_0(X)$$

is long exact. Suppose that there exists a retraction  $r : X \rightarrow A$ . Then we know,  $r \circ \iota = \text{id}_A$ . Therefore,  $r_* \circ \iota_* = \text{id}_{\pi_n(A)}$ . Therefore,  $\iota_*$  is an injection. Applying the result of problem 6 to this long exact sequence, we have the following short exact sequence for each  $n$ ,

$$0 \longrightarrow \pi_n(A) \xrightarrow{\iota_*} \pi_n(X) \longrightarrow \pi_n(X, A) \longrightarrow 0$$

However,  $r_* : \pi_n(X) \rightarrow \pi_n(A)$  is a left inverse of  $\iota_*$  so by problem 7 this short exact sequence splits. Therefore,  $\pi_n(X) \cong \pi_n(A) \oplus \pi_n(X, A)$ .