

Work over \mathbb{C} and let $g \geq 2$. Let $\mathcal{H}_{g,n} \subset \mathcal{M}_{g,n}$ be the space of hyperelliptic curves with n marked points. The points are labeled and may or may not lie in hyperelliptic pairs or at the intersections.

$$\chi_{0,c}^{S_n}(\mathcal{H}_{g,n}) = \sum_{i \geq 0} (-1)^i W_0 H_c^i(\mathcal{H}_{g,n}, \mathbb{Q}) \in K_0(\mathbb{Q}[S_n])$$

for the mixed Hodge structure weight filtration. We view this in the Grothendieck group of S_n -representations as a virtual representation or character.

The main theorem (BCK 2023+) gives a sum-of-graphs formula,

$$z_g = \sum_{n \geq 0} \text{char}_n(\chi_{0,c}^{S_n}(\mathcal{H}_{g,n})) \in \hat{\Lambda}_{\mathbb{Q}} = \varprojlim \mathbb{Q}[[x_1, \dots, x_n]]^{S_n}$$

in the degree-completed ring of symmetric functions where,

$$\text{char}_n \left(\sum_{\lambda n} a_{\lambda} S^{\lambda} \right) = \sum_{\lambda n} a_{\lambda} s_{\lambda} \in \Lambda_n$$

where s_{λ} is the Shur function,

$$s_{\lambda} = \sum_{T \in SSYT(\lambda)} x_1^{c_1(T)} x_2^{c_2(T)} \dots$$

these are semistandard fillings of the Young tableaux. Or equivalently,

$$\text{char}_n(W) = \frac{1}{n!} \sum_{\sigma \in S_n} \text{tr } \sigma |W \cdot \psi(\sigma)$$

where,

$$\psi(\sigma) = p_{a_1} \cdots p_{a_k}$$

where σ has cycle type $a_1 \geq a_2 \geq \dots \geq a_k$ and,

$$p_i = x_1^i + x_2^i + \dots$$

Sample theorem $g = 2$ (Faber, '08),

$$z_2 = -\frac{1}{12} P_1 + \frac{1}{2} \frac{P_1}{P_2} - \frac{1}{6} \frac{P_1^2}{P_3} - \frac{1}{12} \frac{P_1}{3} P_2^2 - \frac{1}{6} \frac{P_2 P_3}{P_6}$$

where $P_i = 1 + p_i$.

Related work:

- (a) $\chi(\mathcal{H}_{g,n})$ and $\chi(M_{g,n})$ Bini '07 Bini-Harer 11, $\chi^{\text{Orb}}(\mathcal{M}_{g,n})$ Harer-Zagier '86.
- (b) $\chi^{S_n}(\mathcal{H}_{g,n})$ and $\chi^{S_n}(M_{g,n})$ Gorsky '09, '14.

0.1 Boundary Complexes

Let $U \subset X$ be a nc compactification of smooth varieties. Then there is a combinatorial object $\Delta(U \subset X)$ called the boundary complex. If $D = D_1 \cup \cdots \cup D_t$ with $D = X \setminus U$ with each D_i smooth (snc case) is a simplicial complex whose p -simplices are irreducible components of $D_{i_0} \cap \cdots \cap D_{i_p}$ for i_0, \dots, i_p .

Example 0.1.1. $U = \mathbb{G}_m^2$. Then we can consider \mathbb{P}^2 or $\mathbb{P}^1 \times \mathbb{P}^1$ are toric compactifications. We could also blow up \mathbb{P}^2 at not a torus fixed point to get a non-toric example. These give a triangle, a square, and a triangle with a tail respectively.

It turns out that the homotopy type of this complex is an invariant of U . From Deligne Hodge II,III we get,

Theorem 0.1.2. $\widetilde{H}^{k-1}(\Delta(U \subset X), \mathbb{Q}) \cong \mathbf{gr}_{(W)_0 H_c^k(U, \mathbb{Q})}.$