

Contents

1	Chapter 1	3
2	Chapter 2	3
2.1	Section 2.1	3
2.1.1	2.2.2	3
2.1.2	2.2.3	4
2.1.3	2.2.5	4
2.1.4	2.2.6 DO!!	4
2.2	Section 2.6	4
2.2.1	2.6.1	4
2.2.2	2.6.2 DO!!	4
2.2.3	2.6.3 DO!!	4
2.2.4	2.6.4 CHECK!!	4
2.2.5	2.6.5 CHECK!!	4
2.2.6	2.6.5 DO!!	5
2.2.7	2.6.7	5
2.2.8	2.6.8 ASK RON!!	5
2.2.9	2.6.9 DO!!	5
2.2.10	2.6.10 DO!!	5
2.2.11	2.6.11 ASK RON!!	5
3	Chapter 3	5
3.1	Section 3.1	5
3.1.1	3.1.1 DO!!	5
3.1.2	3.1.2	5
3.1.3	3.1.3 DO!!	6
3.1.4	3.1.4 DO!!	6
3.1.5	3.1.5 DO!!	6
3.1.6	3.1.7 DO!!	6
3.1.7	3.1.8 DO!!	6
3.1.8	3.1.12 DO!!	6
3.1.9	3.1.13 DO!!	6
3.2	Section 3.2	6
3.2.1	3.2.1	6
3.2.2	3.2.2	6
3.2.3	3.2.4	6
3.2.4	3.2.5 DO!!	6
3.2.5	3.2.6	6
3.2.6	3.2.7 DO!!	6
3.2.7	3.2.8	6
3.2.8	3.2.9 DO!!	7
3.2.9	3.2.10 CHECK!!	7
3.2.10	3.2.11	7
3.2.11	3.2.12 DO!!	8
3.2.12	3.2.13	8
3.2.13	3.2.14 DO!!	8

	3.2.14	3.2.15	8
	3.2.15	3.2.16	8
3.3	Section 3.3		9
	3.3.1	3.3.1 DO!!	9
	3.3.2	3.3.2 DO!!	9
	3.3.3	3.3.3 DO!!	9
4	Chapter 4		9
4.1	Section 4.1		9
	4.1.1	4.1.1	9
	4.1.2	4.1.2	10
	4.1.3	4.1.3 DO!!	10
	4.1.4	4.1.4	10
	4.1.5	4.1.5 DO!!	10
4.2	Section 4.2		10
	4.2.1	4.2.1 DO!!	10
	4.2.2	4.2.2 DO!!	10
	4.2.3	4.2.3 DO!!	10
	4.2.4	4.2.4 DO!!	10
	4.2.5	4.2.5 DO!!	10
	4.2.6	4.2.6	10
	4.2.7	4.2.7 DO!!	11
	4.2.8	4.2.8	11
	4.2.9	4.2.9 DO!!	11
4.3	Section 4.3		11
	4.3.1	4.3.1 DO!!	11
	4.3.2	4.3.2 DO!!	12
	4.3.3	4.3.3 DO!!	12
	4.3.4	4.3.4 DO!!	12
	4.3.5	4.3.5 CHECK!!	12
	4.3.6	4.3.6	13
	4.3.7	4.3.7 DO!!	13
	4.3.8	4.3.8 DO!!	13
	4.3.9	4.3.9	13
	4.3.10	4.3.10 DO!!	13
4.4	Section 4.4		14
	4.4.1	4.4.1 DO!!	14
	4.4.2	4.4.2	14
	4.4.3	4.4.3 ASK RON!!	14
	4.4.4	4.4.4 ASK RON!!	14
	4.4.5	4.4.5 DO!!	14
	4.4.6	4.4.6 DO!!	14
	4.4.7	4.4.7 DO!!	14
	4.4.8	4.4.8 DO!!	14
	4.4.9	4.4.9	14
	4.4.10	4.4.10	15
	4.4.11	4.4.11	15

4.4.12	4.4.12	16
5	Chapter 5	16
5.1	Section 5.1	16
5.1.1	5.1.1 DO!!	16
5.2	Section 5.2	16
5.2.1	5.2.1 DO!!	16
5.3	Section 5.3	16
5.3.1	5.3.1 DO!!	16
6	Chapter 6	16
6.1	Section 6.1	16
6.1.1	6.1.1	16
6.1.2	6.1.2 DO!!	17
6.1.3	6.1.3	17
7	Extra Questions for Ron	18
7.0.1	1	18
7.0.2	2	18

1 Chapter 1

2 Chapter 2

2.1 Section 2.1

2.1.1 2.2.2

Given an exact sequence of vector bundles,

$$0 \longrightarrow L \xrightarrow{\varphi} E \xrightarrow{\psi} F \longrightarrow 0$$

where L is a line bundle. Consider the exact sequence,

$$0 \longrightarrow L \otimes \bigwedge^{i-1} F \longrightarrow \bigwedge^i E \longrightarrow \bigwedge^i F \longrightarrow 0$$

where the first map is defined by taking sections $s_1 \otimes (f_2 \wedge \cdots \wedge f_i)$ and lifting each f_i to a section e_j of E up to a section s_j of L and then map $s_1 \otimes (f_2 \wedge \cdots \wedge f_i) \mapsto s_1 \wedge e_2 \wedge \cdots \wedge e_i$. This map is well-defined because,

$$\begin{aligned} s_1 \wedge e'_2 \wedge \cdots \wedge e'_i &= s_1 \wedge (e_2 + \varphi(s_2)) \wedge \cdots \wedge (e_i + \varphi(s_i)) = s_1 \wedge e_2 \wedge \cdots \wedge e_n + s_1 \wedge s_2 \wedge \cdots \wedge e_n + \cdots \\ &= s_1 \wedge e_2 \wedge \cdots \wedge e_n \end{aligned}$$

because $s_1 \wedge s_j = 0$ since L is a line bundle. Thus this map is well-defined and clearly $\ker \wedge^i \psi$ is the image of this map because $L \subset E$ maps to zero under ψ thus the kernel is exterior products where one factor is in L . Furthermore, dualizing if we have an exact sequence of vector bundles,

$$0 \longrightarrow F \longrightarrow E \longrightarrow L \longrightarrow 0$$

where L is a line bundle then there is an exact sequence,

$$0 \longrightarrow \bigwedge^i F \longrightarrow \bigwedge^i E \longrightarrow L \otimes \bigwedge^{i-1} F \longrightarrow 0$$

2.1.2 2.2.3

Let E be a holomorphic vector bundle E of rank r there exists a non-degenerate pairing,

$$\bigwedge^k E \times \bigwedge^{r-k} E \rightarrow \det E$$

via $(e_1 \wedge \cdots \wedge e_k, e_{k+1} \wedge \cdots \wedge e_r) \mapsto e_1 \wedge \cdots \wedge e_r$. Locally $E \cong \mathcal{O}_X^{\oplus r}$ and thus the pairing is nondegenerate because we can take e_i to be the standard basis of $\mathcal{O}_X^{\oplus r}$. Then $\det E \cong \mathcal{O}_X$ and $(e_1 \wedge \cdots \wedge e_k, e_{k+1} \wedge \cdots \wedge e_r) \mapsto e_1 \wedge \cdots \wedge e_r$ is a generator of $\det E \cong \mathcal{O}_X$.

Since the above pairing is nondegenerate, we get an isomorphism $\bigwedge^k E \xrightarrow{\sim} \bigwedge^{r-k} E^* \otimes \det E$.

2.1.3 2.2.5

Let L, L^* be holomorphic line bundles on a compact complex manifold X . Suppose that L and L^* admit nonzero global holomorphic sections s, s' . Then consider $s \otimes s'$ a global section of $L \otimes L^* \cong \mathcal{O}_X$. However, all nonzero sections of \mathcal{O}_X are nonvanishing because X is compact and thus $H^0(X, \mathcal{O}_X) = \mathbb{C}$. Therefore, s and s' are nonvanishing meaning that $L \cong L^* \cong \mathcal{O}_X$.

2.1.4 2.2.6 DO!!

2.2 Section 2.6

2.2.1 2.6.1

I think f is holomorphic iff $df(Iv) = idf(v)$

2.2.2 2.6.2 DO!!

2.2.3 2.6.3 DO!!

2.2.4 2.6.4 CHECK!!

Let $f : X \rightarrow Y$ be a surjective holomorphic map between connected complex manifolds. We want to look at the smooth locus of f .

I claim the following is true: for a morphism of vector bundles (not necessarily constant rank) $\phi : \mathcal{E}_1 \rightarrow \mathcal{E}_2$ then ϕ has full rank $k = \min\{m, n\}$ iff the morphism $\phi' : \bigwedge^k \mathcal{E}_1 \rightarrow \bigwedge^k \mathcal{E}_2$ is nonzero (is this true).

Therefore, the locus where ϕ is not full rank is the vanishing of the section

$$\phi' \in \mathcal{HOM}_{\mathcal{O}_X} \left(\bigwedge^k \mathcal{E}_1, \bigwedge^k \mathcal{E}_2 \right)$$

Now apply this to the map $f^* \Omega_Y \rightarrow \Omega_X$ to get the nonsmooth locus.

2.2.5 2.6.5 CHECK!!

The cousins' problem has a solution because $H^1(X, \mathcal{O}_X) = 0$. Question: why is every hypersurface defined by a $H^0(K^\times / \mathcal{O}_X^\times)$. Question: how are we supposed to use the pincare lemma.

2.2.6 2.6.5 DO!!

2.2.7 2.6.7

We define,

$$H_{\text{BC}}^{p,q}(X) = \frac{\{\alpha \in \mathcal{A}^{p,q}(X) \mid d\alpha = 0\}}{\partial\bar{\partial}\mathcal{A}^{p-1,q-1}(X)}$$

This makes sense because if $\alpha = \partial\bar{\partial}\gamma$ then

$$d\alpha = \partial^2\bar{\partial}\gamma - \bar{\partial}^2\partial\gamma = 0$$

Now, the inclusion of d-closed forms into $\bar{\partial}$ -closed forms induces a map,

$$H_{\text{BC}}^{p,q}(X) \rightarrow H^{p,q}(X)$$

which is well-defined because if $\alpha = \partial\bar{\partial}\gamma$ then $\alpha = -\bar{\partial}\partial\gamma$ and is thus $\bar{\partial}$ -exact. If X is furthermore compact Kahler then by the $\partial\bar{\partial}$ -lemma we see if α maps to zero i.e. $\alpha = \partial\bar{\partial}\beta$ and $d\alpha = 0$ then $\alpha = d\gamma$ so the map is injective. Furthermore, by the Hodge decomposition, $H^{p,q}(X)$ can be represented by Harmonic forms which are d-closed and thus this map is surjective as well.

2.2.8 2.6.8 ASK RON!!

Is this just because we can take $M \rightarrow M$ via complex conjugation.

2.2.9 2.6.9 DO!!

2.2.10 2.6.10 DO!!

2.2.11 2.6.11 ASK RON!!

3 Chapter 3

3.1 Section 3.1

3.1.1 3.1.1 DO!!

Let X be a complex manifold with an almost complex structure (M, I) . We need to find a Riemannian structure g on M such that g is compatible with I meaning $g(I-, I-) = g(-, -)$. (FINISH)

3.1.2 3.1.2

Let X be a connected complex manifold of dimension $n > 1$ and let g be a Kähler metric. Suppose that $g' = e^f \cdot g$ for some real smooth function $f \in \mathcal{A}^0(X)$ is also a Kähler metric. Then the associated Kähler forms satisfy $\omega' = e^f \cdot \omega$. Since both are Kähler forms, we must have $d\omega' = 0$ and $d\omega = 0$. However,

$$d\omega' = e^f df \wedge \omega + e^f d\omega = e^f df \wedge \omega$$

Therefore, $df \wedge \omega = 0$ and thus $L(df) = 0$. However, since $n > 1$ the Lefschetz operator is injective on k -forms for $k < n$ and thus $df = 0$. Since X is connected then f is constant so $\omega' = c\omega$.

3.1.3 3.1.3 DO!!

3.1.4 3.1.4 DO!!

3.1.5 3.1.5 DO!!

3.1.6 3.1.7 DO!!

3.1.7 3.1.8 DO!!

3.1.8 3.1.12 DO!!

3.1.9 3.1.13 DO!!

3.2 Section 3.2

3.2.1 3.2.1

3.2.2 3.2.2

3.2.3 3.2.4

What does this really mean?? Ask Ron.

3.2.4 3.2.5 DO!!

3.2.5 3.2.6

Let X be a compact Kähler manifold. Then,

$$H^n(X, \mathbb{C}) = \bigoplus_{p+q=n} H^{p,q}(X)$$

Furthermore, $H^{q,p} = \overline{H^{p,q}}$. Therefore,

$$b_{2k+1} = \sum_{p+q=2k+1} h^{p,q} = \sum_{i=0}^k (h^{2k+1-i,i} + h^{i,2k+1-i}) = 2 \sum_{i=0}^k h^{2k+1-i,i}$$

is even.

3.2.6 3.2.7 DO!!

No! (PROVE IT)

3.2.7 3.2.8

Let X be a compact Kähler manifold. Let $\omega \in H^0(X, \Omega_X^p)$. Clearly, $\bar{\partial}\omega = 0$ since ω is a holomorphic $(p, 0)$ -form. Furthermore,

$$\bar{\partial}^* \omega = -(\bar{\star} \circ \bar{\partial} \circ \bar{\star}) \omega$$

but $\bar{\star}\omega$ is a $(n-p, n)$ -form and thus $\bar{\partial}\bar{\star}\omega = 0$. Therefore, $\bar{\partial}\omega = 0$ and $\bar{\partial}^* \omega = 0$ and thus $\Delta_{\bar{\partial}} \omega = 0$.

3.2.8 3.2.9 DO!!

3.2.9 3.2.10 CHECK!!

Let (X, g) be a compact hermitian manifold. Show that any d-harmonic (p, q) -form is also $\bar{\partial}$ -harmonic.

Since α is d-harmonic, we have $d\alpha = 0$ and $d^*\alpha = 0$. Therefore, $\partial\alpha = 0$ and $\bar{\partial}\alpha = 0$ and $\partial^*\alpha = 0$ and $\bar{\partial}^*\alpha = 0$. Therefore, $\Delta_\partial(\alpha) = 0$ and $\Delta_{\bar{\partial}}(\alpha) = 0$.

3.2.10 3.2.11

Let $X = \mathbb{P}^n$. Consider the Euler sequence,

$$0 \longrightarrow \Omega_X \longrightarrow \mathcal{O}_X(-1)^{\oplus(n+1)} \longrightarrow \mathcal{O}_X \longrightarrow 0$$

Then we may apply exterior powers to get the following sequence,

$$0 \longrightarrow \Omega_X^p \longrightarrow \bigwedge^p \mathcal{O}_X(-1)^{\oplus(n+1)} \longrightarrow \mathcal{O}_X \otimes_{\mathcal{O}_X} \Omega_X^{p-1} \longrightarrow 0$$

However,

$$\bigwedge^p \mathcal{O}_X(-1)^{\oplus(n+1)} = \mathcal{O}_X(-p)^{\oplus \binom{n+1}{p}}$$

and thus we have the sequence,

$$0 \longrightarrow \Omega_X^p \longrightarrow \mathcal{O}_X(-p)^{\oplus \binom{n+1}{p}} \longrightarrow \Omega_X^{p-1} \longrightarrow 0$$

Now applying the cohomology sequence we find,

$$H^{q-1}(X, \mathcal{O}_X(-p))^{\oplus \binom{n+1}{p}} \longrightarrow H^{q-1}(X, \Omega_X^{p-1}) \longrightarrow H^q(X, \Omega_X^p) \longrightarrow H^q(X, \mathcal{O}_X(-p))^{\oplus \binom{n+1}{p}}$$

Therefore, if $0 < q < n$ and $p > 0$ then $H^{q-1}(X, \Omega_X^{p-1}) \xrightarrow{\sim} H^q(X, \Omega_X^p)$. Furthermore, if $q = 0$ and $p > 0$ then $H^0(X, \Omega_X^p) = 0$ because we get an exact sequence,

$$0 \longrightarrow H^0(X, \Omega_X^p) \longrightarrow H^0(X, \mathcal{O}_X(-p))^{\oplus \binom{n+1}{p}}$$

and $H^0(X, \mathcal{O}_X(-p)) = 0$. Finally, if $q = n$ and $p < n + 1$ (which it must) then $H^{q-1}(X, \Omega_X^{p-1}) \xrightarrow{\sim} H^q(X, \Omega_X^p)$ because $H^n(X, \mathcal{O}_X(-p)) = H^0(X, \mathcal{O}_X(p - n - 1))$ by Serre duality.

To finish the base case $p = 0$ we know,

$$H^q(X, \mathcal{O}_X) = \begin{cases} \mathbb{C} & q = 0 \\ 0 & q > 0 \end{cases}$$

Therefore, by induction, $H^p(X, \Omega_X^p) = H^{p-1}(X, \Omega_X^{p-1}) = \mathbb{C}$ for $p \leq n$. Furthermore, if $p \neq q$ then reducing via $H^{q-1}(X, \Omega_X^{p-1}) \xrightarrow{\sim} H^q(X, \Omega_X^p)$ we get to either $H^q(X, \Omega_X^p) = 0$ with $q > 0$ or $H^0(X, \Omega_X^p) = 0$ with $p > 0$. Therefore,

$$H^q(X, \Omega_X^p) = \begin{cases} \mathbb{C} & p = q \leq n \\ 0 & p \neq q \end{cases}$$

Now consider the exponential sequence,

$$H^0(X, \mathcal{O}_X^\times) \longrightarrow H^1(X, \mathbb{Z}) \longrightarrow H^1(X, \mathcal{O}_X) \longrightarrow H^1(X, \mathcal{O}_X^\times) \longrightarrow H^2(X, \mathbb{Z}) \longrightarrow H^2(X, \mathcal{O}_X)$$

but $H^1(X, \mathcal{O}_X) = H^2(X, \mathcal{O}_X) = 0$ and $H^1(X, \mathcal{O}_X^\times) = \text{Pic}(X)$. Therefore, the exponential sequence defines an isomorphism $\text{Pic}(X) \xrightarrow{\sim} H^2(X, \mathbb{Z})$. By the Kähler decomposition $H^2(X, \mathbb{C}) = H^{1,1}(X) = H^1(X, \Omega_X^1)$ and thus $\dim_{\mathbb{C}} H^2(X, \mathbb{C}) = 1$. Therefore, $H^2(X, \mathbb{Z}) = \mathbb{Z}$ because $H^2(X, \mathbb{Z})$ is torsion-free since \mathbb{P}^n is simply connected and therefore $H_1(X, \mathbb{Z}) = 0$.

3.2.11 3.2.12 DO!!

3.2.12 3.2.13

Let X be a compact Kähler manifold and $\alpha \in \mathcal{A}^k(X)$ which is d-closed and d^c-exact where d^c = $i(\bar{\partial} - \partial)$. Notice that $dd^c = 2i\partial\bar{\partial}$. Write $\alpha = \alpha^{k,0} + \dots + \alpha^{0,k}$. Since $d\alpha = 0$ and $d^c\alpha = 0$ we see that $\partial\alpha = 0$ and $\bar{\partial}\alpha = 0$ and this must be true for each $\alpha^{p,q}$ because $\Pi^{p+1,q}\partial\alpha = \partial\alpha^{p,q}$ etc. Then by the $\partial\bar{\partial}$ -lemma we know $\alpha^{p,q} = \partial\bar{\partial}\beta$ for $\beta \in \mathcal{A}^{p-1,q-1}(X)$. Therefore,

$$\alpha = \alpha^{k,0} + \dots + \alpha^{0,k} = \partial\bar{\partial}(\beta^{k,0} + \dots + \beta^{0,k}) = -\frac{i}{2}dd^c\beta$$

where $\beta = \beta^{k,0} + \dots + \beta^{0,k}$.

3.2.13 3.2.14 DO!!

DO! LOOK AT PREVIOUS BC PROBLEM AND CORRESPOND

3.2.14 3.2.15

Let (X, g) be a compact hermitian manifold and let $[\alpha] \in H^{p,q}(X)$ be a cohomology class. If we deform $\alpha' = \alpha + t\bar{\partial}\beta$ by an exact form such that $[\alpha'] = [\alpha]$. Then consider the norm,

$$\|\alpha'\|^2 = \langle \alpha + t\bar{\partial}\beta, \alpha + t\bar{\partial}\beta \rangle = \|\alpha\|^2 + 2t\Re\langle \alpha, \bar{\partial}\beta \rangle + t^2\|\bar{\partial}\beta\|^2$$

Suppose that α has minimal norm then the linear term must be zero so $\Re\langle \alpha, \bar{\partial}\beta \rangle = 0$. Likewise, replacing β by $i\beta$ then $\Re\langle \alpha, \bar{\partial}i\beta \rangle = \text{Im}(\langle \alpha, \bar{\partial}\beta \rangle)$ so we must have $\langle \alpha, \bar{\partial}\beta \rangle = 0$. However, $\langle \alpha, \bar{\partial}\beta \rangle = \langle \bar{\partial}^*\alpha, \beta \rangle = 0$ for arbitrary $(p, q-1)$ -forms β . In particular, we can take,

$$\|\bar{\partial}^*\alpha\|^2 = \langle \bar{\partial}^*\alpha, \bar{\partial}^*\alpha \rangle = 0$$

so $\bar{\partial}^*\alpha = 0$. Furthermore $\bar{\partial}\alpha = 0$ since it defines a cohomology class. Thus $\Delta_{\bar{\partial}}\alpha = 0$ so the representatives of minimal norm are exactly the harmonic representatives.

3.2.15 3.2.16

Let X be a compact Kähler manifold. Let ω and ω' be Kähler forms such that $[\omega] = [\omega'] \in H^2(X, \mathbb{R})$. Then $\eta = \omega - \omega' = d\alpha$ for some real 1-form α . Thus η is a closed real $(1, 1)$ -form which is d-exact and thus by the $\partial\bar{\partial}$ -lemma $\eta = i\partial\bar{\partial}f$ for some $f \in \mathcal{A}^{0,0}$. Notice,

$$\bar{\eta} = -i\bar{\partial}\partial\bar{f} = i\partial\bar{\partial}\bar{f}$$

however η is real so $\bar{\eta} = \eta$ and thus $\bar{f} = f$ so $f \in \mathcal{A}_{\mathbb{R}}^0$ is a real function and,

$$\omega = \omega' + i\partial\bar{\partial}f$$

3.3 Section 3.3

3.3.1 3.3.1 DO!!

3.3.2 3.3.2 DO!!

3.3.3 3.3.3 DO!!

4 Chapter 4

4.1 Section 4.1

4.1.1 4.1.1

Let L be a holomorphic line bundle globally generated by sections $s_1, \dots, s_k \in H^0(X, L)$. Then L admits a canonical hermitian structure h defined by locally choosing a trivialization, $\phi : L|_U \xrightarrow{\sim} \mathcal{O}_U$ then for any local sections $\alpha, \beta \in \mathcal{L}(U)$,

$$h(\alpha, \beta) = \frac{\psi(\alpha) \cdot \overline{\psi(\beta)}}{\sum_i |\psi(s_i)|^2}$$

This is well-defined because for any other choice of local trivialization $\psi' : L|_U \xrightarrow{\sim} \mathcal{O}_U$ gives a transition function $t = \psi' \circ \psi^{-1} : \mathcal{O}_U \rightarrow \mathcal{O}_U$ which is a holomorphic function on U . Then $\psi' = (\psi' \circ \psi^{-1}) \circ \psi = t\psi$ and therefore,

$$\frac{\psi'(\alpha) \cdot \overline{\psi'(\beta)}}{\sum_i |\psi'(s_i)|^2} = \frac{t\psi(\alpha) \cdot \overline{t\psi(\beta)}}{\sum_i |t\psi(s_i)|^2} = \frac{|t|^2 \psi(\alpha) \cdot \overline{\psi(\beta)}}{|t|^2 \sum_i |\psi(s_i)|^2} = \frac{\psi(\alpha) \cdot \overline{\psi(\beta)}}{\sum_i |\psi(s_i)|^2}$$

Then the dual bundle L^* obtains a natural hermitian structure h^* defined on $\alpha, \beta \in L^*(U)$ i.e. maps $L|_U \rightarrow \mathcal{O}_U$ via,

$$h^*(\alpha, \beta) = \alpha(h^{-1}(\beta))$$

viewing h as an \mathbb{C} -antilinear isomorphism $h : \mathcal{L} \xrightarrow{\sim} \mathcal{L}^*$. Furthermore, there is an inclusion $L^* \hookrightarrow \mathcal{O}_X^{\oplus k}$ dual to $\mathcal{O}_X^{\oplus k} \twoheadrightarrow \mathcal{L}$ defined by applying $\alpha \in L^*(U)$ to s_1, \dots, s_n . By restricting the standard hermitian structure on $\mathcal{O}_X^{\oplus k}$ to L^* we get a hermitian structure h' on \mathcal{L} . Explicitly, for $\alpha, \beta \in L^*(U)$,

$$h'(\alpha, \beta) = \sum_i \alpha(s_i) \overline{\beta(s_i)}$$

However, choosing a local trivialization $\psi : \mathcal{L}|_U \xrightarrow{\sim} \mathcal{O}_U$ then $\psi^* : \mathcal{O}_U \xrightarrow{\sim} \mathcal{L}^*|_U$ given by $\psi^*(f) = f\psi$ is an isomorphism. We explicitly find,

$$h^{-1}(\psi^*(f)) = \bar{f}h^{-1}(\psi) = f \sum_i s_i \cdot \overline{\psi(s_i)}$$

Therefore,

$$h^*(\psi^*(f), \psi^*(g)) = f\bar{g}\psi(h^{-1}(\beta)) = f\bar{g} \sum_i |\psi(s_i)|^2 = h'(\psi^*(f), \psi^*(g))$$

Thus $h^* = h'$.

4.1.2 4.1.2

Let L be a holomorphic line bundle of degree $d > 2g(C) - 2$ on a curve C where $\deg K_C = 2g(C) - 2$. By Serre duality $H^1(C, L) = H^0(X, L^* \otimes K_C) = 0$ since $\deg(L^* \otimes K_C) = 2g(C) - 2 - d < 0$. In particular if L is a line bundle with $\deg L > 0$ then $H^1(C, K_C \otimes L) = 0$.

4.1.3 4.1.3 DO!!

Let $X = \mathbb{P}^n$. To compute the cohomology $H^q(X, \mathcal{O}_X(k))$ consider the

4.1.4 4.1.4

Let E be a hermitian holomorphic vector bundle on a compact Kähler manifold X . Consider a global holomorphic section $s \in H^0(X, \Omega^p \otimes E)$. Then we know,

$$\Delta_E(s) = 0 \iff \bar{\partial}_E s = 0 \text{ and } \bar{\partial}_E^* s = 0$$

Notice that $\Omega^p \otimes E$ is the kernel of $\bar{\partial}_E : \mathcal{A}^{p,0}(X, E) \rightarrow \mathcal{A}^{p,1}(X, E)$ therefore $\bar{\partial}_E s = 0$ automatically. Furthermore,

$$\bar{\partial}_E^* s = -\bar{\star}_E \bar{\partial}_{E^*} \bar{\star}_E s$$

However $\bar{\star}_E s \in \mathcal{A}^{n-p,n}(X, E^*)$ and thus $\bar{\partial}_{E^*} \bar{\star}_E s \in \mathcal{A}^{n-p,n+1}(X, E^*) = 0$. Therefore, $\bar{\partial}_E^* s = 0$ so $\nabla_E s = 0$ and thus s is Harmonic.

4.1.5 4.1.5 DO!!

4.2 Section 4.2

4.2.1 4.2.1 DO!!

4.2.2 4.2.2 DO!!

4.2.3 4.2.3 DO!!

4.2.4 4.2.4 DO!!

4.2.5 4.2.5 DO!!

Let (E, h) be a hermitian vector bundle and suppose $E = E_1 \oplus E_2$. Then E_1, E_2 are hermitian with hermitian structures h_1 and h_2 induced by the inclusions. Now let ∇_1 and ∇_2 be the induced connections.

4.2.6 4.2.6

Let E be a vector bundle and ∇ a connection on E . Then there is an induced connection $\nabla' : \bigwedge^2 E \rightarrow \Omega_X^1 \otimes \bigwedge^2 E$ as the quotient of the induced connection on $E^{\otimes 2}$. Explicitly,

$$\nabla'(s_1 \wedge s_2) = \nabla s_1 \wedge s_2 + s_1 \wedge \nabla s_2$$

This is well-defined because,

$$\nabla'(s \wedge s) = \nabla s \wedge s + s \wedge \nabla s = 0$$

Furthermore, there is an induced connection $\nabla' : \det E \rightarrow \Omega_X^1 \otimes \det E$ as the quotient of the induced connection on $E^{\otimes n}$. Explicitly,

$$\nabla'(s_1 \wedge \cdots \wedge s_n) = \nabla s_1 \wedge \cdots \wedge s_n + \cdots + s_1 \wedge \cdots \wedge \nabla s_n$$

which is well-defined as above. Let e_i be a local frame of E then we can write $\nabla = d + A$ where A is a matrix of 1-forms which really means,

$$\nabla(f_j e_j) = df_j \otimes e_j + f_j \nabla e_j = df_j \otimes e_j + f_j A_{ij} \otimes e_i = (df_i + A_{ij} f_j) \otimes e_i$$

where $A_{ij} \otimes e_i = \nabla e_j$. Then we find that,

$$\begin{aligned} \nabla'(e_1 \wedge \cdots \wedge e_n) &= A_{1k} \otimes e_k \wedge \cdots \wedge e_n + \cdots + e_1 \wedge \cdots \wedge A_{nk} \otimes e_k = (A_{11} + \cdots + A_{nn}) \otimes (e_1 \wedge \cdots \wedge e_n) \\ &= \text{tr}(A) \otimes (e_1 \wedge \cdots \wedge e_n) \end{aligned}$$

Therefore, the connection $\nabla' : \det E \rightarrow \Omega_X^1 \otimes \det E$ on the line bundle $\det E$ is locally given by the 1-form $\text{tr}(A)$.

4.2.7 4.2.7 DO!!

4.2.8 4.2.8

First note that if ∇_1 and ∇_2 are connections on E_1 and E_2 we define a connection $\nabla : E_1 \otimes E_2 \rightarrow \Omega \otimes (E_1 \otimes E_2)$ via $\nabla(s_1 \otimes s_2) = \nabla_1 s_1 \otimes s_2 + s_1 \otimes \nabla_2 s_2$ and also $\nabla : \text{Hom}_{\mathcal{O}_X}(E_1, E_2) \rightarrow \Omega \otimes \text{Hom}_{\mathcal{O}_X}(E_1, E_2)$ via $\varphi \mapsto \nabla \varphi$ such that $(\nabla \varphi)(s) = \nabla_2 \varphi(s) - \varphi(\nabla_1(s))$ for any $s \in \Gamma(U, E)$.

Therefore, if ∇ is a connection on E define the dual connection $\nabla^* : E^* \rightarrow \Omega_X^1 \otimes E^*$ on E^* via sending $\varphi : E \rightarrow \mathcal{O}_X$ so $(\nabla^* \varphi)(s) = d\varphi(s) - \varphi(\nabla s)$.

Consider the induced connection ∇ on $(E \otimes \bar{E})^*$. Then for any section $h \in \Gamma(U, (E \otimes \bar{E})^*)$, for instance a hermitian metric, we get ∇h such that for any $s_1, s_2 \in E$,

$$\begin{aligned} (\nabla h)(s_1 \otimes \bar{s}_2) &= dh(s_1 \otimes \bar{s}_2) - h(\nabla(s_1 \otimes \bar{s}_2)) = dh(s_1 \otimes \bar{s}_2) - h(\nabla s_1 \otimes \bar{s}_2) - h(s_1 \otimes \bar{\nabla} \bar{s}_2) \\ &= dh(s_1 \otimes \bar{s}_2) - h(\nabla s_1 \otimes \bar{s}_2) - h(s_1 \otimes \overline{\nabla s_2}) \end{aligned}$$

Therefore, as a metric,

$$(\nabla h)(s_1, s_2) = dh(s_1, s_2) - h(\nabla s_1, s_2) - h(s_1, \nabla s_2)$$

Therefore, ∇ is hermitian with respect to the hermitian structure (E, h) iff $\nabla h = 0$.

4.2.9 4.2.9 DO!!

4.3 Section 4.3

4.3.1 4.3.1 DO!!

Consider $\nabla^2 : \mathcal{A}^k(E) \rightarrow \mathcal{A}^{k+2}(E)$ on $\omega \otimes s$ where ω is a k -form and s is a section of E . Then,

$$\nabla(\omega \otimes s) = d\omega \otimes s + (-1)^k \omega \wedge \nabla(s)$$

and thus,

$$\begin{aligned}\nabla^2(\omega \otimes s) &= d\omega \otimes s + (-1)^{k+1} d\omega \wedge \nabla s + (-1)^k d\omega \wedge \nabla(s) + (-1)^{2k} \omega \wedge \nabla^2(s) \\ &= \omega \wedge \nabla^2(s)\end{aligned}$$

since the $dd\omega = 0$ and the middle terms cancel. Here we use the generalized Leibniz formula,

$$\nabla(\omega \wedge \alpha) = d\omega \wedge \alpha + (-1)^k \omega \wedge \nabla \alpha$$

where ω is a k -form and $\alpha \in \mathcal{A}^\ell(E)$. Furthermore $\nabla^2(s) = F_\nabla(s)$ so we find,

$$\nabla^2(\omega \otimes s) = \omega \wedge F_\nabla(s)$$

and therefore for a general E -valued k -form α we see that $\nabla^2(\alpha) = \text{tr}(\omega \wedge F_\nabla)$ where we view $\omega \in H^0(X, \Omega_X^k \otimes E)$ and $F_\nabla \in H^0(X, \Omega_X^2 \otimes E \otimes E^*)$ and taking the map $\wedge : \Omega_X^k \otimes \Omega_X^2 \rightarrow \Omega_X^{k+2}$ and contracting the E^* from F_∇ with the E from ω .

4.3.2 4.3.2 DO!!

4.3.3 4.3.3 DO!!

Let (E_1, h_1) and (E_2, h_2) be two hermitian holomorphic vector bundles endowed with hermitian connections ∇_1, ∇_2 such that the curvature of both is (semi)-positive.

- (a) The curvature of ∇^* on E^* is $F_{\nabla^*} = -F_\nabla^\top$ which is (semi)-negative since F_∇ is (semi)-positive.
- (b) The curvature of ∇ on $E_1 \otimes E_2$ is $F_\nabla = F_{\nabla_1} \otimes \text{id} + \text{id} \otimes F_{\nabla_2}$ which is (semi)-positive since F_{∇_1} and F_{∇_2} are. Furthermore if one of F_{∇_i} is positive then F_∇ is positive since the other term is nonnegative.
- (c) The curvature of ∇ on $E_1 \oplus E_2$ is $F_\nabla = F_{\nabla_1} \oplus F_{\nabla_2}$ which is (semi)-positive since F_{∇_1} and F_{∇_2} are.

4.3.4 4.3.4 DO!!

4.3.5 4.3.5 CHECK!!

Let X be complex manifold. Let L be a holomorphic line bundle with a hermitian structure h whose Chern connection has positive curvature. Then $F_\nabla \in \mathcal{A}^{1,1}(X)$ is an imaginary $(1,1)$ -form. Furthermore, note that $F_\nabla = \bar{\partial}\partial \log h$ and thus,

$$dF_\nabla = (\partial + \bar{\partial})\bar{\partial}\partial \log h = 0$$

because $\bar{\partial}^2 = 0$ and $\partial\bar{\partial}\partial = -\partial^2\bar{\partial} = 0$. Since $\omega = iF_\nabla$ is positive, it is a Kähler form. Furthermore if X is compact then,

$$\int_X A(L)^n = \int_X F_\nabla^n = \int_X \omega^n = n! \int_X \text{vol}_\omega > 0$$

(CHECK THIS! FACTORS OF I)

4.3.6 4.3.6

Let $X = \mathbb{P}^n$ and $\omega_X = \mathcal{O}_X(-n-1)$ be the canonical bundle. The sections $x_0, \dots, x_n \in \Gamma(X, \mathcal{O}_X(1))$ define a canonical hermitian structure h on $\mathcal{O}_X(1)$ which has the property that $\frac{i}{2\pi}F_\nabla = \omega_{\text{FS}}$ which is positive. Then $\omega_X = \mathcal{O}_X(-n-1)$ has a canonical hermitian structure $(h^*)^{\otimes n+1}$ which has curvature $\frac{i}{2\pi}F_{\nabla'} = -(n+1)\omega_{\text{FS}}$ which is therefore negative.

4.3.7 4.3.7 DO!!

4.3.8 4.3.8 DO!!

4.3.9 4.3.9

Let X be a compact Kähler manifold with $b_1(X) = 0$. Suppose that ∇ is a flat connection on \mathcal{O}_X with $\nabla^{0,1} = \bar{\partial}$. Then $\nabla = d + \omega$ where $\omega : \mathcal{A}^0(X) \rightarrow \mathcal{A}^1(X)$ is $\mathcal{A}^0(X)$ -linear and thus $\omega \in \mathcal{A}^1(X)$. Furthermore, $\nabla^{0,1} = \bar{\partial}$ so ω is a smooth $(1,0)$ -form. Now consider the curvature,

$$F_\nabla = \nabla \circ \nabla(1) = \nabla(\omega \otimes 1) = d\omega \otimes 1 - \omega \wedge \nabla(1) = d\omega \otimes 1 - \omega \wedge \omega \otimes 1 = d\omega$$

Since ∇ is flat we must have $d\omega = 0$. Thus ω defines a de Rham cohomology class $[\omega] \in H^1(X, \mathbb{C})$ but $b_1(X) = 0$ so ω is exact. Take $\omega = df$ for some smooth function f . However, ω is a $(1,0)$ -form so f is holomorphic. But X is compact so f is constant and thus $\omega = 0$ showing that $\nabla = d$.

Now suppose that L is a line bundle on X with $c_1(L) = 0$. From the exponential sequence,

$$H^1(X, \mathcal{O}_X) \longrightarrow \text{Pic}(X) \xrightarrow{c_1} H^2(X, \mathbb{Z})$$

and thus $\ker c_1 = \text{Im}(H^1(X, \mathcal{O}_X) \rightarrow \text{Pic}(X))$. However, $b_1(X) = 0$ so by the Kähler decomposition, $H^1(X, \mathcal{O}_X) = 0$. Therefore, $\ker c_1$ is trivial so $L = \mathcal{O}_X$.

4.3.10 4.3.10 DO!!

Let ∇ be a connection on a complex vector bundle E . We want to show that E locally has parallel frames iff $F_\nabla = 0$.

Suppose that E has a local frame e_1, \dots, e_n of parallel sections over U i.e. $\nabla e_i = 0$ and these are independent on each fiber. Since the curvature form $\omega_\nabla(s) = \nabla_1 \circ \nabla(s)$ is \mathcal{O}_X -linear, writing $s = f_i e_i$ we get,

$$\omega_\nabla(f_i e_i) = f_i \omega_\nabla(e_i) = f_i \nabla_1 \circ \nabla e_i = 0$$

Therefore, $\omega_\nabla = 0$ so ∇ must be flat.

Locally write $E|_U \cong \mathcal{O}_U^{\oplus n}$ write e_i for a local frame of $E|_U$. Now write $\nabla e_j = \omega_{ij} \otimes e_i$ thus we see,

$$\nabla(f_j e_j) = df_j \otimes e_j + \omega_{ij} f_j \otimes e_i = (df_j + \omega_{ij} f_j) \otimes e_i$$

Now, applying $\nabla_1 : \Omega_X^1 \otimes E \rightarrow \Omega_X^2 \otimes E$ we get,

$$\begin{aligned} \nabla_1 \circ \nabla(f_j e_j) &= \nabla_1(df_j + \omega_{ij} f_j) \otimes e_i = ddf_j \otimes e_i + d(\omega_{ij} f_j) \otimes e_i - (df_j + \omega_{ij} f_j) \wedge \nabla e_i \\ &= (d\omega_{ij} f_j - \omega_{ij} \wedge df_j) \otimes e_i - (df_j + \omega_{ij} f_j) \wedge \omega_{ki} \otimes e_k \\ &= d\omega_{ij} f_j \otimes e_i + df_j \wedge \omega_{ij} \otimes e_i - df_j \wedge \omega_{ki} \otimes e_k + \omega_{ki} \wedge \omega_{ij} f_j \otimes e_k \\ &= (d\omega_{ij} + \omega_{ik} \wedge \omega_{kj}) f_j \otimes e_i \end{aligned}$$

Therefore,

$$\omega_{\nabla}(f_j e_j) = (d\omega_{ij} + \omega_{ik} \wedge \omega_{kj}) f_j \otimes e_i$$

is linear as it should be. Now assume ∇ is flat i.e. $\omega_{\nabla} = 0$. Thus,

$$d\omega_{ij} + \omega_{ik} \wedge \omega_{kj} = 0$$

First, in the case $n = 1$ the connection is given by a 1-form ω . Then $\omega_{\nabla} = 0 \iff d\omega = 0$ in which case locally $\omega = -df$ and thus $\nabla(fe) = df \otimes e + \omega \otimes e = 0$ so we get a frame of parallel sections.

Now we proceed by induction for the general case. First, using a $\mathrm{GL}_{\mathbb{C}}(n)$ transformation we can

Assume we can find a frame e_1, \dots, e_{n-1}, s such that $\nabla e_i = 0$. (FINISH)

4.4 Section 4.4

4.4.1 4.4.1 DO!!

4.4.2 4.4.2

Let X be a compact complex manifold and L a basepoint-free line bundle. Then L defines a map $f : X \rightarrow \mathbb{P}^N$ such that $f^* \mathcal{O}_{\mathbb{P}^N}(1) = L$. Let h be the standard hermitian structure on $\mathcal{O}_{\mathbb{P}^N}(1)$ so f^*h gives a hermitian structure on L . Taking the Chern connections $\nabla_{f^*h} = f^* \nabla_h$ and thus,

$$F(L, f^*h) = F(f^* \mathcal{O}_{\mathbb{P}^N}(1), f^*h) = f^* F(\mathcal{O}_{\mathbb{P}^N}(1), h) = f^* \omega_{\mathrm{FS}}$$

which is a positive form. Therefore,

$$c_1(L) = f^*[\omega_{\mathrm{FS}}]$$

so we see that,

$$\int_X c_1(L)^n = \int_X (f^* \omega_{\mathrm{FS}})^n = \int_X f^* \omega_{\mathrm{FS}}^n \geq 0$$

4.4.3 4.4.3 ASK RON!!

4.4.4 4.4.4 ASK RON!!

Ask Ron about interpretation!!

4.4.5 4.4.5 DO!!

4.4.6 4.4.6 DO!!

4.4.7 4.4.7 DO!!

4.4.8 4.4.8 DO!!

4.4.9 4.4.9

Note that $\mathrm{End}(E) \cong E^* \otimes E$ then,

$$c_k(\mathrm{End}(E)) = \sum_{i+j=k} c_i(E^*) \cdot c_j(E) = \sum_{i+j} (-1)^i c_i(E) \cdot c_j(E)$$

In particular,

$$c_1(\mathrm{End}(E)) = c_0(E) \cdot c_1(E) - c_1(E) \cdot c_0(E) = 0$$

and likewise,

$$c_2(\text{End}(E)) = c_0(E) \cdot c_2(E) - c_1(E) \cdot c_1(E) + c_2(E) \cdot c_0(E) = 2c_2(E) - c_1(E)^2$$

Then if $E = L \oplus L$ where L is a line bundle we have,

$$c(L) = 1 + c_1(L)$$

and thus,

$$c_1(E) = 2c_1(L) \quad \text{and} \quad c_2(E) = c_1(L)^2$$

Therefore, we see that,

$$(4c_2 - c_1^2)(E) = 4c_1(E)^2 - (4c_1(E))^2 = 0$$

Furthermore, if $E \cong E^*$ then $c_{2k+1}(E) = c_{2k+1}(E^*) = (-1)^{2k+1}c_{2k+1}(E) = -c_{2k+1}(E)$ and thus $c_{2k+1}(E) = 0$.

4.4.10 4.4.10

Let L be a holomorphic line bundle on X a compact Kähler manifold. Suppose that $c_1(L) = [\alpha]$ where α is closed a real $(1, 1)$ -form. Let h_0 be a Hermitian structure on L then,

$$c_1(L, h_0) = \frac{i}{2\pi} \bar{\partial} \partial \log h_0$$

Now consider,

$$\eta = \alpha - c_1(L, h_0)$$

is a real $(1, 1)$ -form and since $[\alpha] = [c_1(L, h_0)]$ also η is d-exact. Thus, by the $\partial\bar{\partial}$ -lemma, we know,

$$\eta = -\frac{i}{2\pi} \partial \bar{\partial} f$$

for $f \in \mathcal{A}_{\mathbb{R}}^{0,0}(X)$ i.e. f is a real smooth function. Therefore,

$$\alpha = \frac{i}{2\pi} \bar{\partial} \partial [f + \log h_0] = \frac{i}{2\pi} \bar{\partial} \partial \log e^f h_0$$

Therefore, let $h = e^f h_0$ be another Hermitian structure (since f is real) then we see $c_1(L, h) = \alpha$.

4.4.11 4.4.11

Let X be compact Kähler and E a vector bundle with a Chern connection ∇ . If we let,

$$\sum_{i=0}^r \tilde{P}_i(B) = \text{tr}(e^B) = \sum_{n=0}^{\infty} \frac{1}{n!} \text{tr}(B^n)$$

so,

$$\tilde{P}_k(B_1, \dots, B_k) = \frac{1}{k!} \text{tr}(B_1 \cdots B_k)$$

and then define,

$$\text{ch}_k(E, \nabla) := \tilde{P}_k\left(\frac{i}{2\pi} F_{\nabla}\right) \in \mathcal{A}_{\mathbb{C}}^{2k}(M)$$

where \tilde{P}_k acts on $\text{End}(E)$ -valued 2-forms via,

$$\tilde{P}_k(\alpha_1 \otimes \varphi_1, \dots, \alpha_k \otimes \varphi_k) = (\alpha_1 \wedge \dots \wedge \alpha_k) \tilde{P}_k(\varphi_1, \dots, \varphi_k) = (\alpha_1 \wedge \dots \wedge \alpha_k) \frac{1}{k!} \text{tr}(\varphi_1 \dots \varphi_k)$$

This is the composition of $(\Omega_X^2)^{\otimes k} \rightarrow \Omega_X^{2k}$ via exterior product and $\text{End}(E)^{\otimes k} \rightarrow \text{End}(E)$ via composition and finally taking trace. We see that,

$$\text{ch}_k(E, \nabla) = \frac{1}{k!} \left(\frac{i}{2\pi} \right)^k \text{tr}(F_{\nabla}^{\otimes k})$$

where $F_{\nabla}^{\otimes k}$ is the image under $(\Omega_X^2 \otimes \text{End}(E))^{\otimes k} \rightarrow \Omega_X^{2k} \otimes \text{End}(E)$. Now taking Dolbeault cohomology classes via $\mathcal{A}_{\mathbb{C}}^{k,k}(\text{End}(E)) \rightarrow H^k(X, \Omega^k \otimes \text{End}(E))$,

$$\text{ch}_k(E) = \frac{1}{k!} \left(\frac{i}{2\pi} \right)^k \text{tr}([F_{\nabla}]^{\otimes k})$$

where $[F_{\nabla}]^{\otimes k}$ is the image under the map,

$$H^1(X, \Omega_X^1 \otimes \text{End}(E)) \times \dots \times H^1(X, \Omega_X^1 \otimes \text{End}(E)) \rightarrow H^k(X, \Omega^k \otimes \text{End}(E))$$

Furthermore $[F_{\nabla}] = A(E)$ so we get,

$$\text{ch}_k(E) = \frac{1}{k!} \left(\frac{i}{2\pi} \right)^k \text{tr}(A(E)^{\otimes k})$$

as a class under the map $H^k(X, \Omega^k \otimes \text{End}(E)) \xrightarrow{\text{tr}} H^k(X, \Omega_X^k) \subset H^{2k}(X, \mathbb{C})$.

4.4.12 4.4.12

Let X be compact Kähler and E a holomorphic vector bundle admitting a holomorphic connection. Then $A(E) = 0$ and therefore $c_k(E) = 0$.

5 Chapter 5

5.1 Section 5.1

5.1.1 5.1.1 DO!!

5.2 Section 5.2

5.2.1 5.2.1 DO!!

5.3 Section 5.3

5.3.1 5.3.1 DO!!

6 Chapter 6

6.1 Section 6.1

6.1.1 6.1.1

- (a) Let $X = \mathbb{P}^n$ then by the Euler sequence,

$$0 \longrightarrow \mathcal{O}_X \longrightarrow \mathcal{O}_X(1)^{\oplus(n+1)} \longrightarrow \mathcal{T}_X \longrightarrow 0$$

giving a long exact sequence,

$$H^1(X, \mathcal{O}_X) \longrightarrow H^1(X, \mathcal{O}_X(1))^{\oplus(n+1)} \longrightarrow H^1(X, \mathcal{T}_X) \longrightarrow H^2(X, \mathcal{O}_X)$$

However, $H^i(X, \mathcal{O}_X(k)) = 0$ for $i > 0$ when $k \geq 0$ and thus we find $H^1(X, \mathcal{T}_X) = 0$.

- (b) Let $X = \mathbb{C}^n/\Gamma$ be a complex torus. Then we know $\mathcal{T}_X = \mathcal{O}_X^{\oplus n}$ and we need to compute $H^1(X, \mathcal{T}_X) = H^1(X, \mathcal{O}_X)^{\oplus n}$. Then $h^{0,1} = H^1(X, \mathcal{O}_X)$ and $b_1 = h^{0,1} + h^{1,0} = 2h^{0,1}$ by Serre duality. However, $b_1 = 2n$ because $H^1(X, \mathbb{Z}) \cong \text{Hom}_{\mathbb{Z}}(\Gamma, \mathbb{Z})$ and Γ has rank $2n$. Therefore $h^{0,1} = n$ and thus $H^1(X, \mathcal{T}_X) = H^1(X, \mathcal{O}_X)^{\oplus n}$ meaning that $\dim_{\mathbb{C}} H^1(X, \mathcal{T}_X) = n^n$.
- (c) Let X be a compact curve of genus g . Then \mathcal{T}_X is a line bundle of degree $2-2g$. Then, by Serre duality, $H^1(X, \mathcal{T}_X) = H^0(X, \Omega_X^{\otimes 2})$. However if $g = 0$ then Ω_X is negative so $H^1(X, \mathcal{T}_X) = 0$. If $g = 1$ then $\mathcal{T}_X \cong \Omega_X \cong \mathcal{O}_X$ in which case $\dim_{\mathbb{C}} H^1(X, \mathcal{T}_X) = 1$. Finally, if $g > 1$ then Ω_X is positive and $\Omega_X^{\otimes 2}$ has degree $4g - 4 > 2g - 2$ so $H^1(X, \Omega_X^{\otimes 2}) = 0$ and thus, by Riemann-Roch,

$$\dim_{\mathbb{C}} H^1(X, \mathcal{T}_X) = \deg \Omega_X^{\otimes 2} + 1 - g = 3g - 3$$

6.1.2 6.1.2 DO!!

Let X be a compact complex manifold (not necessarily Kähler) and $\sigma \in H^0(X, \Omega_X^2)$ an everywhere non-degenerate holomorphic two-form meaning the map $\sigma : \mathcal{T}_X \rightarrow \Omega_X$ is an isomorphism. Since σ is nondegenerate fiberwise, we see that $\dim X = 2r$ is even. Furthermore,

$$\sigma^r = \sigma \wedge \cdots \wedge \sigma \in H^0(X, \Omega_X^{2r})$$

is everywhere nonvanishing and thus $\sigma^r \in H^0(X, K_X)$ trivializes the canonical bundle K_X so X is a Calabi-Yau. Therefore, applying the results of this section with $\Omega = \sigma^r$, if $v \in H^1(X, \mathcal{T}_X)$ is a cohomology class, then there exists a $\bar{\partial}$ -closed lift $\phi_1 \in \mathcal{A}^{0,1}(\mathcal{T}_X)$ representing $H^1(X, \mathcal{T}_X)$ such that the Maurer-Cartan equation admits a formal solution $\sum_i \phi_i t^i$ extending ϕ_1 .

6.1.3 6.1.3

Let X be a compact complex manifold with $H^2(X, \mathcal{T}_X) = 0$. Take a cohomology class $v \in H^1(X, \mathcal{T}_X)$ and lift to some $\bar{\partial}$ -closed $\phi_1 \in \mathcal{A}^{0,1}(\mathcal{T}_X)$ such that $[\phi_1] = v$. Now for induction suppose we have classes $\phi_1, \dots, \phi_n \in \mathcal{A}^{0,1}(\mathcal{T}_X)$ satisfying the Maurer-Cartan equations up to degree n ,

$$\begin{aligned} \bar{\partial}\phi_1 &= 0 \\ \bar{\partial}\phi_2 &= - \sum_{0 < i < 2} [\phi_i, \phi_{2-i}] \\ &\vdots \\ \bar{\partial}\phi_n &= - \sum_{0 < i < n} [\phi_i, \phi_{n-i}] \end{aligned}$$

Now consider,

$$\omega = - \sum_{0 < i < n+1} [\phi_i, \phi_{n+1-i}]$$

and we have,

$$\begin{aligned}
\bar{\partial}\omega &= - \sum_{0 < i < n+1} \bar{\partial}[\phi_i, \phi_{n+1-i}] = - \sum_{0 < i < n+1} ([\bar{\partial}\phi_i, \phi_{n+1-i}] + [\phi_i, \bar{\partial}\phi_{n+1-i}]) \\
&= \sum_{0 < i < n+1} \left(\sum_{0 < j < i} [[\phi_j, \phi_{i-j}], \phi_{n+1-i}] + \sum_{0 < j < n+1-i} [\phi_i, [\phi_j, \phi_{n+1-i-j}]] \right) \\
&= \sum_{0 < i < n+1} \sum_{0 < j < i} [[\phi_j, \phi_{i-j}], \phi_{n+1-i}] + \sum_{0 < i < n+1} \sum_{0 < j < i} [\phi_{n+1-i}, [\phi_j, \phi_i]] = 0
\end{aligned}$$

replacing i by $n+1-i$ in the second sum. Vanishing follows from $[\alpha, \beta] = (-1)^{k\ell+1}[\beta, \alpha]$ for $\alpha \in \mathcal{A}^{0,k}(\mathcal{T}_X)$ and $\beta \in \mathcal{A}^{0,\ell}(\mathcal{T}_X)$ and in our case $k=2$. Therefore $\omega \in \mathcal{A}^{0,2}(\mathcal{T}_X)$ is $\bar{\partial}$ -closed but $H^2(X, \mathcal{T}_X) = 0$ so every $\bar{\partial}$ -closed \mathcal{T}_X -valued 2-form is $\bar{\partial}$ -exact meaning there exists $\phi_{n+1} \in \mathcal{A}^{0,1}(\mathcal{T}_X)$ such that $\omega = \bar{\partial}\phi_{n+1}$. Explicitly,

$$\bar{\partial}\phi_{n+1} = - \sum_{0 < i < n+1} [\phi_i, \phi_{n+1-i}]$$

solving the Maurer-Cartan equation recursively.

7 Extra Questions for Ron

7.0.1 1

Kodaira embedding says that every positive line bundle is ample in the sense of having some power very ample. Does the algebraic geometry definition work here? I.e. L is ample iff for each bundle Q we have $Q \otimes L^n$ generated by global sections for $n \gg 0$. Do we need Q to be arbitrary coherent sheaf.

Yes, in fact we only need this for vector bundles because it then follows by resolution for all coherent sheaves.

7.0.2 2

If we have a big line bundle $H^0(X, L^{\otimes m}) \sim m^n$ then does it follow there is an ample line bundle i.e. X is projective. I am guessing not. This is similar to asking if there are non algebraic examples of compact Moishezon manifolds $a(X) = \dim X$.