

1 CM Types

2 Algebraic de Rham Cohomology

Let X be a smooth quasi-projective variety over K a characteristic zero field. Let $\dim X = n$. The algebraic de Rham complex is,

$$\Omega_X^\bullet := [0 \rightarrow \mathcal{O}_X \rightarrow \Omega_X^1 \rightarrow \Omega_X^2 \rightarrow \cdots]$$

where $\Omega_X^p := \wedge^p \Omega_{X/K}$ is the sheaf of Kahler differentials and differential is defined as,

$$d(f_0 df_1 \wedge \cdots \wedge df_r) = df_0 \wedge df_1 \wedge \cdots \wedge df_r$$

Then we define the algebraic de Rham cohomology,

$$H_{\text{dR}}^i(X/K) = \mathbb{H}^i(X, \Omega_X^\bullet)$$

There is a filtration on Ω_X^\bullet denoted,

$$\Omega_X^{\bullet \geq p} := [0 \rightarrow \cdots \rightarrow 0 \rightarrow \Omega_X^p \rightarrow \Omega_X^{p+1} \rightarrow \cdots]$$

The graded parts are,

$$\Omega_X^{\bullet \geq p} / \Omega_X^{\bullet \geq p+1} = \Omega_X^p[-p]$$

This gives a filtration on $H_{\text{dR}}^i(X/K)$ by,

$$F^p H_{\text{dR}}^i(X/K) := \text{im}(\mathbb{H}(X, \Omega_X^{\bullet \geq p}) \rightarrow \mathbb{H}(X, \Omega_X^\bullet))$$

Then we get the Hodge-de Rham spectral sequence,

$$E_1^{p,q} = H^q(X, \Omega_X^p) \implies \mathbb{H}^{p+q}(X, \Omega_X^\bullet)$$

If X/\mathbb{C} has structure of a complex manifold then we can consider the analogous construction in the analytic topology:

$$\Omega_{X^{\text{an}}}^\bullet := [0 \rightarrow \mathcal{O}_{X^{\text{an}}} \rightarrow \Omega_{X^{\text{an}}}^1 \rightarrow \Omega_{X^{\text{an}}}^2 \rightarrow \cdots]$$

Then GAGA implies (in the proper case),

$$H_{\text{dR}}^i(X^{\text{an}}/\mathbb{C}) := \mathbb{H}^i(X, \Omega_X^{\text{an}}) = H_{\text{dR}}^i(X/\mathbb{C})$$

Note that we need to use the compatibility of the algebraic and analytic Hodge-de Rham spectral sequences since the de Rham complex is not a complex of \mathcal{O}_X -modules (differentials are nonlinear) but we can compare the Hodge cohomology directly from GAGA). This is true in more generality (FIND CITATION)

Proposition 2.0.1 (Grothendieck). Let X be smooth over \mathbb{C} . Then,

$$H_{\text{dR}}^i(X/\mathbb{C}) \cong H_{\text{sing}}^i(X^{\text{an}}, \mathbb{C})$$

Proof. In the analytic topology there is a quasi-isomorphism from the Poincare lemma,

$$\mathbb{C}[0] \rightarrow \Omega_{X^{\text{an}}}^\bullet$$

And therefore,

$$H^i(X^{\text{an}}, \mathbb{C}) \cong H_{\text{dR}}^i(X^{\text{an}}/\mathbb{C})$$

since this is an isomorphism on hypercohomology. □

Moreover, if X is a compact Kahler manifold then degeneration of Hodge-de Rham spectral sequence comes down to the Hodge decomposition in terms of harmonic forms. Therefore,

$$H^n(X, \mathbb{C}) = H_{\text{dR}}^n(X/\mathbb{C}) \cong \bigoplus_{p+q=n} H^q(X, \Omega_{X^{\text{an}}}^p)$$

Then H_{dR}^n is an example of a \mathbb{Q} -Hodge structure of pure weight n .

Definition 2.0.2. Let $Q \subset \mathbb{R}$ be a subring. A \mathbb{Q} -Hodge structure of pure weight k is a projective Q -module V with bigrading of $V_{\mathbb{C}} = V \otimes_Q \mathbb{C}$ meaning,

$$V_{\mathbb{C}} = \bigoplus_{p+q=k} H^{p,q}$$

such that,

$$\overline{H}^{p,q} = H^{q,p}$$

Let X/\mathbb{C} and consider $f : X \rightarrow Y$ a family of varieties over \mathbb{C} with f smooth proper. Then there are cohomology sheaves,

$$\mathcal{H}_{\text{dR}}^q(X/Y) := R^q f_* \Omega_{X/Y}^\bullet$$

finite locally free over Y which is filtered by submodules,

$$H^p \mathcal{H}^q = R^q f_* \Omega_{X/Y}^{\bullet \geq p}$$

These are called Hodge bundles and define a variation of Hodge structures. Spectral sequences applied to $\Omega_{X/\mathbb{C}}^\bullet$ has,

$$E_1^{p,q} = \Omega_Y^p \otimes_{\mathcal{O}_Y} \mathcal{H}_{\text{dR}}^q(X/Y) \implies H_{\text{dR}}^{p+q}(X/\mathbb{C})$$

2.1 Gauss-Manin Connection

$\Omega_{Y/\mathbb{C}}^p \otimes_{\mathcal{O}_Y} \mathcal{H}_{\text{dR}}^q \implies R^{p+q} f_* \Omega_X^\bullet$ differentials on E_1 -page give,

$$d_1^{p,q} : \Omega_Y^p \otimes_{\mathcal{O}_Y} \mathcal{H}_{\text{dR}}^q \rightarrow \Omega_Y^{p+1} \otimes_{\mathcal{O}_Y} \mathcal{H}_{\text{dR}}^q$$

For $p = 0$ this is the Gauss-Manin connection ∇ . This is an algebraic definition so if $f : X \rightarrow Y$ is defined over $K \subset \mathbb{C}$ then ∇ is defined over K and $\mathcal{H}_{\text{dR}}^q$ is defined over K .

Cycle class map on de Rham or Betti cohomology,

$$H^i(X^{\text{an}}, \mathbb{Q}) \otimes_{\mathbb{Q}} \mathbb{C} \xrightarrow{\sim} H_{\text{dR}}^i(X/K) \otimes_K \mathbb{C}$$

rational structures on both sides not the same (even if $K = \mathbb{Q}$) there are at least factors of $2\pi i$. Then we define cycle class maps, for a line bundle \mathcal{L} we get,

$$c_1(\mathcal{L}) \in H^2(X, \mathbb{Z}(1))$$

Indeed from the exponential sequence,

$$0 \longrightarrow \mathbb{Z}(1) \longrightarrow \mathcal{O}_X \xrightarrow{\exp} \mathcal{O}_X^\times \longrightarrow 0$$

then we get,

$$H^1(X, \mathcal{O}_X) \longrightarrow \text{Pic}(X) \longrightarrow H^2(X, \mathbb{Z}(1)) \longrightarrow H^2(X, \mathcal{O}_X)$$

from the long exact sequence. Alternatively, we get a map,

$$\text{dlog} : \mathcal{O}_X^\times \rightarrow \Omega_{X/K}^1$$

and this gives,

$$H^1(X, \mathcal{O}_X^\times) \rightarrow H^1(X, \Omega_{X/K}^1)$$

However we want a class in de Rham cohomology. To do this, notice that dlog lands inside closed forms so we get a map of complexes,

$$\text{dlog} : \mathcal{O}_X^\times[-1] \rightarrow \Omega_X^\bullet$$

and taking $\mathbb{H}^2(X, -)$ gives,

$$c_1 : \text{Pic}(X) = H^1(X, \mathcal{O}_X^\times) \rightarrow H_{\text{dR}}^2(X/K)$$

In general for a vector bundle \mathcal{E} we want to define,

$$c_p(\mathcal{E}) \in F^p H_{\text{dR}}^{2p}(X/K)$$

and compatibly get a class,

$$c_p(\mathcal{E}) \in H^{2p}(X^{\text{an}}, \mathbb{Z}(p))$$

We define these inductively from c_1 using,

$$H^{2p}(\mathbb{P}(\mathcal{E}), \mathbb{Z}(p)) = \bigoplus_{i=0}^{r-1} \xi^i \cdot \pi^* H^{2p-2i}(X^{\text{an}}, \mathbb{Z}(p-i))$$

where $\xi := c_1(\mathcal{O}(-1))$ and then,

$$\xi^r + \pi^*(c_1)\xi^{r-1} + \cdots + \pi^*(c_p) = 0$$

where we define $c_r(\mathcal{E}) := c_r$. This definition extends from vector bundle to Grothendieck group of coherent sheaves,

$$K_0(X) = G_0(X)$$

since X is smooth. Then for some subvariety $Z \subset X$ we get $[\mathcal{O}_Z] \in G_0(X) = K_0(X)$. Then we define,

$$[Z] = \frac{(-1)^{p-1}}{(p-1)!} c_p([\mathcal{O}_Z]) \in H^{2p}(X^{\text{an}}, \mathbb{Z}(p))$$

where p is the codimension of Z .