

# 1 TODO!!

- (a) Finish symplectic geometry course
  - (a) figure out if symplectic toric is the same as projective toric variety (projectivity needed to come from a polytope and also to be Kahler)
  - (b) review coisotropic reduced and write some notes
  - (c) hyperkahler reduction examples
  - (d) are there examples of noncompact hyperkahlers?
  - (e) work out the kinks in notes on hamiltonian actions
- (b) review killing homotopy groups columbia lectures and write some notes
- (c) figure out those damn jet bundles and connections on principal bundles
  - (a) RMK:  $\pi^*E$  is NOT trivial for a vector bundle let alone a fiber bundle. it does get equipped with a canonical section but for a vector bundle this is just the trivial section, only for a principal bundle does giving a section trivialize it.
  - (b) role of atiyah sequence vs jet bundle sequence
  - (c)
- (d) spectral sequences for tor and ext in derived category (FIND MY NOTES ON THIS!)
  - (a) application to universal coefficient theorem
  - (b) Kunneth spectral sequence
  - (c) Kunneth formula for smash product?
  - (d) why are derived functors triangulated
  - (e) derived functors in terms of Kan extensions (NOTES)
- (e) write notes on universal morphisms
- (f)  $G$ -action of  $X/Y$  induces map Descent data  $X/Y$  to  $G$ -equivariant sheaves
  - (a) isomorphism when  $X/Y$  is a  $G$ -cover i.e.  $X \rightarrow Y$  is a  $G$ -torsor
  - (b) write down explicit  $G$ -equivariant structure on  $\Omega_X$
  - (c) Galois descent derive explicit form
- (g) Weil restriction
  - (a) write down trivialization after going back up
  - (b) Galois descent in explicit form
- (h) notes on Galois actions on schemes
- (i) notes on Frobenii
- (j) notes on universal constructions in math with examples

- (k) fix notes on Tor in category of sheaves and Tor symmetry (do I need symmetry of flat objects a priori?).
- (l) Finish stable homotopy theory course.
- (m) Finish vector bundles and connections notes (in AG folder)
  - (a) Kahler iff  $\nabla I = 0$  where  $\nabla$  is the Levi-Civita connection
  - (b) Ricci tensor and the trace bullshit
  - (c) Riemann-Hilbert and existence of flat frames for integrable connections

## 2 What I Want to Think About

- (a) Flat cohomology equal etale cohomology for smooth (affine groups) apply this to that counting rational points things
- (b) work out the details for the group fixing  $\mathbb{C}$  inside endomorphism group. What does an integrable structure of this kind look like, how close to a complex manifold can we get? In dimension two this should be exactly a conformal (not necessarily orientable) structure.
- (c) FINISH CONFORMAL NOTES!
- (d) Hilbert Class Field of curves (ASK BRIAN FOR REFERENCE)
- (e) Read about Fredholm index and Riemann-Roch
- (f) Cohomology and inclusion-exclusion: cohomology for vectorspaces?

## 3 Some Questions I Have

- (a) Reduction of structure group for a scheme.
  - (a) what about the algebraic group  $SL^\pm = \det^{-1}(\mu)$  what does reduction of structure group give. For a manifold this is supposed to be a pseudo-volume form but obviously that's not right.
  - (b) what about  $\text{Res}_{\mathbb{R}}^{\mathbb{C}}(\mathbb{G}_m) \hookrightarrow GL_2$  from the action  $\mathbb{G}_m \curvearrowright \mathbb{A}_{\mathbb{C}}^1$  restricted giving an action  $\text{Res}_{\mathbb{R}}^{\mathbb{C}}(\mathbb{G}_m) \curvearrowright \mathbb{A}_{\mathbb{R}}^2$ . I feel like this should give an almost complex structure. What properties does it have? What about for other fields?
  - (c) What is an almost complex structure on a scheme look like?
- (b) Is my calculation of an “almost almost complex structure” as reduction of structure group to  $\langle \sigma \rangle \rtimes GL(n, \mathbb{C}) \subset GL(2n, \mathbb{R})$ . For the case  $n = 1$  this should be the conformal group justifying that I think this should correspond to the non-oriented case of a complex manifold since Riemann surfaces are exactly oriented conformal manifolds.

## 4 The Tautological Bundle

Consider the fibre bundle,  $\pi : S^{2n+1} \rightarrow \mathbb{P}_{\mathbb{C}}^n$  given by considering  $S^{2n+1} \subset \mathbb{C}^{n+1}$  and restricting the projection  $\mathbb{C}^{n+1} \rightarrow \mathbb{P}_{\mathbb{C}}^n$ . Then  $\pi$  is a principal  $S^1$ -bundle. Consider the tautological representation  $\rho : U(1) \rightarrow \text{GL}_1(\mathbb{C})$  which is the inclusion  $U(1) \hookrightarrow \mathbb{C}^\times$ , which gives an associated line bundle  $S^{2n+1} \times_{\rho} \mathbb{C}$ . We call this the tautological bundle since its fibre above a point is the line in  $\mathbb{C}^{n+1}$  which that point on  $\mathbb{P}_{\mathbb{C}}^n$  corresponds to.

To see this explicitly, consider the following bundle,

$$T = \{(L, v) \mid L \in \mathbb{P}_{\mathbb{C}}^n \text{ and } v \in L \subset \mathbb{C}^{n+1}\} \subset \mathbb{P}_{\mathbb{C}}^n \times \mathbb{C}^{n+1}$$

with the projection  $\pi : T \rightarrow \mathbb{P}_{\mathbb{C}}^n$  via  $(L, v) \mapsto L$ . I claim that this bundle is isomorphic to the tautological bundle constructed above.

Consider the map  $f : S^{2n+1} \times_{\rho} \mathbb{C} \rightarrow T$  via  $f : [x, \lambda] \mapsto (\text{Span}(x), \lambda x)$ . This is clearly a bundle map since  $\pi([x, \lambda]) = \pi(x) = \text{Span}((\cdot)x) = \pi(\text{Span}(x), \lambda x)$ . Furthermore it is well-defined because  $f([x, \mu\lambda]) = (\text{Span}(x), \mu\lambda x) = (\text{Span}(\mu x), \lambda\mu x) = f([\mu x, \lambda])$ . We need to check that this map is injective and surjective. First, if  $f([x, \lambda]) = f([y, \mu])$  then  $\text{Span}(x) = \text{Span}(y)$  so  $y = \gamma x$  for  $\gamma \in \mathbb{C}^\times$  and  $\lambda x = \mu y$  so  $\lambda = \mu\gamma$  (since these vectors are nonzero) and thus,

$$[x, \lambda] = [x, \gamma\mu] = [\gamma x, \mu] = [y, \mu]$$

For surjectivity note that given  $(L, v)$  with  $v \in L$  then  $L = \text{Span}(x)$  for  $x \in S^{2n+1}$  and  $v = \lambda x$  with  $\lambda \in \mathbb{C}$  since  $L$  is a line. Thus  $f([x, \lambda]) = (L, v)$ .

The tautological bundle has no nonzero (holomorphic) global sections. However, there are  $n + 1$  independent global sections of its dual. To see this consider the global  $\text{Hom}(T, \mathcal{O}_{\mathbb{P}})$ . There exist  $n + 1$  independent functions defined by the  $n + 1$  projections  $p_k : \mathbb{C}^{n+1} \rightarrow \mathbb{C}$  via the construction,

$$T \hookrightarrow \mathcal{O}_{\mathbb{P}}^{n+1} = \mathbb{P}_{\mathbb{C}}^n \times \mathbb{C}^{n+1} \xrightarrow{p_k} \mathbb{P}_{\mathbb{C}}^n \times \mathbb{C} = \mathcal{O}_{\mathbb{P}}$$

These sections are referred to as  $X_k$ , the  $k^{\text{th}}$  coordinate function on  $\mathbb{P}_{\mathbb{C}}^n$ .

Producing the coordinate functions  $X_k$  as sections of the dual  $X^\vee$  identifies the tautological bundle  $T$  with the algebraic twist  $\mathcal{O}_{\mathbb{P}}(-1)$  and thus its dual is the Serre twisting sheaf  $T^\vee = \mathcal{O}_{\mathbb{P}}(1)$ .

## 5 MATH 275A 2021 Lecture 2

Using the Stern-Gerlach boxes we define spin operators  $\hat{S}_i$  on our Hilbert space  $H = \mathcal{C}^2$ . These have eigenstate  $\ker \pm$  along each axis. Furthermore, we have a Hamiltonian  $\hat{H}$ . For a constant magnetic field, up to a constant,

$$\hat{H} = \hat{S} \cdot \vec{B}$$

For  $B$  along the  $z$ -direction,

$$\hat{H} = \hat{S}_z B$$

Then the evolution follows the Schrodinger equation,

$$i\partial_t |\psi\rangle = \hat{H} |\psi\rangle$$

For any observable (i.e. operator  $\hat{A}$ ) we can define the expected value,

$$\langle \hat{A} \rangle_\psi = \langle \psi | \hat{A} | \psi \rangle$$

Then,

$$i\partial_t \langle \hat{A} \rangle_\psi = \langle [\hat{A}, \hat{H}] \rangle_\psi$$

Now for example, we choose  $|\psi(0)\rangle = |+_x\rangle$ . Then we expand,

$$|\psi(0)\rangle = \frac{1}{\sqrt{2}} (|+_z\rangle + |-_z\rangle)$$

Then applying the evolution operator,

$$|\psi(t)\rangle = e^{-iHt} \frac{1}{\sqrt{2}} (|+_z\rangle + |-_z\rangle) = \frac{1}{\sqrt{2}} (e^{-i\frac{B}{2}t} |+_z\rangle + e^{i\frac{B}{2}t} |-_z\rangle)$$

Now we consider,

$$i\partial_t \langle \hat{S}_x \rangle = \langle \psi | \hat{S}_x | \psi \rangle = \langle [\hat{S}_x, \hat{H}] \rangle = B \langle [\hat{S}_x, \hat{S}_z] \rangle = -iB\hat{S}_y$$

and therefore,

$$\partial_t \langle \hat{S}_x \rangle = -B \langle \hat{S}_y \rangle$$

Likewise,

$$\partial_t \langle \hat{S}_y \rangle = B \langle \hat{S}_x \rangle$$

This coupled system has solution,

$$\langle \hat{S}_x \rangle = \cos(Bt) \quad \text{and} \quad \langle \hat{S}_y \rangle = \sin(Bt)$$

### 5.0.1 Operators

Infinite dimensional space  $H = L^2(\mathbb{R}) = \{f : \mathbb{R} \rightarrow \mathbb{C} \mid \int |f|^2 < \infty\}$ . We take observables to be “self-adjoint” operators on  $H = L^2(\mathbb{R})$ . For example,  $\hat{x} = x \cdot$  and  $\hat{p} = -\partial_x$ . However, the eigenfunctions of these operators are not  $L^2$  they are tempered distributions. We say,

$$\left\langle \frac{1}{\sqrt{2\pi}} e^{ipx} \middle| \frac{1}{\sqrt{2\pi}} e^{iqx} \right\rangle = \nabla(p - q)$$

### 5.0.2 Uncertainty Principle

Define,

$$\nabla \hat{A} = \hat{A} = -\langle \hat{x} \rangle I$$

and likewise for  $B$  two self-adjoint operators  $A, B$ . Then,

$$\langle (\nabla \hat{x})^2 \rangle_\psi \langle (\nabla \hat{p})^2 \rangle_\psi \geq \frac{1}{4} |\langle \psi | [\hat{A}, \hat{B}] | \psi \rangle|^2$$

For example,

$$[\hat{x}, \hat{p}] = \hat{x}\hat{p} - \hat{p}\hat{x} = iI$$

because,

$$(\hat{x}\hat{p} - \hat{p}\hat{x})\psi = x(-i\partial_x\psi) + i\partial_x(x\psi) = -i\partial_x\psi + i\psi + x\partial_x\psi = i\psi$$

Therefore,

$$\sigma_x^2 \sigma_p^2 \geq \frac{1}{4}$$

### 5.0.3 Angular Momentum

Classical angular momentum  $\vec{L} = \vec{x} \times \vec{p}$ . We upgrade these to quantum self-adjoint operators. Thus we get, for example,

$$\hat{L}_z = -i(x\partial_y - y\partial_x)$$

Then  $L^2 = L_x^2 + L_y^2 + L_z^2$ .

## 6 The Tautological Bundle

Consider the fibre bundle,  $\pi : S^{2n+1} \rightarrow \mathbb{P}_{\mathbb{C}}^n$  given by considering  $S^{2n+1} \subset \mathbb{C}^{n+1}$  and restricting the projection  $\mathbb{C}^{n+1} \rightarrow \mathbb{P}_{\mathbb{C}}^n$ . Then  $\pi$  is a principal  $S^1$ -bundle. Consider the tautological representation  $\rho : U(1) \rightarrow \text{GL}_1(\mathbb{C})$  which is the inclusion  $U(1) \hookrightarrow \mathbb{C}^\times$ , which gives an associated line bundle  $S^{2n+1} \times_{\rho} \mathbb{C}$ . We call this the tautological bundle since its fibre above a point is the line in  $\mathbb{C}^{n+1}$  which that point on  $\mathbb{P}_{\mathbb{C}}^n$  corresponds to.

To see this explicitly, consider the following bundle,

$$T = \{(L, v) \mid L \in \mathbb{P}_{\mathbb{C}}^n \text{ and } v \in L \subset \mathbb{C}^{n+1}\} \subset \mathbb{P}_{\mathbb{C}}^n \times \mathbb{C}^{n+1}$$

with the projection  $\pi : T \rightarrow \mathbb{P}_{\mathbb{C}}^n$  via  $(L, v) \mapsto L$ . I claim that this bundle is isomorphic to the tautological bundle constructed above.

Consider the map  $f : S^{2n+1} \times_{\rho} \mathbb{C} \rightarrow T$  via  $f : [x, \lambda] \mapsto (\text{Span}(x), \lambda x)$ . This is clearly a bundle map since  $\pi([x, \lambda]) = \pi(x) = \text{Span}((x)) = \pi(\text{Span}(x), \lambda x)$ . Furthermore it is well-defined because  $f([x, \mu\lambda]) = (\text{Span}(x), \mu\lambda x) = (\text{Span}(\mu x), \lambda\mu x) = f([\mu x, \lambda])$ . We need to check that this map is injective and surjective. First, if  $f([x, \lambda]) = f([y, \mu])$  then  $\text{Span}(x) = \text{Span}(y)$  so  $y = \gamma x$  for  $\gamma \in \mathbb{C}^\times$  and  $\lambda x = \mu y$  so  $\lambda = \mu\gamma$  (since these vectors are nonzero) and thus,

$$[x, \lambda] = [x, \gamma\mu] = [\gamma x, \mu] = [y, \mu]$$

For surjectivity note that given  $(L, v)$  with  $v \in L$  then  $L = \text{Span}(x)$  for  $x \in S^{2n+1}$  and  $v = \lambda x$  with  $\lambda \in \mathbb{C}$  since  $L$  is a line. Thus  $f([x, \lambda]) = (L, v)$ .

The tautological bundle has no nonzero (holomorphic) global sections. However, there are  $n + 1$  independent global sections of its dual. To see this consider the global  $\text{Hom}(T, \mathcal{O}_{\mathbb{P}})$ . There exist  $n + 1$  independent functions defined by the  $n + 1$  projections  $p_k : \mathbb{C}^{n+1} \rightarrow \mathbb{C}$  via the construction,

$$T \hookrightarrow \mathcal{O}_{\mathbb{P}}^{n+1} = \mathbb{P}_{\mathbb{C}}^n \times \mathbb{C}^{n+1} \xrightarrow{p_k} \mathbb{P}_{\mathbb{C}}^n \times \mathbb{C} = \mathcal{O}_{\mathbb{P}}$$

These sections are referred to as  $X_k$ , the  $k^{\text{th}}$  coordinate function on  $\mathbb{P}_{\mathbb{C}}^n$ .

Producing the coordinate functions  $X_k$  as sections of the dual  $X^\vee$  identifies the tautological bundle  $T$  with the algebraic twist  $\mathcal{O}_{\mathbb{P}}(-1)$  and thus its dual is the Serre twisting sheaf  $T^\vee = \mathcal{O}_{\mathbb{P}}(1)$ .

## 7 Some Connection Musings

**Definition 7.0.1.** Let  $f : E \rightarrow X$  be a smooth surjection (in the smooth category, what should it be in the algebraic category?) then an *Erhesmann connection* is a splitting of the sequence of vector bundles,

$$0 \longrightarrow \ker df \longrightarrow TE \longrightarrow \pi^*TX \longrightarrow 0$$

where we usually call  $V = \ker df$  the vertical bundle. In algebraic language,  $V$  is the dual of the relative differentials so the connection corresponds to a splitting of,

$$0 \longrightarrow f^*\Omega_X \longrightarrow \Omega_E \longrightarrow \Omega_{E/X} \longrightarrow 0$$

*Remark.* Such splittings are supposed to correspond to smooth sections of the map  $J^1(E) \rightarrow E$ . We now explain how this works. Unfortunately, I don't know a good unified language to describe the jet bundles so I will give the algebraic and smooth definitions.

**Definition 7.0.2.** Given a smooth surjection  $f : E \rightarrow X$ , consider the  $n$ -th thickened diagonal,  $X \rightarrow X_n \rightarrow X \times_S X$ . Then we consider the functor sending  $T \rightarrow S$  to pairs of maps  $T \rightarrow X$  and  $T \times_X X_n \rightarrow E$  such that the diagram,

$$\begin{array}{ccc} E & \xrightarrow{f} & X \\ \uparrow & & \uparrow \pi_1 \\ T \times_X X_n & \longrightarrow & X_n \\ \downarrow & \lrcorner & \downarrow \pi_2 \\ T & \longrightarrow & X \end{array}$$

commutes. Then the jet scheme  $J_n(E/X)$  with maps  $J_n(E/X) \rightarrow X$  and  $J^n(E/X) \times_X X_n \rightarrow E$  represents this functor.

## 8 Counterexamples In Geometry

**Example 8.0.1.** The Hopf surface is the compact complex surface  $H = \mathbb{C}^2 \setminus \{0\}/\mathbb{Z}$  where  $\mathbb{Z} \curvearrowright \mathbb{C}^2$  via  $(z_1, z_2) \mapsto (\lambda z_1, \lambda z_2)$  for  $0 < \lambda < 1$ . This surface has  $h^{1,0} = 1$  but  $h^{0,1} = 0$ . Furthermore,  $H$  is diffeomorphic to  $S^3 \times S^1$ . This provides:

- (a) a compact complex manifold that is not Kähler
- (b) a compact complex manifold without Hodge symmetry
- (c) a compact complex manifold that is not symplectic ( $H^2(H, \mathbb{Z}) = 0$ )

*Remark.* From the exponential exact sequences,

$$0 \longrightarrow \mathbb{Z} \longrightarrow \mathcal{O}_X \xrightarrow{\exp} \mathcal{O}_X^\times \longrightarrow 0$$

we have that,

## 9 Questions I was asked in interviews

### 9.1 Oxford

**Exercise 9.1.1.** Which genus Riemann surfaces have a covering map to a genus 2 Riemann surface.

---

From Riemann-Hurwitz we have  $2g - 2 = 2n$  where  $2h - 2 = 2$  since  $h = 2$ . Thus  $g = n + 1$  so any genus can appear. To show that these are all actually possible, draw a picture with a central donut surrounded by  $g - 1$  donuts. This maps by cyclic quotienting onto a two holed torus.

### 9.2 LSGNT

**Exercise 9.2.1.** Let  $E$  be an elliptic curve over  $\mathbb{F}_p$ . Given  $a_p$  how do you find  $\#E(\mathbb{F}_{p^k})$ ?

---

The zeta function is,

$$\zeta_E(t) = \frac{t^2 - a_p t + p}{(1 - t)(1 - pt)}$$

and therefore,

$$\#E(\mathbb{F}_{p^k}) = 1 + p^k - \alpha^k - \beta^k$$

where  $\alpha$  and  $\beta$  are the roots of  $t^2 - a_p t + p$  which are determined via  $\alpha + \beta = a_p$  and  $\alpha\beta = p$ .

## 10 Questions

- (a) When people write  $\mathcal{M}_\ell(\mathbb{C}) = \mathfrak{h}/\mathrm{SL}2\mathbb{Z}$  isn't this wrong because every point of  $\mathcal{M}_\ell(\mathbb{C})$  is "stacky" i.e. this groupoid has  $\mathbb{Z}/2\mathbb{Z}$  stabilizer at the general point due to the inversion map. However,  $\mathfrak{h}/\mathrm{SL}2\mathbb{Z}$  seems to be a setoid at the general point, oh no that's wrong because  $-I$  stabilizes each point and stabilizes a lattice but acts on the elliptic curve by inversion. AHH!!! This is why we retain the  $\mathrm{SL}2\mathbb{Z}$  and don't pass to  $\mathrm{PSL}2\mathbb{Z}$ .

## 11 The Universal Elliptic Curve and Modular Forms

Let  $\mathfrak{h}$  be the upper half plane with coordinate  $\tau$  which we think of as parametrizing the complex elliptic curve  $\mathbb{C}/(\mathbb{Z} \oplus \tau\mathbb{Z})$ . We first make a universal family of elliptic curves by,

$$\pi : \mathbb{C} \times \mathfrak{h} / \langle (z, \tau) \sim (z + 1, \tau) \sim (z + \tau, \tau) \rangle \rightarrow \mathfrak{h}$$

This map has fiber over  $\tau$  equal to  $\mathbb{C}/(\mathbb{Z} \oplus \tau\mathbb{Z})$ . However, if  $\gamma \in \mathrm{SL}2\mathbb{Z}$  takes  $\gamma \cdot \tau = \tau'$  then I claim that the elliptic curves are isomorphic. We want to encode this. The isomorphism comes from the transformation of a positive ordered basis  $\omega_1, \omega_2$  of a lattice into the form  $\tau, 1$  where  $\tau = \frac{\omega_1}{\omega_2}$ . Then  $\gamma \cdot \tau = \frac{a\tau + b}{c\tau + d}$  is the image of the basis  $\tau, 1$  under  $\gamma$  after dividing by  $(c\tau + d)$  so that we are normalized to  $\gamma \cdot \tau, 1$ . Therefore, the isomorphism  $C_\tau \xrightarrow{\sim} C_{\gamma\tau}$  is given by  $z \mapsto (c\tau + d)^{-1}z$ . Therefore, we mod out our family to get,

$$\mathcal{C} \rightarrow \mathcal{M}$$

where  $\mathcal{M} = \mathfrak{h}/\mathrm{SL}2\mathbb{Z}$  and,

$$\begin{aligned} \mathcal{C} &= \mathbb{C} \times \mathfrak{h} / \langle (z, \tau) \sim (z + 1, \tau) \sim (z + \tau, \tau) \rangle / \mathrm{SL}2\mathbb{Z} \\ &= \mathbb{C} \times \mathfrak{h} / \langle (z, \tau) \sim (z + 1, \tau) \sim (z + \tau, \tau) \sim ((c\tau + d)^{-1}z, \gamma \cdot \tau) \rangle \end{aligned}$$

Now we consider the vertical cotangent bundle of  $\mathcal{C} \rightarrow \mathcal{M}$  which is,

$$\Omega_{\mathcal{C}/\mathcal{M}} = \mathbb{C} \times \mathbb{C} \times \mathfrak{h} / \langle (\omega, z, \tau) \sim (\omega, z+1, \tau) \sim (\omega, z+\tau, \tau) \sim ((c\tau+d)\omega, (c\tau+d)^{-1}z, \gamma \cdot \tau) \rangle$$

Notice that the cotangent fibers change opposite to the coordinate  $z$  because the natural forward map is the inverse pullback which scales oppositely. All this is extremely problematic because of the fixed points of  $\mathrm{SL}_2\mathbb{Z} \curvearrowright \mathfrak{h}$  which give stabilizers and thus too much quotienting we really should be taking groupoid quotients and thus get “stacky” points but the functions in the two cases are basically the same because they reduce to being equivariant maps. Now pulling back this bundle along the zero section of  $\mathcal{C} \rightarrow \mathcal{M}$  (i.e. the map  $\tau \mapsto (0, \tau)$ ) gives,

$$\omega = e^*\Omega_{\mathcal{C}/\mathcal{M}} = \mathbb{C} \times \mathfrak{h} / (\omega, \tau) \sim ((c\tau+d)\omega, \gamma \cdot \tau)$$

Therefore, a section of  $\omega \rightarrow \mathcal{M}$  is an equivariant section of  $\mathbb{C} \times \mathfrak{h} \rightarrow \mathfrak{h}$  and thus a function  $f : \mathfrak{h} \rightarrow \mathbb{C}$  such that,

$$f(\gamma \cdot \tau) = (c\tau + d)f(\tau)$$

which is exactly a weight-one modular function!. Therefore, modular functions of weight  $k$  correspond to sections of  $\omega^{\otimes k}$ .

To get modular forms, we need a holomorphy condition at  $\infty$ . To get this, we need to extend  $\omega$  over the boundary to a line bundle on  $\overline{\mathcal{M}}$ .

## 12 How many nodes does a curve in $\mathbb{P}^3$ have when projected to $\mathbb{P}^2$

Given a smooth curve  $X \subset \mathbb{P}^3$  of genus  $g$  we consider a general projection  $\mathbb{P}^3 \dashrightarrow \mathbb{P}^2$ . Then  $X$  gets a node at some point if the associated line intersects  $X$  at multiple points.

For a curve  $X \subset \mathbb{P}^2$  consider the projection map  $X \rightarrow \mathbb{P}^1$ . This is ramified exactly at the points where a section  $s \in \Gamma(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(1))$  vanishes as well as its derivative on  $X$ . Thus we want to consider  $J^1(\mathcal{O}_X(1))$  and the vanishing locus of a general section. But we don’t want a general line, we want a general line that vanishes at a fixed point  $p \in \mathbb{P}^2$ . Call this linear system  $V_p \subset \Gamma(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(1))$ . Then

## 13 Viewpoints on Čech Cohomology

### 13.1 Čech Cohomology as the Cohomology of a Complex

Let  $(\mathcal{C}, \mathcal{O})$  be a ringed site,  $U \in \mathcal{C}$  and  $\mathfrak{U} = \{U_i \rightarrow U\}_{i \in I}$  be a cover of  $U$ . We set,

$$U_{i_0, \dots, i_p} = U_{i_0} \times_U \cdots \times_U U_{i_p}$$

**Definition 13.1.1.** The Čech complex of a presheaf  $\mathcal{F}$  on  $\mathcal{C}$  is defined by terms,

$$\check{C}^p(\mathfrak{U}, \mathcal{F}) = \prod_{i_0, \dots, i_p \in I} \mathcal{F}(U_{i_0, \dots, i_p})$$

giving a complex,



$$0 \longrightarrow \prod_{i_0 \in I} \mathcal{F}(U_{i_0}) \xrightarrow{d} \prod_{i_0, i_1 \in I} \mathcal{F}(U_{i_0} \times_U U_{i_1}) \xrightarrow{d} \dots$$

where,

$$d(s)_{i_0, \dots, i_{p+1}} = \sum_{j=0}^{p+1} (-1)^j s_{i_0, \dots, \hat{i}_j, \dots, i_{p+1}}|_{U_{i_0, \dots, i_{p+1}}}$$

This is a complex. Then the Čech cohomology is the cohomology of this complex,

$$\check{H}^p(\mathfrak{U}, \mathcal{F}) = H^p(\check{C}^p(\mathfrak{U}, \mathcal{F}))$$

**Proposition 13.1.2.** Formation of the Čech complex is functorial,

$$\check{C}^\bullet(\mathfrak{U}, -) : \mathbf{PSh}_{\mathcal{O}} \rightarrow \mathbf{Ch}(\mathbf{Mod}_{\mathcal{O}(U)})$$

and therefore Čech cohomology is functorial,

$$\check{H}^p(\mathfrak{U}, -) : \mathbf{PSh}_{\mathcal{O}} \rightarrow \mathbf{Mod}_{\mathcal{O}(U)}$$

**Lemma 13.1.3.** If  $\mathcal{I}$  is an injective presheaf then  $\check{H}^p(\mathfrak{U}, \mathcal{I}) = 0$  for all  $p > 0$ .

*Proof.* FIND A GOOD PROOF (e.g. Tag 01EN). □

**Proposition 13.1.4.** The functors  $\check{H}^p(\mathfrak{U}, -) : \mathbf{PSh}_{\mathcal{O}} \rightarrow \mathbf{Mod}_{\mathcal{O}(U)}$  form a universal  $\nabla$ -functor.

*Proof.* Given an exact sequence of presheaves,

$$0 \longrightarrow \mathcal{F}_1 \longrightarrow \mathcal{F}_2 \longrightarrow \mathcal{F}_3 \longrightarrow 0$$

we get an exact sequence of complexes,

$$0 \longrightarrow \check{C}^\bullet(\mathfrak{U}, \mathcal{F}_1) \longrightarrow \check{C}^\bullet(\mathfrak{U}, \mathcal{F}_2) \longrightarrow \check{C}^\bullet(\mathfrak{U}, \mathcal{F}_3) \longrightarrow 0$$

because taking sections of presheaves is exact and products are exact in the category of modules. The associated long exact sequence gives the required connecting maps and exactness showing that  $\check{H}^p(\mathfrak{U}, -)$  form a  $\nabla$ -functor. Furthermore, since  $\mathbf{PSh}_{\mathcal{O}}$  has enough injectives and  $\check{H}^p(\mathfrak{U}, \mathcal{I}) = 0$  for  $p > 0$  we see that  $\check{H}^p(\mathfrak{U}, -)$  are effaceable and thus form a universal  $\nabla$ -functor. □

## 13.2 Čech Cohomology as a Canonical Resolution

## 13.3 Čech Cohomology as a Derived Functor on Presheaves

Let  $(\mathcal{C}, \mathcal{O})$  be a ringed site,  $U \in \mathcal{C}$  and  $\mathfrak{U} = \{U_i \rightarrow U\}_{i \in I}$  be a cover of  $U$  then define the Čech sections functor,

$$\check{H}^0(\mathfrak{U}, -) : \mathbf{PSh}_{\mathcal{O}_X} \rightarrow \mathbf{Mod}_{\mathcal{O}(U)}$$

defined by,

$$\mathcal{F} \mapsto \ker \left( \prod_{i \in I} \mathcal{F}(U_i) \rightarrow \prod_{i, j \in I^2} \mathcal{F}(U_i \times_U U_j) \right)$$

Consider the inclusion functor,

$$\iota : \mathbf{Sh}_{\mathcal{O}} \rightarrow \mathbf{PSh}_{\mathcal{O}_X}$$

Then there is a commutative diagram of functors,

$$\begin{array}{ccc}
\mathbf{PSh}_{\mathcal{O}_X} & \xrightarrow{\quad \iota \quad} & \mathbf{Sh}_{\mathcal{O}_X} \\
& \searrow \check{H}^0 & \swarrow \Gamma_U \\
& \mathbf{Mod}_{\mathcal{O}(U)} & 
\end{array}$$

Furthermore,  $\iota$  is right-adjoint to sheafification which is exact and thus preserves injectives. Therefore, we can apply the Grothendieck spectral sequence to get,

$$E_2^{p,q} = R^p \check{H}^0(\mathfrak{U}, R^q \iota(\mathcal{F})) \implies H^0(U, \mathcal{F})$$

Furthermore, because  $\Gamma_V$  is exact on presheaves, we see that,

$$R^p \Gamma_V = R^p(\Gamma_V \circ \iota) = \Gamma_V \circ R^p \iota$$

and therefore,

$$[(R^p \iota)(\mathcal{F})](V) = \Gamma_V \circ (R^p \iota)(\mathcal{F}) = (R^p \Gamma_V)(\mathcal{F}) = H^p(V, \mathcal{F})$$

Therefore  $(R^p \iota)(\mathcal{F})$  is the presheaf  $V \mapsto H^p(V, \mathcal{F})$  which we call  $\mathcal{H}^p(\mathcal{F})$ . Now I claim that the derived functor  $R^p \check{H}^0$  agrees with  $\check{H}^p$  defined earlier. Since  $\check{H}^0$  is left-exact, there are natural isomorphisms,

$$\check{H}^0 \xrightarrow{\sim} R^0 \check{H}^0$$

Furthermore, because  $\check{H}^p$  and  $R^p \check{H}$  are universal  $\nabla$ -functors, the above map extends to an isomorphism of  $\nabla$ -functors. Therefore, we derived the Čech to derived spectral sequence,

$$E_2^{p,q} = \check{H}^p(\mathfrak{U}, \mathcal{H}^q(\mathcal{F})) \implies H^{p+q}(U, \mathcal{F})$$

Now, we define,

$$\check{H}^0(U, -) = \varinjlim_{\mathfrak{U}} \check{H}^0(\mathfrak{U}, -)$$

Because covers form a filtered poset this filtered colimit is exact. Therefore,

$$\check{H}^p(U, -) := \varinjlim_{\mathfrak{U}} \check{H}^p(\mathfrak{U}, -) = \varinjlim_{\mathfrak{U}} R^p \check{H}^0(\mathfrak{U}, -) = R^p \varinjlim_{\mathfrak{U}} \check{H}^0(\mathfrak{U}, -)$$

Furthermore, since  $\check{H}^0(\mathfrak{U}, -) \circ \iota = \Gamma_U$  for any cover we see that  $\check{H}^0(U, -) \circ \iota = \Gamma_U$ . Therefore, applying the Grothendieck spectral sequence,

$$E_2^{p,q} = \check{H}^p(U, \mathcal{H}^q(\mathcal{F})) \implies H^{p+q}(U, \mathcal{F})$$

### 13.4 The Degree 1 Case

By the locality of cohomology, for each  $s \in H^q(V, \mathcal{F})$  there exists a cover  $\{V_i\}$  of  $V$  such that  $s|_{V_i} = 0$  for each  $i$  and therefore choosing a small enough refinement  $\check{H}^0(U, \mathcal{H}^q(\mathcal{F})) = 0$ . Now consider the differentials  $d_r : E_r^{p,q} \rightarrow E_r^{p+r, q-r+1}$  then for  $(p, q) = (1, 0)$  we see that  $d_r = 0$  for  $r > 1$  because  $(p+r, q-r+1) = (r+1, 1-r)$  is not first quadrant. Furthermore, if  $(p+r, q-r+1) = (1, 0)$  then  $(p, q) = (1-r, r-1)$  is not first quadrant for  $r > 1$  so  $d_r = 0$ . Since  $E_2^{0,1} = 0$  we see that the  $p+q=1$  terms are converged at the  $E_2$ -page. Therefore, from the filtration we see that,

$$\check{H}^1(U, \mathcal{F}) = H^1(U, \mathcal{F})$$

in vast generality.

## 14 Orientability versus Two-Sidedness

In multivariable calculus when you learn about the divergence theorem you may be told “this only works for orientable surfaces because you need to choose a normal vector to integrate over in order to compute the flux” this is only true because the ambient  $\mathbb{R}^3$  is orientable. Furthermore, you are told that the Möbius strip is non-orientable because it only has one side but this is actually a feature of its embedding in  $\mathbb{R}^3$ . In general, orientability, which is intrinsic, is not actually the concept being probed but rather two-sidedness, which is relative to the embedding.

**Definition 14.0.1.** Let  $M$  be a manifold. Then an embedded submanifold  $X \subset M$  is *two-sided* if the normal bundle  $N_M X$  is orientable.

**Proposition 14.0.2.** Suppose that  $X \subset M$  has codimension 1 then  $X$  is two sided if and only if it admits a global nonvanishing normal vector field  $v \in \Gamma(X, N_M X)$ .

*Proof.* The normal bundle  $N_M X$  is a line bundle and thus is orientable if and only if it is trivial if and only if it admits a nonvanishing global section.  $\square$

*Remark.* This motivates the terminology because the nonvanishing normal vector field distinguishes between a “positive” side and a “negative” side of the manifold.

**Proposition 14.0.3.** If  $M$  is orientable and  $X \subset M$  is an embedded submanifold then  $X$  is two-sided if and only if  $M$  is orientable.

*Proof.* From the exact sequence,

$$0 \longrightarrow TX \longrightarrow TM|_X \longrightarrow N_M X \longrightarrow 0$$

we see that,

$$\det TM|_X \cong \det TX \otimes \det N_M X$$

However,  $TM$  is orientable so  $\det TM$  is trivial and thus,

$$\det TM \cong \det N_M X$$

meaning that one bundle is orientable if and only if the other is orientable. Equivalently, we can use Stiefel-Whitney classes,

$$w_1(TX) + w_1(N_M X) = w_1(TM)|_X = 0$$

and therefore (because these live in  $\mathbb{Z}/2\mathbb{Z}$  cohomology),

$$w_1(TX) = w_1(N_M X)$$

Furthermore, the Stiefel-Whitney classes vanish exactly when the bundle is orientable so we see that  $TX$  is orientable if and only if  $N_M X$  is orientable.  $\square$

*Remark.* In general, we see that,

$$w_1(N_M X) = w_1(TX) + w_1(TM)|_X$$

and therefore we can compute the two-sidedness from the orientability of  $X$  together with the pullback of the Stiefel-Whitney class of  $TM$ .

*Remark.* This paper gives lots of examples of non-orientable surfaces such as Möbius strips and Klein bottles with two-sided embeddings into non-orientable 3-manifolds.

## 15 Stable Parallelizability of Spheres

**Proposition 15.0.1.** Let  $X$  be a  $n$ -dimensional oriented surface and  $\iota : X \rightarrow \mathbb{R}^{n+1}$  an immersion. Then  $TX$  is stably trivial.

*Proof.* The canonical exact sequence,

$$0 \longrightarrow TX \longrightarrow \iota^*TY \longrightarrow N_YX \longrightarrow 0$$

splits (every sequence splits) to give  $\iota^*TY = TX \oplus N_YX$  but  $TY \cong \varepsilon^{n+1}$  is trivial. Because  $X$  and  $Y$  are orientable, the embedding  $\iota : X \rightarrow Y$  is two-sided so  $N_YX$  is orientable. However,  $N_YX$  is a line bundle since  $\dim Y = \dim X + 1$  and thus  $N_YX$  is trivial. Therefore,

$$TX \oplus \mathbb{R} \cong \mathbb{R}^{n+1}$$

and thus  $TX$  is stably trivial so  $X$  is stably parallelizable.  $\square$

**Corollary 15.0.2.** All spheres have stably trivial tangent bundles in fact  $TS^n \oplus \mathbb{R}$  is trivial.

*Remark.* This shows that  $S^n \times \mathbb{R}$  is parallelizable. For the case  $n = 2$ , this has to be true because it is a theorem of Thurston that every orientable 3-manifold has trivial tangent bundle.

## 16 Some Questions

### 16.1 Can I use Miracle Flatness at Only Closed Points

### 16.2 Do Regular Functions Separate Points

Throughout we assume that  $X$  is separated. Otherwise points of  $X$  cannot be separated by rational functions let alone regular functions. For simplicity, we assume that  $X$  is noetherian.

I can reduce to the case of an integral scheme as follows. First, if  $X$  is nonreduced then  $X_{\text{red}} \rightarrow X$ . Suppose that  $X_{\text{red}}$  separated points then any function mapping to it on  $X$  will work. Now we assume  $X$  is reduced. If  $X$  is reducible then there are two cases to consider. If  $x, y \in X$  lie on different irreducible components then there exist disjoint opens containing the two points so they can clearly be separated by regular functions. If  $x, y \in X$  lie in the same irreducible component then we reduce to that irreducible component. If there is some  $x, y \in U \subset Z$  open in  $Z$  with a regular function separating  $x, y$  then (UGH NOT QUITE).

#### 16.2.1 The Integral Case

Let  $X$  be an integral separated scheme. Then points of  $X$  are separated by rational functions. Indeed, I claim that if  $\mathcal{O}_{X,x} = \mathcal{O}_{X,y}$  inside the function field  $K(X)$  then  $x = y$ . In fact, I will show that neither can dominate the other. Suppose that  $\mathcal{O}_{X,x} \subset \mathcal{O}_{X,y}$  with  $\mathfrak{m}_y \cap \mathcal{O}_{X,x} = \mathfrak{m}_x$  (we say  $\mathcal{O}_{X,y}$  dominates  $\mathcal{O}_{X,x}$ ) then there is a valuation ring  $A$  dominating  $\mathcal{O}_{X,y}$  inside  $K(X)$  giving maps,

$$\begin{array}{ccc} \text{Spec}(K(X)) & \longrightarrow & X \\ \downarrow & \nearrow & \downarrow \\ \text{Spec}(A) & \longrightarrow & \text{Spec}(\mathbb{Z}) \end{array}$$

via sending  $\mathfrak{m}_A \mapsto x$  and the local ring map  $\mathcal{O}_{X,x} \hookrightarrow A$  and by  $\mathfrak{m}_A \mapsto y$  and the local ring map  $\mathcal{O}_{X,y} \hookrightarrow A$ . By the valuative criterion of separatedness, there is at most one such dotted map and thus  $x = y$  and  $\mathcal{O}_{X,x} = \mathcal{O}_{X,y}$ .

Now I claim that if  $\mathcal{O}_{X,x} \subset \mathcal{O}_{X,y}$  then  $x, y$  are contained in the same affine open and  $y \rightsquigarrow x$ . Indeed, the prime ideal  $\mathfrak{p} = \mathfrak{m}_y \cap \mathcal{O}_{X,x}$  corresponds to some point  $y' \in \text{Spec}(\mathcal{O}_{X,x})$  which lies in every affine open containing  $x$ . Then  $\mathcal{O}_{X,x} \subset \mathcal{O}_{X,y'} \subset \mathcal{O}_{X,y}$  because  $\mathcal{O}_{X,y'} = (\mathcal{O}_{X,x})_{\mathfrak{p}}$  which are units in  $\mathcal{O}_{X,y}$ . Then  $\mathcal{O}_{X,y'} \hookrightarrow \mathcal{O}_{X,y}$  is local so by our previous result  $y = y'$ .

I would like to improve this to give a rational function whose value at  $x$  is 1 and at  $y$  is 0. We need to know that  $\mathfrak{m}_x \cap \mathcal{O}_{X,y} \neq \mathfrak{m}_y \cap \mathcal{O}_{X,x}$ .

## 17 Finite Intersection Property

**Definition 17.0.1.** A collection of sets  $\{K_\alpha\}$  has the finite intersection property if every finite intersection is nonempty,

$$K_{\alpha_1} \cap \cdots \cap K_{\alpha_n} \neq \emptyset$$

**Proposition 17.0.2.** A space is compact if and only if every collection  $\{K_\alpha\}$  of closed subsets with the finite intersection property has,

$$\bigcap_{\alpha} K_{\alpha} \neq \emptyset$$

*Proof.* We let  $U_\alpha = K_\alpha$  where  $U_\alpha$  is open iff  $K_\alpha$  is closed. Then  $\{U_\alpha\}$  has no finite subcover iff each,

$$U_{\alpha_1} \cap \cdots \cap U_{\alpha_n} \subsetneq X \iff K_{\alpha_1} \cap \cdots \cap K_{\alpha_n} \neq \emptyset$$

iff  $\{K_\alpha\}$  has FIP. Furthermore,  $\{U_\alpha\}$  is a cover iff,

$$\bigcup_{\alpha} U_{\alpha} = X \iff \bigcap_{\alpha} K_{\alpha} = \emptyset$$

Therefore every cover has a finite subcover is equivalent to every collection of opens without a finite subcover is not a cover which is equivalent to every collection of closed sets with FIP has nonempty intersection.  $\square$

**Proposition 17.0.3.** Let  $X$  be a topological space. Let  $\{K_\alpha\}$  be a collection of sets with the FIP such that one of the following holds,

- (a) the  $K_\alpha$  are closed and some  $K_{\alpha_0}$  is compact
- (b)  $X$  is Hausdorff and  $K_\alpha$  are compact.

Then,

$$\bigcap_{\alpha} K_{\alpha} \neq \emptyset$$

*Proof.* Because compact sets are closed in a Hausdorff space condition (b) implies (a). Therefore, it suffices to prove the conclusion assuming (a). By FIT  $K_{\alpha_0}$  is nonempty and  $\{K_{\alpha_0} \cap K_\alpha\}$  is a collection of closed sets with FIP in the compact set  $K_{\alpha_0}$  and therefore,

$$\bigcap_{\alpha} K_{\alpha} = \bigcap_{\alpha} (K_{\alpha_0} \cap K_{\alpha}) \neq \emptyset$$

by the previous proposition.  $\square$

**Corollary 17.0.4.** Let  $I_n \subset \mathbb{R}$  be a sequence of nonempty nested closed intervals. Then,

$$\bigcap_n I_n \neq \emptyset$$

## 18 When Are Isometries Smooth

**Definition 18.0.1.** Let  $f : X \rightarrow Y$  be a map between metric spaces. We say that  $f$  is *isometric* if for all  $x_1, x_2 \in X$ ,

$$d_Y(f(x_1), f(x_2)) = d_X(x_1, x_2)$$

**Proposition 18.0.2.** Isometric maps are uniformly continuous with uniformly continuous inverse on their image.

*Proof.* Uniform continuity is immediate because,

$$d_X(x_1, x_2) < \epsilon \iff d_X(f(x_1), f(x_2))$$

Furthermore,  $f$  is injective because if  $f(x_1) = f(x_2)$  then,

$$d_X(x_1, x_2) = d_Y(f(x_1), f(x_2)) = 0$$

so  $x_1 = x_2$ . Finally, it is clear that  $f^{-1}$  is also isometric and thus uniformly continuous.  $\square$

**Proposition 18.0.3.** Isometric maps  $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$  are affine orthogonal transformations.

*Proof.* First we may translate  $f$  such that  $f(0) = 0$ . For  $x, y \in \mathbb{R}^n$  consider,

$$\|f(x) - f(y)\|^2 = \|f(x)\|^2 - 2\langle f(x), f(y) \rangle + \|f(y)\|^2 = \|x\|^2 - 2\langle f(x), f(y) \rangle + \|y\|^2$$

however,

$$\|f(x) - f(y)\|^2 = \|x - y\|^2 = \|x\|^2 - 2\langle x, y \rangle + \|y\|^2$$

and therefore,

$$\langle f(x), f(y) \rangle = \langle x, y \rangle$$

so  $f$  preserves the inner product. Furthermore we can show that  $f$  is linear as follows.

$$\begin{aligned} \|f(x+y) - f(x) - f(y)\|^2 &= \|f(x+y)\|^2 - 2\langle f(x+y), f(x) \rangle - 2\langle f(x+y), f(y) \rangle + \|f(x) + f(y)\|^2 \\ &= \|x+y\|^2 - 2\langle x+y, x \rangle - 2\langle x+y, y \rangle + \|f(x)\|^2 + 2\langle f(x), f(y) \rangle + \|f(y)\|^2 \\ &= \|x+y\|^2 - 2\langle x+y, x+y \rangle + \|x\|^2 + \langle x, y \rangle + \|y\|^2 \\ &= \|x+y\|^2 - 2\|x+y\|^2 + \|x+y\|^2 = 0 \end{aligned}$$

and therefore  $f(x+y) = f(x) + f(y)$ . Furthermore,

$$\begin{aligned} \|f(\lambda x) - \lambda f(x)\|^2 &= \|f(\lambda x)\|^2 - 2\langle f(\lambda x), \lambda f(x) \rangle + \lambda^2 \|f(x)\|^2 = \|\lambda x\|^2 - 2\lambda \langle \lambda x, x \rangle + \lambda^2 \|x\|^2 \\ &= \lambda^2 \|x\|^2 - 2\lambda \|x\|^2 + \|x\|^2 = 0 \end{aligned}$$

and therefore  $f(\lambda x) = \lambda f(x)$  so  $f$  is linear and  $\langle f(x), f(y) \rangle = \langle x, y \rangle$  so  $f$  is orthogonal and therefore invertible.  $\square$

**Theorem 18.0.4** (Myers-Steenrod Theorem). Let  $M$  and  $N$  be Riemannian manifolds with induced metrics and  $\phi : M \rightarrow N$  is a surjective distance preserving map then  $\phi$  is a smooth isometry.

## 19 Tangent Spaces a la EGA

Grothendieck defines the tangent space in a kinda funny way. First define the tangent bundle,

$$T_{X/S} = \mathbb{V}_X(\Omega_{X/S}) = \mathbf{Spec}_X \left( \mathrm{Sym}(\Omega_{X/S}) \right)$$

Then the tangent space at a point are the  $\mathrm{Spec}(\kappa(x))$ -points over  $X$  (which form a vector space). This is,

$$T_{X/S}(x) = \mathrm{Hom}_X \left( \mathrm{Spec}(\kappa(x)), \mathbb{V}_X(\Omega_{X/S}) \right) = \mathrm{Hom}_{\mathcal{O}_X} \left( \Omega_{X/S}, \iota_* \mathcal{O}_{\kappa(x)} \right) = \mathrm{Hom}_{\kappa(x)} \left( (\Omega_{X/S})_x \otimes \kappa(x), \kappa(x) \right)$$

is the  $\kappa(x)$ -dual of  $(\Omega_{X/S})_x \otimes_{\mathcal{O}_{X,x}} \kappa(x)$  which does make sense but doesn't agree with the Zariski tangent space in general. Maybe this is better because it has the correct dimension at the generic point unlike the Zariski tangent space.

Grothendieck shows [EGA IV<sub>4</sub> 16.5.13.2] that if  $\kappa(s) \rightarrow \kappa(x)$  is an isomorphism then,

$$T_{X/S}(X) \xrightarrow{\sim} \mathrm{Hom}_{\kappa(x)} \left( \mathfrak{m}'_x / \mathfrak{m}_x^2, \kappa(x) \right)$$

where  $\mathfrak{m}'_x$  is the maximal ideal of the local ring  $\mathcal{O}_{X_s, x} = \mathcal{O}_{X, x} / \mathfrak{m}_s \mathcal{O}_{X, x}$ . This means that  $T_{X/S}(x) = T_{X_s/s}(x)$  as it should. This is also Pset 9 problem 2 in Johan's first class.

I claim that this result actually holds more generally. Because  $T_{X/S}(X) = T_{X_s/s}(x)$  in general I can work with  $X_s \rightarrow \mathrm{Spec}(\kappa(s))$  and assume that  $X$  is over a field  $k$ . Then I claim that if  $\kappa(x)/k$  is a separable algebraic extension then,

$$T_{X/k}(x) \xrightarrow{\sim} \mathrm{Hom}_{\kappa(x)} \left( \mathfrak{m}_x / \mathfrak{m}_x^2, \kappa(x) \right)$$

Indeed this is an immediate consequence of [H, Ex. II 8.1(a)].

Notice that if  $\kappa(x)/k$  is not algebraic this immediately fails e.g. the generic  $x \in \mathbb{A}_k^1$  point has  $\mathfrak{m}_x / \mathfrak{m}_x^2 = (0)$  but  $T_{\mathbb{A}^1/k}(x) \cong k(t)$ . Furthermore, if  $\kappa(x)/k$  is not separable this also fails e.g. consider  $X = \mathrm{Spec}(\mathbb{F}_p(t^{\frac{1}{p}})) \rightarrow \mathrm{Spec}(\mathbb{F}_p(t))$ . Then  $\Omega_{X/k} \cong \mathbb{F}_p(t^{\frac{1}{p}})$  but  $\mathfrak{m}_x / \mathfrak{m}_x^2 = 0$ .

### 19.0.1 The Submersion Theorem

DO THIS TOMORROW!!

## 20 Genus and Reduction

**Lemma 20.0.1.** Let  $A_1, A_2$  be rings. Then there is an equivalence of categories,

$$\mathbf{Mod}_{A_1} \times \mathbf{Mod}_{A_2} \rightarrow \mathbf{Mod}_{A_1 \times A_2}$$

given by sending  $(M_1, M_2) \mapsto (M_1)_{A_1 \times A_2} \oplus (M_2)_{A_1 \times A_2}$

*Proof.* Faithfulness is clear. For fullness, notice that if  $\phi : M_1 \oplus M_2 \rightarrow N_1 \oplus N_2$  is a morphism of  $A_1 \times A_2$ -modules then letting  $e_i \in A_i$  be the identity because  $\phi(e_i \cdot m) = e_i \cdot \phi(m)$  we see that  $\phi$  is represented by a diagonal matrix of morphisms proving fullness. Finally, let  $M$  be an  $A_1 \times A_2$ -module. Then  $M = e_1 M \oplus e_2 M$  because  $e_1 + e_2 = 1$  and  $e_1^2 = e_1$  and  $e_2^2 = e_2$  and  $e_1 e_2 = 0$  and  $e_i M$  is naturally an  $A_i$ -module proving fullness.  $\square$

**Lemma 20.0.2.** Let  $A$  be a Noetherian ring and  $M$  a finite  $A$ -module such that  $\dim \operatorname{Supp}_A(M) = 0$ . Then  $\operatorname{Supp}_A(M)$  is finite and,

$$M \cong \bigoplus_{\mathfrak{p} \in \operatorname{Supp}_A(M)} M_{\mathfrak{p}}$$

*Proof.* Because  $M$  is finite type we have  $\operatorname{Supp}_A(M) = V(\operatorname{Ann}_A(M))$ . Let  $B = A/\operatorname{Ann}_A(M)$  then  $B$  is noetherian and  $\dim B = \dim \operatorname{Supp}_A(M) = 0$  so  $B$  is Artinian. Therefore  $\operatorname{Supp}_A(M) = \operatorname{Spec}(B)$  is finite and consists of the maximal ideals of  $B$ . Then by the Chinese remainder theorem,

$$B \cong \prod_{\mathfrak{m} \in \operatorname{Spec}(B)} B_{\mathfrak{m}}$$

and therefore,

$$M \cong \bigoplus_{\mathfrak{m} \in \operatorname{Spec}(B)} M_{\mathfrak{m}}$$

□

**Lemma 20.0.3.** Let  $X$  be a Noetherian scheme and  $\mathcal{F}$  a coherent  $\mathcal{O}_X$ -module with,

$$\dim \operatorname{Supp}_{\mathcal{O}_X}(\mathcal{F}) = 0$$

Then,  $\operatorname{Supp}_{\mathcal{O}_X}(\mathcal{F})$  is finite and,

$$\mathcal{F} \cong \bigoplus_{x \in \operatorname{Supp}_{\mathcal{O}_X}(\mathcal{F})} (\iota_x)_* \mathcal{F}_x$$

where  $\iota_x : \operatorname{Spec}(\mathcal{O}_{X,x}) \rightarrow X$  is the canonical map and  $\mathcal{O}_{X,x}/\operatorname{Ann}_{\mathcal{O}_{X,x}}(\mathcal{F}_x)$  is Artin local.

*Remark.* Notice that because the  $\mathcal{O}_{X,x}$ -module  $\mathcal{F}_x$  is supported only over the maximal ideal that  $(\iota_x)_* \mathcal{F}_x$  is the same (viewing  $\mathcal{F}_x$  as an abelian sheaf) as pushing forward along the map  $x \hookrightarrow X$ .

*Proof.* Because  $X$  is quasi-compact, we can choose a finite affine open cover  $U_i$  on which  $\mathcal{F}|_{U_i} = \widetilde{M}_i$ . Then the result follows immediately from the previous lemma. Notice further that there is a canonical map,

$$\mathcal{F} \rightarrow \prod_{x \in \operatorname{Supp}_{\mathcal{O}_X}(\mathcal{F})} (\iota_x)_* \mathcal{F}_x$$

from adjunction and the universal property of the product. Finiteness shows that this is a direct sum and we can check locally that it is an isomorphism. (DO THIS BETTER!!) □

**Proposition 20.0.4.** Suppose that  $X$  is a finite type scheme over  $k$  and  $\mathcal{I} \subset \mathcal{O}_X$  a quasi-coherent ideal sheaf which is supported only at closed points. Let  $Z \subset X$  be the closed subscheme determined by  $Z$ . Then,  $\operatorname{Supp}_{\mathcal{O}_X}(\mathcal{I})$  is a finite set and,

$$\chi(X, \mathcal{O}_X) - \chi(Z, \mathcal{O}_Z) = \sum_{x \in \operatorname{Supp}_{\mathcal{O}_X}(\mathcal{I})} \dim_k \mathcal{I}_x$$

*Proof.* Consider the exact sequence of the closed immersion  $\iota : Z \hookrightarrow X$ ,

$$0 \longrightarrow \mathcal{I} \longrightarrow \mathcal{O}_X \longrightarrow \iota_* \mathcal{O}_Z \longrightarrow 0$$



Therefore,

$$\chi(X, \mathcal{O}_X) - \chi(Z, \mathcal{O}_Z) = \chi(X, \mathcal{I})$$

using that  $\iota$  is affine so  $H^i(X, \iota_* \mathcal{F}) = H^i(Z, \mathcal{F})$ . Furthermore, because  $X$  is noetherian,  $\mathcal{I}$  is coherent and thus  $\text{Supp}_{\mathcal{O}_X}(\mathcal{I})$  is closed but only contains closed points. Therefore, writing  $\text{Supp}_{\mathcal{O}_X}(\mathcal{I})$  as a union of finitely many irreducible components we see that these must be points (they are irreducible and only contain closed points) and thus  $Y = \text{Supp}_{\mathcal{O}_X}(\mathcal{I})$  consists of a finite number of closed points and is zero dimensional. Therefore,

$$\mathcal{I} \cong \bigoplus_{x \in \text{Supp}_{\mathcal{O}_X}(\mathcal{I})} (\iota_x)_* \mathcal{I}_x$$

Thus,

$$\chi(X, \mathcal{I}) = \sum_{x \in \text{Supp}_{\mathcal{O}_X}(\mathcal{I})} \dim_k \mathcal{I}_x$$

where the higher cohomology of  $\mathcal{I}$  vanishes because  $\text{Supp}_{\mathcal{O}_X}(\mathcal{I})$  is zero dimensional.  $\square$

**Corollary 20.0.5.** Suppose that  $X$  is a finite type scheme over  $k$  which is reduced at all non-closed points. Then, the ideal sheaf  $\mathcal{N}$  of  $X_{\text{red}} \hookrightarrow X$  is supported at finitely many points and,

$$\chi(X, \mathcal{O}_X) - \chi(X_{\text{red}}, \mathcal{O}_{X_{\text{red}}}) = \sum_{x \in \text{Supp}_{\mathcal{O}_X}(\mathcal{N})} \dim_k \mathcal{N}_x$$

*Proof.* Consider the exact sequence of the closed immersion  $X_{\text{red}} \hookrightarrow X$  which is also a homeomorphism,

$$0 \longrightarrow \mathcal{N} \longrightarrow \mathcal{O}_X \longrightarrow \mathcal{O}_{X_{\text{red}}} \longrightarrow 0$$

Therefore,

$$\chi(X, \mathcal{O}_X) - \chi(X_{\text{red}}, \mathcal{O}_{X_{\text{red}}}) = \chi(X, \mathcal{N})$$

Furthermore, because  $\mathcal{O}_{X,x}$  is reduced unless  $x \in X$  is closed, we see that  $\text{Supp}_{\mathcal{O}_X}(\mathcal{N})$  is zero dimensional. Explicitly, there is an exact sequence,

$$0 \longrightarrow \mathcal{N}_x \longrightarrow \mathcal{O}_{X,x} \longrightarrow \mathcal{O}_{X_{\text{red}},x} \longrightarrow 0$$

and  $\mathcal{O}_{X_{\text{red}},x} = (\mathcal{O}_{X,x})_{\text{red}}$  so we have  $\mathcal{N}_x = \text{nilrad}(\mathcal{O}_{X,x})$  vanishes if and only if  $\mathcal{O}_{X,x}$  is reduced. Therefore,  $\text{Supp}_{\mathcal{O}_X}(\mathcal{N})$  is supported only at closed points so we can apply the proposition to conclude.  $\square$

**Corollary 20.0.6.** Let  $X$  be a projective scheme with ample line bundle  $\mathcal{O}_X(1)$ . Let  $\mathcal{I}$  be an ideal sheaf on  $X$  which is supported only at closed points. Let  $Z \subset X$  be the closed subscheme determined by  $\mathcal{I}$ . Then  $\text{Supp}_{\mathcal{O}_X}(\mathcal{I})$  is a finite set and,

$$P_X(n) - P_Z(n) = \sum_{x \in \text{Supp}_{\mathcal{O}_X}(\mathcal{I})} \dim_k \mathcal{I}_x$$

where  $P_X$  and  $P_Z$  are the Hilbert polynomials. In particular, if  $\dim X > 0$  we see that  $X$  and  $Z$  have the same dimension and degree.

*Proof.* This follows from the fact that,

$$0 \longrightarrow \mathcal{I}(n) \longrightarrow \mathcal{O}_X(n) \longrightarrow \iota_* \mathcal{O}_Z(n) \longrightarrow 0$$

is exact and  $\mathcal{J}(n) \cong \mathcal{J}$  because  $\mathcal{O}_X(n)$  is a line bundle and thus its stalk is free of rank 1 at each  $x \in X$ . Therefore,

$$P_X(n) - P_Z(n) = \chi(X, \mathcal{O}_X(n)) - \chi(Z, \mathcal{O}_Z(n)) = \chi(X, \mathcal{J})$$

so we conclude by the previous calculation.  $\square$

**Corollary 20.0.7.** Let  $X$  be a projective scheme with ample line bundle  $\mathcal{O}_X(1)$ . Suppose that  $X$  is reduced at all non-closed points. Then  $\mathcal{N}$  is supported at finitely many points and,

$$P_X(n) - P_{X_{\text{red}}}(n) = \sum_{x \in \text{Supp}_{\mathcal{O}_X}(\mathcal{N})} \dim_k \mathcal{N}_x$$

In particular, if  $\dim X > 0$  we see that  $X$  and  $X_{\text{red}}$  have the same dimension and degree.

## 20.1 The Application to Projective Cones

Let  $X \subset \mathbb{P}^n = \text{Proj}(S)$  be cut out by the saturated ideal  $I \subset S$ . Then we consider the projective cone  $C(X) = \text{Proj}(S[z]/I[z])$ . If  $X$  is integral then  $S/I \hookrightarrow \Gamma_*(\mathcal{O}_X)$  which is a domain and thus  $S/I$  is a domain so  $I$  is prime. Thus  $S[z]/I[z] = (S/I)[z]$  is a domain so  $C(X)$  is integral. Likewise if  $X$  is reduced then  $S/I \hookrightarrow \Gamma_*(\mathcal{O}_X)$  is reduced so  $I$  is a radical ideal.

*Remark.* This only holds for the *saturated* ideal  $I$ . For example, consider  $I = (xy, x^2) \subset k[x, y]$ . Then  $I^{\text{sat}} = (x)$  is a prime ideal and  $\text{Proj}(k[x, y]/I) = \text{Proj}(k[x, y]/I^{\text{sat}}) = \text{Spec}(k)$  is integral but  $I$  is clearly not prime.

If we choose the wrong ideal  $I$  then we get the wrong cone  $C_I(X) = \text{Proj}(S[z]/I[z])$  because  $I[z]$  and  $I^{\text{sat}}[z]$  need not have the same saturation. However, the problem is completely supported at the origin because  $I$  and  $I^{\text{sat}}$  become equal after localization at any ideal not containing some  $x_i$ . Therefore, the closed immersion  $C(X) \hookrightarrow C_I(X)$  is cut out by the nilpotent ideal sheaf  $I^{\text{sat}}/I$  supported at the origin and we see  $C_I(X)_{\text{red}} = C(X)_{\text{red}}$ . Therefore,

$$P_{C_I(X)}(n) - P_{C(X)}(n) = \dim_k(I^{\text{sat}}/I)_{(\mathfrak{m})}$$

I SHOULD CHECK THIS PART!!!

**Example 20.1.1** (H, Example, 9.8.4). we have a twisted cubic degenerating to a nodal cubic with reduced shit. Let  $X_0$  have an affine patch cut out by,

$$I_0 = (z^2, yz, xz, y^2 - x^2(x+1))$$

Then the reduction is cut out by the ideal  $N = (z)$  so  $\dim_k N/I_0 = 1$ . Let  $C = (X_0)_{\text{red}}$  be the nodal cubic. Then,

$$P_C(n) = 3n + 0$$

because it is a Cartier divisor in  $\mathbb{P}^2$  of degree 3 so,

$$\chi(\mathcal{O}_C) = 1 - g_a = 0$$

Because  $X \rightarrow \mathbb{A}^1$  is a flat family,  $P_{X_0}$  is equal to  $P_{X_1}$  where  $X_1$  is the twisted cubic given parametrically by  $(t^2 - 1, t^3 - t, t)$  which is isomorphic to the twisted cubic curve  $(t, t^2, t^3)$ . This has ideal  $I$  the relations for the functions  $s^3, s^2t, st^2, t^3$  meaning the kernel of the map,

$$k[x_0, x_1, x_3, x_4] \rightarrow k[s, t]$$

sending  $x_0 \mapsto s^3, x_1 \mapsto s^2t, x_3 \mapsto st^2, x_4 \mapsto t^3$  and therefore the quotient is isomorphic to its image,

$$S/I \cong k[s^3, s^2t, st^2, t^3]$$

where we give  $s, t$  degree  $\frac{1}{3}$  (i.e. view it as a subring of  $(k[s, t])^{(3)}$ ) to make this a graded isomorphism. We see that the ideals  $I_a$  are homogeneous and prime since they are the kernel of a map of graded domains (the quotient of such a kernel is a subring of a domain and hence a domain) and therefore saturated (if  $\mathfrak{p}$  is a prime ideal not containing the irrelevant ideal and  $x_i^n f \in \mathfrak{p}$  then either  $x_i \in \mathfrak{p}$  for each  $i$  or  $f \in \mathfrak{p}$ ). Therefore,

$$\dim_k(S_0/I_0)_d = \dim_k(k[s^4, s^3t, st^3, t^4])_d = \begin{cases} 1 & d = 0 \\ 3d + 1 & d > 0 \end{cases}$$

Therefore,

$$P_{X_0} = 3d + 1$$

showing that,

$$P_{X_0}(n) - P_C(n) = 1 = \dim_k(N/I_0)$$

as expected.

## 21 Local Systems

*Remark.* We want to prove the following claim: let  $\mathcal{L}$  be a local system of  $A$ -modules valued in  $M$  on a topological space  $X$  (or Grothendieck topology) then  $\chi(X, \mathcal{L}) = \chi(X, \underline{A})$ .

**Definition 21.0.1.** Suppose that  $G_0(A)$  is equipped with a rank function  $\text{rank}_A : K_0(A) \rightarrow \mathbb{Z}$ . Then for a sheaf of  $A$ -modules we define,

$$\chi(X, \mathcal{F}) = \sum_{i=0}^{\infty} \text{rank}_A H^i(X, \mathcal{F})$$

when  $H^i(X, \mathcal{F})$  are finite  $A$ -modules and there is vanishing of  $H^i(X, \mathcal{F})$  for sufficiently large  $i$ .

*Remark.* In the case that  $A$  is a domain there is always a rank function  $M \mapsto \dim_K(M \otimes_A K)$  where  $K = \text{Frac}(A)$  which descends to  $G_0(A)$  because it is additive over short exact sequences.

### 21.1 The Case of a Topological Space

**Proposition 21.1.1** (Mayer-Vietoris).

**Proposition 21.1.2.** Let  $\mathcal{L}$  be a local system valued in an abelian group  $A$  on a topological space  $X$  then  $\chi(X, \mathcal{L}) = \chi(X, \underline{A})$ .

*Proof.* Consider the □

**Corollary 21.1.3.** If  $X$  is a (INSET CORRECT TOP PROPERTY) space and  $\mathcal{L}$  is a local system on  $X$  valued in an abelian group  $A$  then,

$$\chi(X, \mathcal{L}) = \chi(X, \underline{A}) = \sum_{i=0}^n \text{rank}$$

## 21.2 The General Case of a Site

## 22 Images of Maximal Ideals

**Lemma 22.0.1.** Let  $X$  be a locally finite type scheme over  $k$ . Then  $x \in X$  is closed if and only if  $\kappa(x)/k$  is finite.

*Proof.* If  $x \in X$  is closed then choose an affine open  $x \in \text{Spec}(A)$  with  $A$  a finite type  $k$ -algebra. Then  $x = \mathfrak{m} \in \text{Spec}(A)$  is closed so  $\mathfrak{m}$  is maximal so  $A/\mathfrak{m}$  is a finitely generated  $k$ -algebra and a field so  $\kappa(x) = A/\mathfrak{m}$  is a finite  $k$ -extension by the Nullstellensatz.

Conversely, if  $\kappa(x)/k$  is finite then for every affine open  $x \in \text{Spec}(A)$  we see that  $A/\mathfrak{p} \hookrightarrow (A/\mathfrak{p})_{\mathfrak{p}}$  because  $A/\mathfrak{p}$  is a domain but  $\kappa(x) = (A/\mathfrak{p})_{\mathfrak{p}}$  is finite dimensional over  $k$  and so  $A/\mathfrak{p}$  is a finite dimensional  $k$ -algebra domain and hence a field so  $\mathfrak{p}$  is maximal. Thus  $x$  is closed in each  $\text{Spec}(A)$  and since these cover  $X$  we see that  $x \in X$  is closed.  $\square$

**Proposition 22.0.2.** A map of schemes locally of finite type over a field  $k$  sends closed points to closed points.

*Proof.* A point  $x \in A$  being closed is equivalent to  $\kappa(x)/k$  being finite by the Nullstellensatz. Then  $\kappa(f(x)) \hookrightarrow \kappa(x)$  so the image of a closed point is closed.  $\square$

*Remark.* In general this is false even for finite type maps. For example, consider  $\text{Spec}(\mathbb{Q}_p) \rightarrow \text{Spec}(\mathbb{Z}_p)$  which is finite type since  $\mathbb{Q}_p = \mathbb{Z}_p[p^{-1}]$ . However, we give an extension of this result to arbitrary schemes.

**Lemma 22.0.3.** Let  $\varphi : A \rightarrow B$  be a finite type ring map and  $\mathfrak{m} \subset B$  a maximal ideal and  $\mathfrak{p} = \varphi^{-1}(\mathfrak{m})$  the image. Then there exists  $t \in A \setminus \mathfrak{p}$  such that  $\mathfrak{p}A_t$  is maximal.

*Proof.* Let  $K = \text{Frac}(A/\mathfrak{p})$ . The maximal ideal  $\mathfrak{m} \in \text{Spec}(B)$  corresponds to a maximal ideal  $\bar{\mathfrak{m}} \in \text{Spec}(B')$  with  $B' = B \otimes_A (A/\mathfrak{p})_{\mathfrak{p}} = B \otimes_A K$  which is the fiber. Then  $B'/\bar{\mathfrak{m}}' = (B/\mathfrak{m})_{\mathfrak{p}} = B/\mathfrak{m}$  because  $B/\mathfrak{m}$  is a field. Now  $B'$  is a finite type  $K$ -algebra because  $A \rightarrow B$  is finite type and thus  $K \rightarrow B \otimes_A K$  is finite type. Therefore, by the Nullstellensatz,  $B/\mathfrak{m}$  is a finite extension of  $K$ . Hence through  $A_{\mathfrak{p}} \rightarrow A_{\mathfrak{p}}/\mathfrak{p}A_{\mathfrak{p}} = K \rightarrow B/\mathfrak{m}$  we see that  $B/\mathfrak{m}$  is a finite  $A_{\mathfrak{p}}$ -module.

Choose generators  $x_1, \dots, x_n \in B/\mathfrak{m}$  as an  $A$ -algebra. Write  $B/\mathfrak{m} = A[x_1, \dots, x_n]/I$ . Since  $B/\mathfrak{m}$  is finite over  $A_{\mathfrak{p}}$  each  $x_i$  is integral over  $A_{\mathfrak{p}}$  so it satisfies some monic  $p_i \in A_{\mathfrak{p}}[x]$ . Let  $t \in A \setminus \mathfrak{p}$  be the product of the denominators of the coefficients of all  $p_i$ . Then  $p_i \in A_t[x]$  and thus  $x_i \in B/\mathfrak{m}$  is integral over  $A_t$  and hence  $A/\mathfrak{m}$  is a finite  $A_t$ -module since  $B/\mathfrak{m}$  is generated by finitely many integral elements as an  $A$ -algebra and hence as an  $A_t$ -algebra.

Consider  $A_t/\mathfrak{p}A_t = (A/\mathfrak{p})_t$ . Since  $\mathfrak{p}B \subset \mathfrak{m}$  we see that  $B/\mathfrak{m}$  is a finite  $A_t/\mathfrak{p}A_t$ -module. Then  $(A/\mathfrak{p})_t \subset K \subset B/\mathfrak{m}$  and thus  $K$  is finite over  $(A/\mathfrak{p})_t$ . Therefore,  $(A/\mathfrak{p})_t$  is a field because  $(A/\mathfrak{p})_t \subset K$  is an integral extension of domains with  $K$  a field and hence  $(A/\mathfrak{p})_t = K$  since  $K = \text{Frac}(A/\mathfrak{p})$ . Therefore  $\mathfrak{p}A_t$  is a maximal ideal.  $\square$

**Proposition 22.0.4.** Let  $f : X \rightarrow Y$  be a locally finite type map of schemes. The image of a locally closed point is locally closed.

*Proof.* Let  $x \in X$  be locally closed. Choose some affine open  $U = \text{Spec}(B)$  with  $x \in U$  closed and  $\text{Spec}(B) \rightarrow \text{Spec}(A)$  for  $\text{Spec}(A) \subset Y$  affine open. It corresponds to some maximal ideal  $\mathfrak{m} \in \text{Spec}(A)$  and therefore under the finite type ring map  $\varphi : A \rightarrow B$  there is  $t \in A \setminus \mathfrak{p}$  with  $\mathfrak{p} = \varphi^{-1}(\mathfrak{m})$  such that  $f(x) = \mathfrak{p}$  is closed in  $\text{Spec}(A_t)$  by the lemma.  $\square$

*Remark.* This is a corollary of Chevallay's theorem.

## 23 Degree of a Pullback of Curves

**Proposition 23.0.1.** Let  $f : X \rightarrow Y$  be a finite locally free morphism of proper schemes over  $k$ . Let  $\mathcal{E}$  be a vector bundle on  $Y$  then,

$$\chi(X, f^* \mathcal{E}) =$$

(HMMMMMM)

## 24 Unimodular Lattices

**Definition 24.0.1.** Let  $(V, \langle -, - \rangle)$  be a real inner-product space with  $n = \dim V$  finite. Then a *lattice* is a subgroup  $\Lambda \subset V$  such that  $\Lambda \otimes_{\mathbb{Z}} \mathbb{R} = V$ .

**Definition 24.0.2.** Let  $\Lambda$  be a lattice. Then we define the *dual Lattice*,

$$\Lambda^* \subset V^* \quad \text{where } \Lambda^* = \{\varphi \in V^* \mid \forall \gamma \in \Lambda : \varphi(\gamma) \in \mathbb{Z}\}$$

However,  $V$  is equipped with an inner product and under the natural isomorphism  $V \xrightarrow{\sim} V^*$  defined by  $v \mapsto \langle v, - \rangle$  we can identify,

$$\Lambda^* \subset V \quad \text{via} \quad \Lambda^* = \{v \in V \mid \forall \gamma \in \Lambda \mid \langle v, \gamma \rangle \in \mathbb{Z}\}$$

Thus we can write,

$$\begin{array}{ccc} \Lambda^* & \xrightarrow{\sim} & \text{Hom}(\Lambda, \mathbb{Z}) \\ \downarrow & \lrcorner & \downarrow \\ V^* & \longrightarrow & \text{Hom}(\Lambda, \mathbb{R}) \end{array}$$

**Definition 24.0.3.** The *covolume* or **DEFINE**

**Proposition 24.0.4.**  $|\Lambda| \cdot |\Lambda^*| = 1$

*Proof.* DO THIS!!

□

**Definition 24.0.5.** A lattice  $\Lambda$  is,

- (a) *integral* if  $\langle \gamma, \gamma' \rangle \in \mathbb{Z}$  for all  $\gamma, \gamma' \in \Lambda$
- (b) *unimodular* if  $|\Lambda| = 1$
- (c) *even* if  $\|\gamma\|^2 \in 2\mathbb{Z}$  for all  $\gamma \in \Lambda$
- (d) *self-dual* if  $\Lambda^* = \Lambda$  inside  $V$ .

**Lemma 24.0.6.** A lattice  $\Lambda$  is self-dual if and only if  $\Lambda$  is integral and unimodular.

*Proof.* If  $\Lambda$  is integral,  $\Lambda \subset \Lambda^*$  and if  $\Lambda$  is unimodular then  $|\Lambda^*| = |\Lambda| = 1$  proving that  $\Lambda = \Lambda^*$ . Conversely, if  $\Lambda = \Lambda^*$  then  $|\Lambda| = |\Lambda^*| = 1$  and  $\Lambda \subset \Lambda^*$  proving that  $\Lambda$  is unimodular and integral. □

**Definition 24.0.7.** Let  $\Lambda$  be a lattice. We define the theta function,

$$\Theta_\Lambda : \mathfrak{h} \rightarrow \mathbb{C}$$

via the infinite summation,

$$\Theta_\Lambda(\tau) = \sum_{\gamma \in \Lambda} e^{i\pi\tau\|\gamma\|^2}$$

**Proposition 24.0.8.** The summation form of  $\Theta_\Lambda$  is everywhere absolutely convergent on  $\mathfrak{h}$ .

*Proof.* Notice that,

$$|e^{i\pi\tau\|\gamma\|^2}| = e^{-\pi\|\gamma\|^2 \operatorname{Im}(\tau)}$$

Since  $\operatorname{Im}(\tau) > 0$  we see that  $0 < e^{-\pi\operatorname{Im}(\tau)} < 1$  and therefore because the number of lattice points of bounded norm grows polynomially the sum is convergent.  $\square$

**Theorem 24.0.9** (Poisson Summation). Let  $f : V \rightarrow \mathbb{C}$  be a Schwartz function with Fourier transform  $\hat{f} : V^* \rightarrow \mathbb{C}$ . Then,

$$\sum_{\gamma \in \Lambda} f(\gamma) = \frac{1}{|\Lambda|} \sum_{\varphi \in \Lambda^*} \hat{f}(\varphi)$$

**Proposition 24.0.10.** Let  $\Lambda$  be a lattice. For any  $\tau \in \mathfrak{h}$ ,

$$\Theta_{\Lambda^*}(-1/\tau) = \left(\frac{\tau}{i}\right)^{\frac{n}{2}} |\Lambda| \cdot \Theta_\Lambda(\tau)$$

*Proof.* This is a direct application of Poisson summation for  $f(v) = e^{i\pi\tau\|v\|^2}$ . A direct calculation shows that,

$$\hat{f}(v) = \left(\frac{i}{\tau}\right)^{\frac{n}{2}} e^{-i\pi\|v\|^2/\tau}$$

Then,

$$\Theta_\Lambda(\tau) = \sum_{\gamma \in \Lambda} f(\gamma) = \frac{1}{|\Lambda|} \sum_{\varphi \in \Lambda^*} \hat{f}(\varphi) = \frac{1}{|\Lambda|} \left(\frac{i}{\tau}\right)^{\frac{n}{2}} \Theta_{\Lambda^*}(-1/\tau)$$

$\square$

**Corollary 24.0.11.** If  $\Lambda$  is self-dual then,

$$\Theta_\Lambda(-1/\tau) = \left(\frac{\tau}{i}\right)^{\frac{n}{2}} \Theta_\Lambda(\tau)$$

**Proposition 24.0.12.** If  $\Lambda$  is integral then  $\Theta_\Lambda(\tau+2) = \Theta_\Lambda(\tau)$ . If  $\Lambda$  is even then  $\Theta_\Lambda(\tau+1) = \Theta_\Lambda(\tau)$ .

*Proof.* If  $\|\gamma\|^2 \in \mathbb{Z}$  then,

$$e^{i\pi(\tau+2)\|\gamma\|^2} = e^{2\pi i\|\gamma\|^2} e^{i\pi\tau\|\gamma\|^2} = e^{i\pi\tau\|\gamma\|^2}$$

Likewise, if  $\|\gamma\|^2 \in 2\mathbb{Z}$  then,

$$e^{i\pi(\tau+1)\|\gamma\|^2} = e^{\pi i\|\gamma\|^2} e^{i\pi\tau\|\gamma\|^2} = e^{i\pi\tau\|\gamma\|^2}$$

$\square$

**Corollary 24.0.13.** If  $\Lambda$  is self-dual and even then  $\Theta_\Lambda$  is modular.

**Theorem 24.0.14.** Let  $\Lambda$  be an even integral unimodular lattice. Then  $8 \mid \dim \Lambda$ .

*Proof.* Since integral unimodular lattices are self-dual we see that  $\Theta_\Lambda$  is modular.  $\square$

*Proof.* Let  $S, T \in \text{SL}2\mathbb{Z}$  describe  $T : \tau \mapsto \tau + 1$  and  $S : \tau \mapsto -1/\tau$ . The relation  $(ST)^3 = \text{id}$  describes the trajectory,

$$\tau \mapsto -\frac{1}{\tau} \mapsto \frac{\tau-1}{\tau} \mapsto \frac{\tau}{1-\tau} \mapsto \frac{1}{1-\tau} \mapsto \tau-1 \mapsto \tau$$

Using the modularity properties,

$$\Theta_\Lambda(\tau) = \left(\frac{i}{\tau-1}\right)^{\frac{n}{2}} \left(\frac{\tau-1}{i\tau}\right)^{\frac{n}{2}} \left(\frac{\tau}{i}\right)^{\frac{n}{2}} \Theta_\Lambda(\tau) = \left(\frac{1}{i}\right)^{\frac{n}{2}} \Theta_\Lambda(\tau)$$

Since  $\Theta_\Lambda(\tau) \neq 0$  because,

$$\Theta_\Lambda(i) = \sum_{\gamma \in \Lambda} e^{-\pi \|\gamma\|^2} > 0$$

and therefore, we must have  $i^{\frac{n}{2}} = 1$  and hence  $n$  is divisible by 8.  $\square$

*Remark.* All these numbers lie in a wedge on the complex plane (indeed  $\text{Re}(z) > 0$ ) and thus the function  $(-)^{\frac{n}{2}}$  is well-defined and is multiplicative.

## 25 Gradient Descent

### 25.1 Convex Functions

**Definition 25.1.1.** Let  $\Omega \subset V$  be a convex subset of a real vectorspace. Then  $f : \Omega \rightarrow \mathbb{R}$  is *convex* if,

$$f((1-t)x + ty) \leq (1-t)f(x) + tf(y)$$

for all  $t \in [0, 1]$  and *strictly convex* if for  $x \neq y$ ,

$$f((1-t)x + ty) < (1-t)f(x) + tf(y)$$

for  $t \in (0, 1)$ .

**Proposition 25.1.2.** Suppose that  $f : \Omega \rightarrow \mathbb{R}$  is convex and  $M$  the set of local minima. Then  $f(M)$  is at most one point. If  $f$  is strictly convex then  $M$  is at most one point.

*Remark.*  $f = 0$  is convex but not strictly convex showing that  $M$  can be large. Furthermore,  $M$  can be empty even for strictly convex functions. For example for  $f : \mathbb{R} \rightarrow \mathbb{R}$  via  $f(x) = x$ .

*Proof.* Suppose that  $x, y \in M$  with  $x \neq y$  and suppose that  $f(x) \neq f(y)$ . WLOG let  $\nabla = f(x) - f(y) > 0$ . Then,

$$g(t) = f((1-t)x + ty) \leq (1-t)f(x) + tf(y) = f(x) - t\nabla$$

By assumption, there exists some  $\epsilon > 0$  such that  $x$  is the minimum of  $f$  on  $B_\epsilon(x)$ . Choosing,

$$t = \frac{\epsilon}{2\|x - y\|}$$

We see that  $(1-t)x + ty = x + t(y-x) \in B_\epsilon(x)$  and therefore,

$$f(x) \leq f((1-t)x + ty) \leq f(x) - t\nabla$$

which is a contradiction to  $t > 0$  and  $\nabla > 0$ . Thus  $f(M)$  is empty or a single point. Furthermore, if  $f$  is strictly convex then let  $x, y \in M$  and assume that  $x \neq y$ . We showed that  $f(x) = f(y)$  so we see that for,

$$t \leq \frac{\epsilon}{\|x - y\|}$$

we see that,

$$f(x) \leq f((1-t)x + ty) < (1-t)f(x) + tf(y) = f(x)$$

which is a contradiction. Thus  $x = y$ . □

**Proposition 25.1.3.** If  $f$  is convex and  $(\nabla f)_x = 0$  then  $x$  is a global minimum.

*Proof.* For any  $y \in \Omega$  let  $\nabla = f(y) - f(x)$ . It suffices to prove that  $\nabla \geq 0$ . Consider,

$$g(t) = f((1-t)x + ty) \leq (1-t)f(x) + tf(y) = f(x) + \nabla t$$

Notice that  $g'(0) = 0$  and,

$$\frac{g(t) - g(0)}{t} \leq \nabla$$

and thus taking the limit,

$$g'(0) = \lim_{t \rightarrow 0} \frac{g(t) - g(0)}{t} \leq \nabla$$

and thus  $\nabla \geq g'(0) = 0$ . □

**Corollary 25.1.4.** Let  $f$  be convex and differentiable at  $x \in \Omega$ . Then for any  $y \in \Omega$ ,

$$f(y) \geq f(x) + (y - x) \cdot (\nabla f)_x$$

*Proof.* Define,

$$g(y) = f(y) - f(x) - (y - x) \cdot (\nabla f)_x$$

Then we see that  $g$  is convex and  $(\nabla g)_x = 0$  so  $x$  is a global minimum of  $g$  and  $g(x) = 0$  so  $g \geq 0$  proving the claim. □

## 25.2 Lipschitz Condition

**Definition 25.2.1.** We say that  $f : \Omega \rightarrow \mathbb{R}$  is Lipschitz-differentiable if  $f$  is differentiable and there is a constant  $L > 0$  such that for all  $x, y \in \Omega$ ,

$$\|(\nabla f)_x - (\nabla f)_y\| \leq L\|x - y\|$$

**Definition 25.2.2.** Let  $f : \Omega \rightarrow \mathbb{R}$  be a differentiable function. Then for some fixed  $\eta > 0$  we define the *gradient descent iteration* of  $x$  as,

$$x^+ = x - \eta(\nabla f)_x$$

**Lemma 25.2.3.** If  $f$  is Lipschitz-differentiable and  $\eta \leq L^{-1}$  then,

$$-\frac{3}{2}\eta\|(\nabla f)_x\|^2 \leq f(x^+) - f(x) \leq -\frac{1}{2}\eta\|(\nabla f)_x\|^2$$



*Proof.* Define,

$$g(t) = f((1-t)x + tx^+) = f(x - t\eta(\nabla f)_x)$$

Because  $g$  is continuously differentiable since  $f$  is continuously differentiable,

$$f(x^+) - f(x) = g(1) - g(0) = \int_0^1 g'(t) dt$$

However,

$$g'(t) = -\eta(\nabla f)_x \cdot (\nabla f)_{x(t)}$$

Therefore,

$$\begin{aligned} |f(x^+) - f(x) + \eta||(\nabla f)_x|^2| &= \eta \left| (\nabla f)_x \cdot \int_0^1 [(\nabla f)_x - (\nabla f)_{x(t)}] dt \right| \\ &\leq \eta ||(\nabla f)_x|| \int_0^1 ||(\nabla f)_x - (\nabla f)_{x(t)}|| dt \\ &\leq \eta L ||(\nabla f)_x|| \int_0^1 ||x - x(t)|| dt \\ &= \eta^2 L ||(\nabla f)_x||^2 \int_0^1 t dt \\ &= \frac{1}{2} \eta^2 L ||(\nabla f)_x||^2 \end{aligned}$$

If we take  $\eta < L^{-1}$  then completing the proof we find,

$$|f(x^+) - f(x) + \eta||(\nabla f)_x|^2| \leq \frac{1}{2} \eta ||(\nabla f)_x||^2$$

□

*Remark.* Therefore, defining a sequence  $x_{k+1} = x_k^+$  then either  $x_k$  hits a point with  $(\nabla f)_{x_k} = 0$  at which point the sequence stabilizes or  $f(x_k)$  forms a strictly decreasing sequence.

*Remark.* Consider  $f : \mathbb{R} \rightarrow \mathbb{R}$  given by  $f(x) = x$ . This has a constant derivative and hence we can take  $L$  to be any positive constant. However, we get the sequence  $x_k = x_0 - k\eta$  does not converge and gives an unbounded decreasing sequence  $f(x_k)$ .

**Corollary 25.2.4.** Suppose that in addition,  $f$  is bounded below. Then the decreasing sequence  $f(x_k)$  must converge to a limit. However, I claim that the sequence  $x_k$  need not converge to a limit. Indeed, consider the following example.

**Example 25.2.5.** Consider  $f : \mathbb{R} \rightarrow \mathbb{R}$  defined by,

$$f(x) = \begin{cases} 0 & x \leq 0 \\ \varphi(x) & 0 < x < 1 \\ \frac{1}{x} & x \geq 1 \end{cases}$$

where  $\varphi$  is a smooth function which interpolates to make  $f$  smooth. Then  $f$  has second derivative bounded so it is Lipschitz-differentiable and notice that  $f$  is even strictly convex on  $[1, \infty)$ . However, let  $x_0 = 1$ . Then we get the recurrence,

$$x_{k+1} = x_k + \frac{\eta}{x_k^2}$$

This is increasing so we just need to show that it is unbounded. Suppose that  $x_k < n$  for all  $n$ . Then  $x_{k+1} - x_k > \frac{\eta}{n^2}$  so for  $N > \eta^{-1}n^2(n - x_k)$  we would have  $x_{k+N} \geq n$  giving a contradiction proving that this sequence is indeed unbounded. However,  $f$  on  $\mathbb{R}$  we see that  $f$  achieves it minimum but is not convex on this region. We will see that if we have both convexity and achieving a minimum then this cannot happen.

*Remark.* Now we suppose that  $f$  achieves its minimum at some point  $x^*$ . Then we want to bound how quickly the distance between  $x$  and  $x^*$  decreases under  $x \mapsto x^+$ .

**Proposition 25.2.6.** Suppose that  $f : \Omega \rightarrow \mathbb{R}$  is convex and Lipschitz-differentiable with  $\eta < L^{-1}$ . Fix  $x^* \in \Omega$  then,

$$\|x - x^*\|^2 - \|x^+ - x^*\|^2 \geq 2\eta(f(x^+) - f(x^*)) \geq 0$$

*Proof.* We expand,

$$\|x^+ - x^*\|^2 = \|(x^+ - x) + (x - x^*)\|^2 = \|x^+ - x\|^2 + 2(x^+ - x) \cdot (x - x^*) + \|x - x^*\|^2$$

Therefore,

$$\|x - x^*\|^2 - \|x^+ - x^*\|^2 = -2(x^+ - x) \cdot (x - x^*) - \|x^+ - x\|^2 = 2\eta(\nabla f)_x \cdot (x - x^*) - \eta^2\|(\nabla f)_x\|^2$$

However, by previous lemmas,

$$f(x^*) - f(x) \geq (\nabla f)_x \cdot (x^* - x)$$

and also,

$$f(x^+) - f(x) \leq -\frac{1}{2}\eta\|(\nabla f)_x\|^2$$

Therefore,

$$\|x - x^*\|^2 - \|x^+ - x^*\|^2 \geq 2\eta(f(x) - f(x^*)) + 2\eta(f(x^+) - f(x)) = 2\eta(f(x^+) - f(x^*))$$

proving the claim.  $\square$

*Remark.* If  $x^*$  is a global (local suffices since  $f$  is convex) minimum of  $x^*$  (and hence  $f$  is bounded below) The above proposition implies that the sequence  $\|x - x^*\|$  is decreasing.

**Proposition 25.2.7.** Suppose that  $f : \Omega \rightarrow \mathbb{R}$  is convex and has a global minimum (and hence  $f$  is bounded below) and  $f$  is Lipschitz-differentiable with  $\eta < L^{-1}$ . Furthermore assume that the unit ball in  $V$  is precompact (e.g. if  $V$  is finite dimensional) then  $x_k \rightarrow x^*$  converges where  $x^*$  is a global minimum and,

$$f(x_k) - f(x^*) \leq \frac{\|x_0 - x^*\|^2}{2\eta k}$$

*Proof.* Let  $x'$  achieve the global minimum. Then the previous lemma shows that  $x_k \in B_r(x')$  where  $r = 2\eta(f(x_0) - f(x'))$ . Since  $\overline{B_r(x')}$  is compact we see that  $x_k$  has a convergent subsequence  $x_{k_i} \rightarrow x^*$ . I claim that  $x^*$  is a global minimum of  $f$ . First, since  $f^* = f(x')$  is a global minimum,  $f$  is bounded below and thus the decreasing sequence  $f(x_k)$  converges and hence is Cauchy. Therefore,

$$\lim_{k \rightarrow \infty} \|(\nabla f)_{x_k}\|^2 \leq 2\eta^{-1} \lim_{k \rightarrow \infty} |f(x_{k+1}) - f(x_k)| = 0$$

Therefore,

$$\|(\nabla f)_{x^*}\| \leq \|(\nabla f)_{x^*} - (\nabla f)_{x_k}\| + \|(\nabla f)_{x_k}\| \leq L\|x^* - x_k\| + \|(\nabla f)_{x_k}\|$$

The first term goes to zero on the subsequence  $x_{k_i}$  and the second term goes to zero and therefore  $(\nabla f)_{x^*} = 0$  so  $x^*$  is a global minimum of  $f$ . Therefore we can apply the previous lemma to conclude,

$$\|x_k - x^*\|^2 - \|x_{k+1} - x^*\|^2 \geq 2\eta(f(x_{k+1}) - f(x^*)) \geq 0$$

meaning that,

$$\|x_{k+1} - x^*\| \leq \|x_k - x^*\|$$

Therefore, since  $x_{k_i} \rightarrow x^*$  we see that  $x_k \rightarrow x^*$  converges. Finally, because  $f(x_k)$  is a decreasing sequence,

$$\begin{aligned} f(x_k) - f(x^*) &\leq \frac{1}{k} \sum_{j=1}^k [f(x_j) - f(x^*)] \leq \frac{1}{2\eta k} \sum_{j=1}^k (\|x_{j-1} - x^*\|^2 - \|x_j - x^*\|^2) \\ &= \frac{1}{2\eta k} (\|x_0 - x^*\|^2 - \|x_k - x^*\|^2) \leq \frac{\|x_0 - x^*\|^2}{2\eta k} \end{aligned}$$

□

## 26 Relationships between Geometric Categories

*Remark.* We consider the properties of the following morphisms of categories,

$$\mathbf{AffSch} \hookrightarrow \mathbf{Sch} \hookrightarrow \mathbf{LRS} \hookrightarrow \mathbf{RingSp} \rightarrow \mathbf{Top} \rightarrow \mathbf{Set}$$

## 27 $G$ -Structure on the Cotangent Bundle

Consider an action  $G \curvearrowright X$  over a base  $S$ .

The map  $a_G = (\text{id}, a) : G \times X \rightarrow G \times X$  defines a morphism  $\psi : a_G^* \Omega_{G \times X/S} \xrightarrow{\sim} \Omega_{G \times X/S}$ . Now we have,

$$\Omega_{G \times X/S} \cong \pi_1^* \Omega_{G/S} \oplus \pi_2^* \Omega_{X/S}$$

Furthermore,

$$a_G^* \Omega_{G \times X/S} \cong \pi_1^* \Omega_{G/S} \oplus a^* \Omega_{X/S}$$

Therefore  $\psi$  is a matrix,

$$\psi = \begin{pmatrix} \psi_{11} & \psi_{21} \\ \psi_{12} & \psi_{22} \end{pmatrix}$$

Furthermore,  $\pi_1 \circ a_G = \pi_1$  so  $\psi \circ a_G^* d\pi_1 = d\pi_1$  which is the inclusion  $\pi_1^* \Omega_{G/S} \rightarrow \Omega_{G \times X/S}$  and therefore  $\psi$  on the  $\pi_1^* \Omega_{G/S}$  factor is trivial meaning the matrix is of the form,

$$\psi = \begin{pmatrix} \text{id} & \psi_{21} \\ 0 & \psi_{22} \end{pmatrix}$$

therefore,

$$\psi_{22} : a^* \Omega_{X/S} \rightarrow \pi_2^* \Omega_{X/S}$$

is an isomorphism since the matrix is an isomorphism giving the required structure.

## 28 Vector Bundles on Vector Bundles

Given a vector bundle  $\pi : E \rightarrow X$  and a vector bundle  $\pi' : E' \rightarrow E$  is it true that  $\pi' : E' \rightarrow X$  in, in some sense, a vector bundle.

If  $E'$  is pulled back from a vector bundle  $V \rightarrow X$  then I think  $E'$  as an  $X$ -vector bundle is  $V \oplus E$ . Therefore if  $s : X \rightarrow E$  is the zero section, we should ask if  $E' = \pi^* s^* E$ .

In the topological category this is true because  $\pi : E \rightarrow X$  is a homotopy equivalence and indeed  $s \circ \pi$  is homotopic to the identity and therefore  $E' \cong \text{id}^* E' \cong \pi^* s^* E'$ .

However, this is false in the holomorphic/algebraic category. Indeed consider  $X = \mathbb{P}^1$  and the trivial bundle so  $E = \mathbb{P}^1 \times \mathbb{A}^1 \rightarrow \mathbb{P}^1$ . Then consider the vector bundle,

$$0 \longrightarrow \pi_1^* \mathcal{O}_{\mathbb{P}^1}(-1) \longrightarrow \mathcal{E} \longrightarrow \pi_1^* \mathcal{O}_{\mathbb{P}^1}(1) \longrightarrow 0$$

corresponding to the extension class,

$$\begin{aligned} \xi \in \text{Ext}_E^1(\pi_1^* \mathcal{O}_{\mathbb{P}^1}(-1), \pi_1^* \mathcal{O}_{\mathbb{P}^1}(1)) &= H^1(E, \pi_1^* \mathcal{O}_{\mathbb{P}^1}(-2)) = H^1(\mathbb{P}^1, (\pi_1)_* \pi_1^* \mathcal{O}_{\mathbb{P}^1}(-2)) \\ &= H^1(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(-2)) \otimes_k k[t] \cong k[t] \end{aligned}$$

which is given by  $\xi = t$ . Therefore, at  $t = 0$  we have the trivial extension so  $\mathcal{E}_t \cong \mathcal{O}(-1) \oplus \mathcal{O}(1)$  and for  $t \neq 1$  we have the nontrivial extension  $\mathcal{E}_t \cong \mathcal{O}^{\oplus 2}$ .

Therefore,  $\mathcal{E}$  is not the pullback of any vector bundle on  $\mathbb{P}^1$  otherwise its fibers would be constant. However, it is not clear if  $\mathcal{E}$  can be thought of as a vector bundle over  $\mathbb{P}^1$ . (WAIT)

## 29 Why the Stabilizers in the Moduli Space of Elliptic Curves are important

“The moduli space of elliptic curves is just  $\mathbb{A}^1$  because an elliptic curve is determined uniquely by its  $j$ -invariant”. This is wrong even over  $\mathbb{C}$  but it becomes clear if we try to work in a field that doesn't contain all square roots (or even  $\mathbb{Z}$ ). If  $k$  does not have all square roots then it is false that  $E$  is determined by  $j(E)$ . Indeed,

$$y^2 = x^3 + ax + b$$

and

$$y^2 \cdot d = x^3 + ax + b$$

where  $d \in k$  is a nonsquare are nonisomorphic curves which become isomorphic over  $k(\sqrt{d})$ . Indeed, we see that away from the special values  $j(E) \neq 0, 1728$  the elliptic curves with fixed  $j$ -invariant are classified by  $k^\times / (k^\times)^2$  and for  $j = 0, 1728$  they are classified by  $k^\times / (k^\times)^6$  and  $k^\times / (k^\times)^4$ .

Indeed, since these curves become isomorphic over  $\bar{k}$  they are all forms of some  $E_0$ . Then  $\text{Aut}(E_0) \cong \mu_2, \mu_6, \mu_4$  (note I think it really should be  $\mathbb{Z}/2\mathbb{Z}$  not  $\mu_2$  but these are the same in characteristic not 2 and indeed I am not sure what happens for  $p \leq 5$  anyway) as above and therefore isomorphism classes of curves over  $k$  with fixed  $j(E)$  are classified by,

$$H^1(k, \mu_n) \cong k^\times / (k^\times)^n$$

by Kummer theory.

What does this mean at the level of stacks. A morphism  $\mathrm{Spec}(k) \rightarrow \mathcal{M}_{1,1}$  landing in the topological point determined by  $j(E)$  factors through the residual Gerbe which is usually  $B(\mathbb{Z}/2\mathbb{Z})$  (at the special points it is  $B\mu_6$  or  $B\mu_4$ ) and therefore there are actually many such maps classified by  $(\mathbb{Z}/2\mathbb{Z})$ -bundles on  $k$  i.e. Galois 2-covers which are, via Kummer theory, classified by  $H^1(k, \mathbb{Z}/2\mathbb{Z}) = k^\times/(k^\times)^2$  and ditto for the special points. Therefore, we indeed see that the stabilizers of this stack carry useful arithmetic data.

## 30 $E$ -Fibrations

For simplicity assume that  $k$  has characteristic zero. I think everything here works in every characteristic except for possibly 2 and 3.

**Definition 30.0.1.** Let  $E$  be an elliptic curve over a field  $k$ . An  $E$ -fibration is a smooth proper morphism  $\mathcal{E} \rightarrow S$  with  $S$  over  $k$  such that each geometric fiber  $\mathcal{E}_K$  is isomorphic to  $E_K$  as a curve (not as a group scheme). We say that an  $E$ -fibration is trivial if it is isomorphic to  $E \times S$  as an  $S$ -scheme.

**Lemma 30.0.2.** If  $\mathcal{E} \rightarrow S$  has a section then it is trivial after a finite étale cover.

*Proof.* If  $\mathcal{E} \rightarrow S$  has a section then it is a family of elliptic curves over a base and therefore defines a morphism  $S \rightarrow \mathcal{M}_{1,1}$  whose image is topologically a single point. Since  $\mathcal{M}_{1,1}$  is a DM-stack this implies that there is a finite étale cover  $S' \rightarrow S$  such that  $S' \rightarrow \mathcal{M}_{1,1}$  is constant (the residual gerbe is finite étale) and thus  $\mathcal{E}' \rightarrow S'$  is trivial.  $\square$

**Proposition 30.0.3.** An  $E$ -fibration is locally trivial in the étale topology.

*Proof.* Because  $\mathcal{E} \rightarrow S$  is smooth, étale locally on  $S$  it has sections and hence after a further finite étale extension it is trivial.  $\square$

*Remark.* Here is a weird argument which works in some more general cases but only when  $S$  is regular. We can similarly define an  $F$ -fibration for any scheme  $F$  over a field  $k$ . Suppose that  $F$  is smooth and has smooth aut scheme  $\mathrm{Aut}(F)$ . Then if  $S$  is regular I claim that every  $F$ -fibration over  $S$  is locally trivial in the étale topology.

Let  $X \rightarrow S$  be an  $F$ -fibration with regular  $S$ . Then consider  $I = \mathrm{Isom}_S(X, S \times_k F)$  which is automatically an  $\mathrm{Aut}(F)$ -pseudo-torsor. I want to show that  $I$  is actually an étale torsor. The strategy is to use the fact that (representable) pseudotorsors for a smooth group scheme which are fppf are locally trivial in the étale topology. This is because using fppf descent and the pseudotorsor condition the morphisms is smooth and hence admits sections locally in the étale topology locally trivializing it as a torsor. Thus it suffices to show that  $I \rightarrow S$  is fppf. Because the geometric fibers of  $X \rightarrow S$  are all isomorphic to  $F_K$  the fibers of  $I \rightarrow S$  are isomorphic to  $\mathrm{Aut}(F_K)$  and hence have constant dimension. In particular,  $I \rightarrow S$  is surjective. (I DON'T THINK THE NEXT STEP WORKS) I want to use miracle flatness but  $I$  could be highly singular so ...

**Proposition 30.0.4.** Let  $S$  be a curve. Then every  $E$ -fibration over  $S$  is trivial after a finite (not necessarily étale) extension.

*Proof.*  $\mathcal{E} \times_S \mathcal{E} \rightarrow \mathcal{E}$  has a section so if we can produce a subscheme  $S' \subset \mathcal{E}$  which is finite over  $S$  then  $\mathcal{E}_{S'} \rightarrow S'$  admits a section and hence is trivial after an additional finite étale extension. There is an étale open  $U \rightarrow S$  with a section and hence a map  $U \rightarrow \mathcal{E}$  over  $S$ . Taking the scheme theoretic image gives a closed subscheme  $S' \subset \mathcal{E}$  such that  $S' \rightarrow S$  is nonconstant and hence finite since it is a map of curves.  $\square$

**Example 30.0.5.** Consider  $x \in E$  a nontorsion point. Then take  $E \times \mathbb{P}^1 \rightarrow \mathbb{P}^1$  and identify the fibers over  $\pm 1$  via translation by  $x$  to give  $X \rightarrow S$  where  $S$  is a nodal curve. This exists as a scheme by (REF) but is not projective (example on p.198 of Raynaud's thesis). In fact it is easy to see that  $X$  cannot be projective over  $S$ . Otherwise  $X$  would be projective and thus would admit a line bundle  $\mathcal{L}$ . Pulling back gives a line bundle  $\mathcal{L}'$  on  $E \times \mathbb{P}^1$  which takes the form  $\mathcal{L}_1 \otimes \mathcal{O}(n)$  for some  $n$  and  $\mathcal{L}_1 \in \text{Pic}(E)$ . However, the line bundle  $\mathcal{L}_1$  must be isomorphic to its pullback by  $\varphi : E \rightarrow E$  given by translation by  $x$ . Now,

$$\varphi^* \mathcal{O}_E([p_1] + \cdots + [p_n]) = \mathcal{O}_E([p_1 + x] + \cdots + [p_n + x]) = \mathcal{O}_E([p_1] + \cdots + [p_n] + n[x] - n[e])$$

and therefore we need that  $n[x] = n[e]$  meaning that  $nx = e$  so  $x$  must be a torsion point giving a contradiction unless  $\mathcal{L}_1 = \mathcal{O}_E$ . Therefore  $\mathcal{L}' \cong \mathcal{O}_{\mathbb{P}^1}(n)$  which is not ample. Since  $E \times \mathbb{P}^1 \rightarrow X$  is finite and  $\mathcal{L}$  is ample, its pullback  $\mathcal{L}'$  is ample giving a contradiction.

*Remark.* Since an  $E$ -fibration is étale locally trivial they are forms of  $E$  and therefore classified by  $H_{\text{ét}}^1(S, \text{Aut}(E))$ .

**Proposition 30.0.6.** An  $E$ -fibration is equivalent to the data of an  $E$ -torsor after a finite étale extension of the base.

*Proof.* Consider the sequence,

$$1 \longrightarrow E \longrightarrow \text{Aut}(E) \longrightarrow G \longrightarrow 1$$

I claim that  $G$  is a finite étale group scheme over  $k$ . (IS THIS TRUE!!) Therefore, there is an exact sequence,

$$H^1(S, E) \longrightarrow H^1(S, \text{Aut}(E)) \longrightarrow H^1(S, G)$$

Given a class  $[\mathcal{E} \rightarrow S] \in H^1(S, \text{Aut}(E))$  its image in  $H^1(S, G)$  is killed by a finite étale extension (since  $G$  is finite étale every  $G$ -torsor is finite étale and thus kills itself) hence we can assume that  $[\mathcal{E} \rightarrow S]$  is in the image of  $H^1(S, E) \rightarrow H^1(S, \text{Aut}(E))$  after a finite étale extension of the base.  $\square$

**Lemma 30.0.7.** If  $[\mathcal{E} \rightarrow S] \in H^1(S, E)$  is torsion then it is killed by a finite étale extension.

*Proof.* Consider the sequence,

$$0 \longrightarrow E[n] \longrightarrow E \xrightarrow{n} E \longrightarrow 0$$

where  $E[n]$  is a finite étale group scheme over  $k$  (IS THIS TRUE). Therefore we get a sequence,

$$H^1(S, E[n]) \longrightarrow H^1(S, E) \xrightarrow{n} H^1(S, E)$$

therefore if  $[\mathcal{E} \rightarrow S] \in H^1(S, E)$  is torsion it lies in some kernel and hence in the image of some  $[c'] \in H^1(S, E[n])$  which is killed by a finite étale cover because  $E[n]$  is a finite étale group scheme.  $\square$

**Proposition 30.0.8.** If  $S$  is regular then every  $E$ -fibration over  $S$  is projective and trivial in the finite étale topology.

*Proof.* By [Raynaud, Cor. XIII 2.4] every  $E$ -torsor is projective and torsion in  $H^1(S, E)$  so this follows from the previous results.  $\square$

(WANT AN EXAMPLE OF NORMAL SURFACE WITH LOC-TORSION TORSION WHICH IS NOT TORSION)

## 31 Review of Basics

*Remark.* The Zariski topology on  $\text{Spec}(A)$  makes the relations between  $I$  and  $Z$  definitional.

**Proposition 31.0.1.** For any  $Q \subset \text{Spec}(A)$ ,

$$\overline{Q} = V(\bigcap Q) = V(I(Q)) \quad \text{where} \quad I(Q) = \{f \in A \mid \forall \mathfrak{p} \in Q : f \in \mathfrak{p}\}$$

*Proof.* By definition, if  $\mathfrak{p} \in Q$  then  $I(Q) \subset \mathfrak{p}$  so  $\mathfrak{p} \in V(I(Q))$  meaning  $Q \subset V(I(Q))$ . Furthermore, if  $Q \subset V(J)$  for some ideal  $J$  then  $J \subset \mathfrak{p}$  for each  $\mathfrak{p} \in Q$  meaning,

$$J \subset \bigcap_{\mathfrak{p} \in Q} \mathfrak{p} = I(Q)$$

and thus  $V(I(Q)) \subset V(J)$  proving the claim.  $\square$

**Proposition 31.0.2.** for any  $S \subset A$  we have  $I(V(S)) = \sqrt{S}$  where,

$$\sqrt{S} = \bigcap_{\mathfrak{p} \supset S} \mathfrak{p}$$

*Proof.* This is by definition since  $V(S) = \{\mathfrak{p} \mid \mathfrak{p} \supset S\}$  and

$$I(V(S)) = \{f \in A \mid \forall \mathfrak{p} \in V(S) : f \in \mathfrak{p}\}$$

$\square$

**Proposition 31.0.3.** Let  $\varphi^* : \text{Spec}(B) \rightarrow \text{Spec}(A)$  be a map of affine schemes. Then,

$$(a) \quad (\varphi^*)^{-1}(D(f)) = D(\varphi(f))$$

$$(b) \quad (\varphi^*)^{-1}(V(I)) = V(\varphi(I))$$

$$(c) \quad \overline{\varphi^*(V(I))} = V(\varphi^{-1}(I))$$

*Proof.* First,

$$\varphi^*(\mathfrak{p}) \in D(f) \iff f \notin \varphi^*(\mathfrak{p}) \iff \varphi(f) \in \mathfrak{p} \iff \mathfrak{p} \in D(\varphi(f))$$

Second,

$$\varphi^*(\mathfrak{p}) \in V(I) \iff \varphi^*(\mathfrak{p}) \supset I \iff \mathfrak{p} \supset \varphi(I) \iff \mathfrak{p} \in V(\varphi(I))$$

For (c),

$$\overline{\varphi^*(V(I))} = V(\bigcap \varphi^*(V(I))) = V(\bigcap_{\mathfrak{p} \supset I} \varphi^{-1}(\mathfrak{p})) = V(\varphi^{-1}(\bigcap_{\mathfrak{p} \supset I} \mathfrak{p})) = V(\varphi^{-1}(\sqrt{I})) = V(\sqrt{\varphi^{-1}(I)}) = V(\varphi^{-1}(I))$$

where,

$$\varphi^{-1}(\sqrt{I}) = \sqrt{\varphi^{-1}(I)}$$

because,

$$x \in \varphi^{-1}(\sqrt{I}) \iff \varphi(x) \in \sqrt{I} \iff \varphi(x)^n \in I \iff \varphi(x^n) \in I \iff x^n \in \varphi^{-1}(I) \iff x \in \sqrt{\varphi^{-1}(I)}$$

$\square$

*Remark.* Dually, one might expect  $\varphi^*(D(f))^\circ = D(\varphi^{-1}(f))$  but this is false. Indeed, consider  $\varphi : \mathbb{Z} \rightarrow \mathbb{Q}$  then  $\varphi^*(D_{\mathbb{Q}}(2)) = \{(0)\}$  has empty interior but  $D_{\mathbb{Z}}(2)$  is nonempty.

*Remark.* Some of the sets on the RHS are not ideals but it is clear that for any subset  $S \subset A$  we have  $D(S) = D(SA)$  with  $SA$  the ideal generated by  $S$ .

*Remark.* For any multiplicative subset  $S \subset A$  we have  $\text{Spec}(S^{-1}A) \rightarrow \text{Spec}(A)$  identifies the subset where  $S \cap \mathfrak{p} = \emptyset$  in general this cannot be expressed as the condition  $I \not\subset \mathfrak{p}$  for some ideal  $I \subset A$  so this is not open. For example, let  $A = \mathbb{Z}$  and  $S = \mathbb{Z} \setminus \{0\}$  then  $S \cap \mathfrak{p} = \emptyset$  if and only if  $\mathfrak{p} = (0)$  but we need some  $I \neq (0)$  which is contained in every prime, impossible since  $\mathbb{Z}$  is reduced.

**Corollary 31.0.4.** If  $\varphi : A \rightarrow B$  satisfies  $\ker \varphi \subset \text{nilrad}(A)$  if and only if  $\varphi^* : \text{Spec}(B) \rightarrow \text{Spec}(A)$  is dominant.

*Proof.* Since  $\varphi^{-1}(0) = \ker \varphi$  we have,

$$\overline{\varphi^*(\text{Spec}(B))} = V(\ker \varphi)$$

and the result follows. Let's do this directly. First, let  $\ker \varphi \subset \text{nilrad}(A)$  we need to show any nonempty open  $U \subset \text{Spec}(A)$  contains some element of the image. Choose  $D(f) \subset U$  with  $f \notin \text{nilrad}(A)$  so  $D(f)$  is nonempty then  $\varphi(f) \notin \text{nilrad}(B)$  (see the remark) so  $D(\varphi(f))$  is nonempty and  $D(\varphi(f)) \rightarrow D(f)$  proving the claim. Conversely, if  $\varphi^*$  is dominant and  $f \in \ker \varphi$  then if  $D(f)$  is nonempty there is some  $\mathfrak{p} \in \text{Spec}(B)$  mapping into  $D(f)$  meaning  $\varphi(f) \notin \mathfrak{p}$  but  $\varphi(f) = 0$  giving a contradiction so  $D(f)$  is empty meaning  $f \in \text{nilrad}(A)$ .  $\square$

*Remark.* Notice that,

$$\varphi^{-1}(\text{nilrad}(B)) \subset \text{nilrad}(A) \iff \ker \varphi \subset \text{nilrad}(A)$$

The forward implication is obvious. Conversely, if  $\ker \varphi \subset \text{nilrad}(A)$  then if  $\varphi(x) \in \text{nilrad}(B)$  then  $\varphi(x)^n = 0$  so  $\varphi(x^n) = 0$  and hence  $x^n \in \ker \varphi \subset \text{nilrad}(A)$  so  $x \in \text{nilrad}(A)$  proving the claim.

## 32 Factoring Through a Point

**Proposition 32.0.1.** Let  $f : X \rightarrow Y$  be a morphism of schemes with  $f(X) = \{y\}$  one point. Then there is a factorization,

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ \uparrow & & \uparrow \\ X_{\text{red}} & \longrightarrow & \text{Spec}(\kappa(y)) \end{array}$$

*Proof.* It suffices to show that  $X_{\text{red}} \rightarrow Y$  factors through  $\text{Spec}(\kappa(y))$ . Since  $f(X) = \{y\}$  reduce to affine opens  $U \rightarrow V$  is  $\varphi^* : \text{Spec}(B) \rightarrow \text{Spec}(A)$  image is  $\mathfrak{p} \subset A$ . Thus  $\mathfrak{p} = \varphi^{-1}(\mathfrak{q})$  for all primes  $\mathfrak{q} \subset B$ . Thus,

$$\mathfrak{p} = \varphi^{-1}(\text{nilrad}(B))$$

and also if  $f \notin \mathfrak{p}$  then  $\varphi(f) \notin \mathfrak{q}$  so  $\varphi(f)$  is a unit. Therefore we get a factorization,

$$A \rightarrow (A/\mathfrak{p})_{\mathfrak{p}} \rightarrow B/\text{nilrad}(B) = B_{\text{red}}$$

$\square$



### 33 The Stack of All Curves is Not Separated

*Remark.* What do we mean by separated for a map of stacks? First of all, for a map of algebraic stacks  $f : \mathcal{X} \rightarrow \mathcal{Y}$  the diagonal  $\Delta_f$  is always representable by algebraic spaces ([Tag 04XS](#)) so we can talk about its properties as long as we know about the property for algebraic spaces. It might seem natural to impose that the diagonal is a closed embedding but it turns out this is too restrictive, in particular this will imply that a separated DM-stack is an algebraic space (I THINK) for the following reason.

**Proposition 33.0.1.** The following is a 2-pullback diagram,

$$\begin{array}{ccc} \mathrm{Isom}_{\mathcal{X}}(a, b) & \longrightarrow & T \\ \downarrow & \lrcorner & \downarrow (ab) \\ \mathcal{X} & \xrightarrow{\Delta} & \mathcal{X} \times \mathcal{X} \end{array}$$

*Proof.* □

*Remark.* In particular, if  $\Delta$  is a closed immersion then we require that  $\mathrm{Isom}_{\mathcal{X}}(a, b) \rightarrow T$  is a closed immersion. Consider the case that  $T = \mathrm{Spec}(k)$  then this says that all objects in  $\mathcal{X}$  have only trivial automorphisms. In particular, if  $\mathcal{X}$  is an algebraic-stack then it is an algebraic space ([Tag 04SZ](#)). This is clearly not satisfactory.

*Remark.* One way to fix this problem is to notice the following: if  $f : X \rightarrow Y$  is a map of schemes then because  $\Delta_{X/Y} : X \rightarrow X \times_Y X$  is an immersion always the following are equivalent,

- (a)  $f$  is separated
- (b)  $\Delta_{X/Y}$  is a closed immersion
- (c)  $\Delta_{X/Y}$  is finite
- (d)  $\Delta_{X/Y}$  is proper
- (e)  $\Delta_{X/Y}$  is universally closed
- (f)  $\Delta_{X/Y}$  is closed.

Therefore, we could take any of these as a definition for stacks and retain the same notion for schemes (and algebraic spaces by similar reasoning). Because topological conditions behave somewhat badly for stacks, finite and proper are the convenient notions. Following the stacks project we take the latter although some authors prefer finiteness of the diagonal.

**Definition 33.0.2.** A map of algebraic stacks  $f : \mathcal{X} \rightarrow \mathcal{Y}$  is,

- (a) *separated* if  $\Delta_f : \mathcal{X} \rightarrow \mathcal{X} \times_{\mathcal{Y}} \mathcal{X}$  is proper
- (b) *quasi-separated* if  $\Delta_f : \mathcal{X} \rightarrow \mathcal{X} \times_{\mathcal{Y}} \mathcal{X}$  is quasi-compact and quasi-separated.

where properties of  $\Delta_f$  are in the sense of representable maps.

*Remark.* There are two ways to know we have the “correct” definition of separatedness. The first is that, when the map  $f$  is representable by algebraic spaces, this definition agrees with the separated in the sense of representable maps.

**Lemma 33.0.3.** Let  $f : \mathcal{X} \rightarrow \mathcal{Y}$  be a representable (by algebraic spaces) morphism of algebraic stacks. Then the following are equivalent,

- (a)  $f$  is separated in the sense of representable maps
- (b)  $\Delta_f$  is a cloed immersion
- (c)  $\Delta_f$  is proper
- (d)  $\Delta_f$  is universally closed.

*Proof.* This is basically because these properties are the same for maps of algebraic spaces and are preserved under base change. See [Tag 04YS](#) for details.  $\square$

*Remark.* Part of the reason we require  $\Delta_f$  to be proper and not just universally closed is that although  $\Delta_f$  is automatically finite type it can be nonseparated. For example, let  $G$  be a nonseparated group algebraic space over  $S = \operatorname{Spec}(k)$  (see [Tag 06E9](#)) and consider  $\mathcal{X} = BG = [S/G] \rightarrow S$ . Then consider,

$$\begin{array}{ccc} G & \longrightarrow & S \\ \downarrow & \lrcorner & \downarrow \\ BG & \xrightarrow{\Delta} & BG \times_S BG \end{array}$$

shows that  $\Delta$  is not separated.

*Remark.* The best reason we know that we have the right notion of separatedness is that the following valuative criterion holds.

**Proposition 33.0.4.** Let  $f : \mathcal{X} \rightarrow \mathcal{Y}$  be a quasi-separated morphism of algebraic stacks. If  $f$  satisfies the uniqueness part of the valuative criterion then  $f$  is separated where this criterion is for every 2-commutative diagram,

$$\begin{array}{ccc} \operatorname{Spec}(K) & \longrightarrow & \mathcal{X} \\ \downarrow j & \nearrow \text{dashed} & \downarrow f \\ \operatorname{Spec}(A) & \longrightarrow & \mathcal{Y} \end{array}$$

with  $A$  a valuation ring and  $K = \operatorname{Frac}(A)$  the category of maps  $\operatorname{Spec}(A) \rightarrow \mathcal{X}$  equipped with natural transformations making the diagrams 2-commute is equivalent to a  $(-1)$ -category (either empty or one element).

*Remark.* See [Tag 0CL9](#) we define what this uniqueness criterion means. The category of dotted arrows has objects  $(a, \alpha, \beta)$  where  $a : \operatorname{Spec}(A) \rightarrow \mathcal{X}$  is a morphism and  $\alpha : a \circ j \rightarrow x$  and  $\beta : y \rightarrow f \circ a$  are two morphisms witnessing the commutativity of the triangles. These must be compatible in the sense that,

$$\begin{array}{ccc} & f \circ a \circ j & \\ \beta * \operatorname{id}_j \nearrow & & \searrow \operatorname{id}_f * a \\ y \circ j & \xrightarrow{\gamma} & f \circ x \end{array}$$

commutes. A morphism  $(a, \alpha, \beta) \rightarrow (a', \alpha', \beta')$  is a 2-arrow  $\theta : a \rightarrow a'$  compatible with commutativity meaning  $\alpha = \alpha' \circ (\theta * \operatorname{id}_j)$  and  $\beta = (\operatorname{id}_f * \theta) \circ \beta$ .

### 33.1 The Stack of All Curves

*Remark.* Work in the category of  $k$ -schemes so all our stacks are equipped with a forgetful map to  $\mathrm{Spec}(k)$ .

**Definition 33.1.1.** The stack of all curves  $\mathcal{M}^{\mathrm{all}}$  is the fibered category of flat proper finitely presented morphism of algebraic spaces  $\pi : \mathcal{C} \rightarrow S$  whose geometric fibers are 1-dimensional.

**Example 33.1.2.** Consider  $A = k[[t]]$  and  $K = \mathrm{Frac}(A) = k((t))$ . Then consider,

$$X = \mathrm{Proj} \left( A[x, y, z] / (zy^2 - x^3 - t^4xz^2) \right) \rightarrow \mathrm{Spec}(A)$$

which is flat because it is an integral scheme over a DVR and hence torsion-free as an  $A$ -module. The special fiber,

$$X_k \cong \mathrm{Proj} \left( k[x, y, z] / (zy^2 - x^3) \right)$$

is a nodal curve while the generic fiber,

$$X_k \cong \mathrm{Proj} \left( K[x, y, z] / (zy^2 - x^3 - t^4xz^2) \right) \cong \mathrm{Proj} \left( K[x, y, z] / (zy^2 - x^3 - xz^2) \right) \cong E_K$$

where the isomorphism takes,

$$x \mapsto t^2x \quad y \mapsto t^3y \quad z \mapsto z$$

is an elliptic curve base changed from the elliptic curve over  $k$ ,

$$E = \mathrm{Proj} \left( k[x, y, z] / (zy^2 - x^3 - xz^2) \right)$$

Therefore, this shows that  $\mathcal{M}^{\mathrm{all}}$  is not separated because it violates the valuative criterion,

$$\begin{array}{ccc} \mathrm{Spec}(K) & \xrightarrow{x} & \mathcal{M}^{\mathrm{all}} \\ j \downarrow & \nearrow & \downarrow f \\ \mathrm{Spec}(A) & \xrightarrow{y} & \mathrm{Spec}(k) \end{array}$$

Such a diagram corresponds to a curve  $C \rightarrow \mathrm{Spec}(K)$  and because  $\mathrm{Spec}(k)$  is a scheme there are no nontrivial 2-morphisms so the square is commutative with a unique 2-morphism  $\gamma = \mathrm{id}$ . Lifts are tripples  $(a, \alpha, \beta)$  where  $a$  corresponds to a curve  $\mathcal{C} \rightarrow \mathrm{Spec}(A)$  and  $\alpha : a \circ j \rightarrow x$  is an isomorphism  $\alpha : \mathcal{C}_K \rightarrow C$  over  $K$  and  $\beta : y \rightarrow f \circ x$  must be the identity because  $\mathrm{Spec}(k)$  is representable. Finally these much satisfy,

$$\gamma = (\mathrm{id}_f * \alpha) \circ (\beta * \mathrm{id}_j)$$

which is automatic because these are 2-morphisms on  $\mathrm{Spec}(k)$  (also  $\gamma = \mathrm{id}$  and  $\beta = \mathrm{id}$  and  $\mathrm{id}_f * \alpha = \mathrm{id}$  since  $\mathcal{M}^{\mathrm{all}} \rightarrow \mathrm{Spec}(k)$  collapses all isomorphisms to  $\mathrm{id}$ ). Therefore, the category of arrows is equivalent to the category of curves  $\mathcal{C} \rightarrow \mathrm{Spec}(A)$  equipped with an isomorphism  $\alpha : \mathcal{C}_K \rightarrow C$  with morphisms as isomorphisms of families respecting the identification of the special fiber. Therefore, this is exactly the category of models. However, above we have  $X$  and  $E_A$  which are nonisomorphic models (have nonisomorphic generic fibers) so  $\mathcal{M}^{\mathrm{all}}$  is not separated.

Here is another way to understand the failure of separatedness in this example. The models  $X, E_A$  determine morphisms  $a, b : \mathrm{Spec}(A) \rightarrow \mathcal{M}^{\mathrm{all}}$  so consider,

$$\begin{array}{ccc}
\mathrm{Isom}_A(X, E_A) & \longrightarrow & \mathrm{Spec}(A) \\
\downarrow & \lrcorner & \downarrow \\
\mathcal{M}^{\mathrm{all}} & \longrightarrow & \mathcal{M}^{\mathrm{all}} \times_k \mathcal{M}^{\mathrm{all}}
\end{array}$$

and we know  $\mathrm{Isom}_A(X, E_A)$  is nonempty over the generic fiber but empty on the special fiber so its image in  $\mathrm{Spec}(A)$  is not closed and hence  $\mathrm{Isom}_A(X, E_A) \rightarrow \mathrm{Spec}(A)$  is not proper proving that  $\Delta_f$  is not proper.

*Remark.* The semi-stable reduction theorem tells us that a semi-stable reduction when it exists is always unique and also after a finite extension does exist. This proves the uniqueness and existence parts of the valuative criteria for properness for the stack of semi-stable curves.

## 34 Moduli of Smooth Fanos

**Definition 34.0.1.** A family of smooth fanos is a smooth proper finitely presented morphism  $\pi : X \rightarrow S$  such that each fiber  $X_s$  is a smooth Fano meaning a smooth projective variety with ample  $\omega_{X_s}^\vee$ .

## 35 Isotrivial Families

**Definition 35.0.1.** A *polarized family* if a proper, flat, finitely presented morphism  $\pi : X \rightarrow S$  equipped with a relatively ample invertible sheaf  $\mathcal{L}$  on  $X$ . A morphism of polarized families is a cartesian diagram,

$$\begin{array}{ccc}
X' & \longrightarrow & X \\
\downarrow f & & \downarrow \\
S' & \xrightarrow{g} & S
\end{array}$$

and an isomorphism  $\varphi : f^*\mathcal{L} \rightarrow \mathcal{L}'$ .

*Remark.* Since  $\pi : X \rightarrow T$  is finite type and  $\mathcal{L}$  is  $\pi$ -relatively ample there is a (Zariski) open cover  $T_i \rightarrow T$  such that  $\mathcal{L}$  is ample for  $X_i \rightarrow T_i$  and thus we get a closed embedding  $X_i \hookrightarrow \mathbb{P}_{T_i}^N$  over  $T_i$  defined by  $\mathcal{L}^{\otimes n_i}$  for some  $n_i > 0$ . Therefore,  $\pi$  is locally projective (in the sense of Hartshorne).

**Definition 35.0.2.** The stack of polarized proper schemes  $\mathcal{M}_{\mathrm{pol}}$  is the stack of polarized families. Explicitly, it is the category fibered over  $(\mathbf{Sch}_{\mathbb{Z}})_{\mathrm{fppf}}$  whose objects are pairs  $(X \rightarrow S, \mathcal{L})$  with,

- (a)  $X \rightarrow S$  a proper, flat, finitely presented morphism
- (b)  $\mathcal{L}$  an invertible  $\mathcal{O}_X$ -module relatively ample for  $X \rightarrow S$

and morphisms  $(X' \rightarrow S', \mathcal{L}') \rightarrow (X \rightarrow S, \mathcal{L})$  are given by  $(f, g, \varphi)$  with,

- (a)  $f : X' \rightarrow X$  and  $g : S' \rightarrow S$  morphisms of schemes such that,

$$\begin{array}{ccc}
X' & \xrightarrow{f} & X \\
\downarrow & & \downarrow \\
S' & \xrightarrow{g} & S
\end{array}$$

is a commutative cartesian diagram,

(b)  $\varphi : f^* \mathcal{L} \rightarrow \mathcal{L}'$  is an isomorphism.

**Theorem 35.0.3.** The fibered category  $\mathcal{M}_{\text{pol}}$  is a locally noetherian algebraic stack and the canonical morphism  $\mathcal{M}_{\text{pol}} \rightarrow \text{Spec}(\mathbb{Z})$  is quasi-separated and locally of finite presentation.

*Proof.* See [Tag 0D4X](#) and [Tag 0DPU](#). Then  $\mathcal{M}_{\text{pol}}$  is locally noetherian because  $\text{Spec}(\mathbb{Z})$  is and we apply [Tag 06R6](#).

DO THE IDEA!!

□

**Lemma 35.0.4.** The morphism  $\mathcal{I}_{\mathcal{M}_{\text{pol}}} \rightarrow \mathcal{M}_{\text{pol}}$  is quasi-compact.

*Proof.* For each morphism  $T \rightarrow \mathcal{M}_{\text{pol}}$  from a scheme defining a polarized family  $(X \rightarrow T, \mathcal{L})$  we get the 2-fiber square,

$$\begin{array}{ccc} \text{Aut}_{\mathcal{M}_{\text{pol}}}(X) & \longrightarrow & T \\ \downarrow & & \downarrow \\ \mathcal{I}_{\mathcal{M}_{\text{pol}}} & \longrightarrow & \mathcal{M}_{\text{pol}} \end{array}$$

therefore it suffices to show that  $\text{Aut}_{\mathcal{M}_{\text{pol}}}(X) \rightarrow T$  is quasi-compact. Since quasi-compactness is local on the base, we may assume that  $\pi : X \rightarrow T$  is projective with  $X \hookrightarrow \mathbb{P}_T^n$  via  $\mathcal{L}$ . Since polarized automorphisms of  $X$  fix  $\mathcal{L}$  we see that any automorphism of  $X$  extends to an automorphism of  $\mathbb{P}_T^n$  giving a map of sheaves,

$$\text{Aut}_{\mathcal{M}_{\text{pol}}}(X) \rightarrow \text{PGL}_{n+1}$$

whose kernel is given by automorphisms of  $X$  which fix  $\mathbb{P}_T^n$  and hence of the form  $(\text{id}, \text{id}, \varphi)$  where  $\varphi : \mathcal{L} \rightarrow \mathcal{L}$  is an automorphism. Therefore, we get a sequence,

$$1 \longrightarrow \mathbb{G}_m \longrightarrow \text{Aut}_{\mathcal{M}_{\text{pol}}}(X) \longrightarrow \text{PGL}_{n+1}$$

(DO TRANSPORTERS EXIST IN GENRAL?)

□

*Remark.* This is quite special to automorphisms of polarized varieties. For example,  $A = E \times E$  where  $E$  is an ordinary elliptic curve has  $\text{Aut}(A) = \text{GL}_2(\mathbb{Z})$  which is not quasi-compact but  $A$  has finitely many polarized automorphisms.

**Lemma 35.0.5.** Let  $\mathcal{X}$  be a locally noetherian algebraic stack then for each  $x \in |X|$  the residual gerbe  $\mathcal{Z}_x$  of  $\mathcal{X}$  at  $x$  exists and  $\mathcal{Z}_x \rightarrow \mathcal{X}$  is a closed embedding.

*Proof.*

□

**Corollary 35.0.6.** For each  $x \in |\mathcal{M}_{\text{pol}}|$  the residual gerbe  $\mathcal{Z}_x$  of  $\mathcal{M}_{\text{pol}}$  at  $x$  exists and  $\mathcal{Z}_x \rightarrow \mathcal{M}_{\text{pol}}$  is a closed embedding.

*Proof.* Using lemma [Tag 06UH](#).

DO THE IDEA

□

### 35.1 Using the Gerbes

**Lemma 35.1.1.** Let  $\mathcal{X}$  be a reduced algebraic stack and  $\mathcal{Y}$  be a locally noetherian algebraic stack with  $y \in |\mathcal{Y}|$  is a closed point. Let  $f : \mathcal{X} \rightarrow \mathcal{Y}$  be a morphism of stacks such that  $f(|\mathcal{X}|) = \{y\}$  then there is a factorization,

$$\begin{array}{ccc} \mathcal{X} & \xrightarrow{f} & \mathcal{Y} \\ & \searrow & \nearrow \\ & \mathbb{Z}_y & \end{array}$$

*Proof.* Because  $\mathbb{Z}_y \hookrightarrow \mathcal{Y}$  is a closed substack with  $|\mathbb{Z}_y| = \{y\}$ , we can apply [Tag 050B](#).  $\square$

**Proposition 35.1.2.** Let  $f : T \rightarrow \mathcal{X}$  be a morphism from a reduced noetherian Jacobson scheme to a locally noetherian algebraic stack

### 35.2 WORK TO DO

Here is a problem, because of the extra  $\mathbb{G}_m$  factor the DM-locus is empty. We would like to consider the DM-locus inside here and see the stack of curves living inside or something. Can we rigidify this stack somehow. What sort of level structure will kill these automorphisms of the line bundle.

## 36 Is Syntomic the topology that was promised?

I want the smallest topology containing  $\mathrm{Spec}(A[x]/(x^p - a)) \rightarrow \mathrm{Spec}(A)$  in characteristic  $p$  and Zariski covers. I would also be interested in the “finite” version of this topology.

Perhaps there is a good notion of a “finite flat topology”.

## 37 Connections on Principle Bundles

**Proposition 37.0.1.** Let  $\pi : G \rightarrow S$  be a group scheme. Then,

$$\Omega_{G/S} = \pi^* e^* \Omega_{G/S}$$

so if we set,

$$\omega_{G/S} = e^* \Omega_{G/S}$$

then,

$$\Omega_{G/S} = \pi^* \omega_{G/S}$$

Furthermore, by the projection form,

$$\pi_* \Omega_{G/S} = \omega_{G/S} \otimes_{\mathcal{O}_S} \mathcal{O}_G$$

and thus if  $\pi$  is  $\mathcal{O}$ -connected (e.g. for  $G/S$  an abelian scheme) then,

$$\pi_* \Omega_{G/S} = \omega_{G/S}$$

*Proof.* Consider the Cartesian diagram,

$$\begin{array}{ccccc}
G \times_S G & & \xrightarrow{m} & & G \\
& \searrow \text{dashed} & & \searrow \pi_2 & \\
& G \times_S G & & G & \\
& \downarrow \pi_1 & & \downarrow & \\
& G & \xrightarrow{\quad} & S &
\end{array}$$

(Note: A curved arrow labeled  $\pi_1$  also goes from  $G \times_S G$  to  $G$  in the bottom row.)

because the dashed arrow is an isomorphism, the outside square is Cartesian so  $m^*\Omega_{G/S} = \pi_1^*\Omega_{G/S}$ . Then,

$$\pi^*e^*\Omega_{G/S} = (e \circ \pi, \text{id})^*\pi_1^*\Omega_{G/S} = (e \circ \pi, \text{id})^*m^*\Omega_{G/S} = \text{id}^*\Omega_{G/S} = \Omega_{G/S}$$

□

*Remark.* If  $S = \text{Spec}(k)$  then  $\omega_{G/S} = \mathfrak{g}$  is the Lie algebra.

**Definition 37.0.2.** Let  $P$  be an object on an  $S$ -scheme  $\pi : X \rightarrow S$ . Then an  $S$ -connection on  $P$  is an isomorphism  $\varphi : \pi_1^*P \rightarrow \pi_2^*P$  of objects over  $X^{(1)}$  such that  $\Delta^*\varphi = \text{id}$  where,

$$X^{(1)} \hookrightarrow X \times_S X$$

is the first infinitesimal diagonal. Consider,

$$X \hookrightarrow X \times_S X \times_S X$$

and the first-order neighborhood  $X_3^{(1)}$  equipped with three projections  $\pi_{ij} : X_3^{(1)} \rightarrow X^{(1)}$ . We say that  $\varphi$  is *integrable* if it satisfies the cocycle condition,

$$\pi_{23}^*\varphi \circ \pi_{12}^*\varphi = \pi_{13}^*\varphi$$

**Proposition 37.0.3.** Let  $G \rightarrow S$  be a smooth group scheme and  $\pi : P \rightarrow X$  be a  $G$ -bundle. Consider the sequence,

$$0 \longrightarrow \pi^*\Omega_{X/S} \longrightarrow \Omega_{P/S} \longrightarrow \Omega_{P/X} \longrightarrow 0$$

This is an exact sequence of descent data and therefore arises as  $\pi^*$  of an exact sequence,

$$0 \longrightarrow \Omega_X \longrightarrow Q \longrightarrow \text{ad}(P) \longrightarrow 0$$

where  $\text{ad}(P)$  is the adjoint bundle,

$$\text{ad}(P) = P \times_G \omega_{G/S}$$

defined via the adjoint action of  $G$  on  $\omega_{G/S}$ .

## 38 When are Flag Varieties Toric?

Let  $G$  be a simple reductive group and  $P \subset G$  a parabolic subgroup. When is the flag variety  $G/P$  toric? We know that any algebraic group is unirational (at least if  $G$  is reductive or  $k$  is perfect see Lemma 7.2.3 in Brian's second course) and thus  $G/P$  is unirational so it might seem reasonable that it could be toric.

**Proposition 38.0.1.** If  $G$  is simple with trivial center and  $P \subset G$  is a parabolic subgroup then  $\text{Aut}(G/P) = G \rtimes A$  where  $A$  is a finite group determined by the automorphisms of the Dynkin diagram of  $G$  except for the following exceptional cases,

- (a)  $X = G_2/U_2$  where  $\text{Aut}^0(X) = \text{SO}_7$
- (b)  $X = \text{Sp}_r/\text{Sp}_{r-1}U_1$  where  $\text{Aut}^0(X) = \text{PSL}_{2r}$
- (c)  $X = \text{SO}_{2n+1}/U_n$  where  $\text{Aut}^0(X) = \text{PSO}_{2n+2}$ .

*Proof.* There is a reference [here](#). □

Therefore, except for these cases the maximal torus of  $\text{Aut}(X)$  is the maximal torus  $T \subset G$ . Hence if  $\dim T < \dim G/P$  then it is impossible for  $G/P$  to be toric. For example, let  $G = \text{PGL}_n$  and  $B \subset G$  the standard Borel so that  $G/B$  is the complete flag variety. Then  $\dim T = n - 1$  and  $\dim G/B = \frac{1}{2}n(n - 1)$  so we see that,

$$\dim T < \dim G/B \iff n > 2$$

and indeed for  $n = 2$  we get  $G/B = \mathbb{P}^1$  which is toric.

This proves that projective bundles over toric varieties need not be toric. Indeed,  $F_n = \text{PGL}_n/B$  is an iterated projective bundle over  $\text{Spec}(k)$  and hence at some point a projective bundle must take a toric variety to a non-toric variety since  $F_n$  is not toric for  $n > 2$ . For  $n = 3$  we get a counter-example for the projectivization of a rank 2 vector bundle on  $\mathbb{P}^2$  in fact  $F_3 = \mathbb{P}_{\mathbb{P}^2}(\mathcal{Q})$  where  $\mathcal{Q} = \Omega_{\mathbb{P}^2}(1)$  is the canonical subbundle and  $\mathbb{P}^2$  is toric but  $F_3$  is not toric by the above discussion.

## 39 May 27 Orders of Points on genus 1 curves

**Definition 39.0.1.** Let  $X \rightarrow \text{Spec}(k)$  be a  $k$ -scheme. For a point  $x \in X$ , let  $\deg(x) = [\kappa(x) : k]$  and,

- (a)  $\text{radix } X = \min\{\deg x \mid x \in X\}$
- (b)  $\text{ind } X = \gcd\{\deg x \mid x \in X\}$ .

**Proposition 39.0.2.** Let  $G$  be a  $k$ -group and  $T$  a  $G$ -torsor. Then  $\text{ind } T$  is the gcd of the degrees of all extensions such that  $T$  becomes trivial.

*Proof.*  $T_{k'}$  is trivial iff  $T(k') \neq \emptyset$  proving the claim. □

**Proposition 39.0.3.** If  $T \in H^1(k, G)$  is torsion, denote its order per  $T$ , then  $\text{per } T \mid \text{ind } T$ .

*Proof.* It suffices to show that if  $T_{k'}$  is trivial then  $\text{per } T \mid [k' : k]$ . Indeed, using that,

$$H^1(k, G) \xrightarrow{\text{res}} H^1(k', G) \xrightarrow{\text{cor}} H^1(k, G)$$

is multiplication by  $n = [k' : k]$  we see that  $T \in H^1(k, G)[n]$  and hence its order per  $T \mid n$ . □

**Proposition 39.0.4.** Let  $C$  be a genus 1 curve over a field  $k$ . Then  $\forall x \in C : \text{ind}(C) \mid \deg x$  so in particular  $\text{radix } C = \text{ind } C$ .



*Proof.* Let  $x \in C$  be a point achieving the minimum and  $y \in C$  another point. Assume the theorem is false then  $\gcd(\deg x, \deg y) < \deg x$ . Therefore, there is a divisor,

$$D = p[x] + q[y]$$

for  $p, q \in \mathbb{Z}$  such that  $0 < \deg D < \deg x$ . Then by Riemann-Roch,

$$h^0(C, \mathcal{O}_C(D)) = \deg D > 0$$

meaning that  $D$  is equivalent to an effective divisor proving there is some point with smaller degree than  $x$ .  $\square$

**Proposition 39.0.5.** Let  $k$  be a field and  $E$  an elliptic curve over  $k$  such that for every finite extension  $k'/k$  and all sufficiently large primes  $p \gg 0$  the abelian group  $k'^\times/(k'^\times)^n$  is infinite and  $E(k)/pE(k)$  is finite (e.g.  $k$  is a number field and  $E$  is any elliptic curve). Then for any  $n > 0$  there is an  $E$ -torsor  $C$  with  $\text{ind } C \geq n$ .

*Proof.* It suffices to prove that for any sufficiently large prime  $\ell$  there is a nontrivial  $C \in H^1(k, E)[\ell]$  since then  $\text{per } C = \ell$  and thus  $\text{ind } C \geq \ell$ . Choose  $\ell$  large enough to be invertible in  $k$  (i.e. not the characteristic) then consider the sequence,

$$0 \longrightarrow E[n] \longrightarrow E \xrightarrow{n} E \longrightarrow 0$$

exact in the étale topology. Thus we find an exact sequence,

$$0 \longrightarrow E(k)/\ell E(k) \longrightarrow H^1(k, E[\ell]) \longrightarrow H^1(k, E)[\ell] \longrightarrow 0$$

Since the first term is finite, to show that  $H^1(k, E)[\ell] \neq 0$  it suffices to show that  $H^1(k, E[\ell])$  is infinite. Let  $k'/k$  be a field extension such that  $E[\ell]$  is split and  $k'$  has all  $\ell^{\text{th}}$ -roots of unity. Then from the inflation-restriction sequence,

$$0 \longrightarrow H^1(\text{Gal}(k'/k), E[\ell]^{G_{k'}}) \longrightarrow H^1(k, E[\ell]) \longrightarrow H^1(k', E[\ell]) \longrightarrow H^2(\text{Gal}(k'/k), E[\ell]^{G_{k'}})$$

However,  $\text{Gal}(k'/k)$  is finite and  $E[\ell]^{G_{k'}}$  is a finite abelian group (equipped with the trivial action) so  $H^1(\text{Gal}(k'/k), E[\ell]^{G_{k'}})$  and  $H^2(\text{Gal}(k'/k), E[\ell]^{G_{k'}})$  are finite. Therefore,  $H^1(k, E[\ell])$  is infinite iff  $H^1(k', E[\ell]) = \text{Hom}(G_{k'}, E[\ell])$  is infinite. However,  $E[\ell]_{k'} \cong (\mathbb{Z}/\ell\mathbb{Z})^2 \cong \mu_\ell^2$  and therefore from the Kummer sequence,

$$H^1(k', \mu_\ell) = (k'^\times)/(k'^\times)^\ell$$

is infinite proving that  $H^1(k', E[\ell])$  is infinite.  $\square$

*Remark.* Can we strengthen the statement to  $\text{ind } C = n$ ?

## 40 TEST

## 41 Introduction

Let  $X$  be a minimal variety of dimension  $n$  over  $\mathcal{C}$ . Throughout this note, we will assume that there are  $k$  linearly independent nowhere-vanishing holomorphic 1-forms on  $X$ . In other words, there exists a short exact sequence

$$0 \rightarrow \mathcal{O}^{\oplus k} \rightarrow \Omega_X^1 \rightarrow \mathcal{F}^\vee \rightarrow 0$$

of vector bundles.

**Lemma 41.0.1.** The dual short exact sequence

$$0 \rightarrow \mathcal{F} \rightarrow T_X \rightarrow \mathcal{O}_X^{\oplus k} \rightarrow 0$$

gives a foliation.

**Q1.** When is  $\mathcal{F}$  algebraic?

This is very much related to a paper of Campana and Peternell, where they ask this question in the context

## 41.1 Foliations

**Definition 41.1.1.** A foliation  $\mathcal{F} \subset T_X$  is *regular* if it is a subbundle.

Recall,

**Theorem 41.1.2** (Frobenius). Let  $M$  be a smooth manifold and  $\mathcal{F}$  a regular foliation. Then there is a collection  $\{L_\alpha\}_\alpha$  of connected injectively immersed smooth manifolds  $L_\alpha \rightarrow M$  (not closed) such that,

- (a)  $M$  is a disjoint union of the  $L_\alpha$
- (b) for each  $p \in M$  there is a chart  $U_p \subset M$  such that  $U_p \cap L_\alpha$  is a countable union of slices (there are coordinates  $(x^1, \dots, x^n)$  such that the components are  $x^{r+1} = c_1, \dots, x^n = c_{n-r}$ )
- (c) for each  $p \in L_\alpha$  we have  $T_p L_\alpha = \mathcal{F}_p$  inside  $T_M$ .

If  $X$  is a complex manifold and  $\mathcal{F} \subset T_X$  is a complex regular foliation then we can assume that the leaves  $L_\alpha$  are immersed complex submanifolds.

**Definition 41.1.3.** Let  $X$  be a smooth algebraic variety. We say an immersed manifold  $\iota : L \subset X$  is *algebraic* if  $\iota(L) \cap Z^{\text{sm}} \subset Z$  is (analytically) open where  $Z$  is the Zariski closure of  $\iota(L)$ .

*Remark.* I choose this slightly strange condition to capture the following phenomenon. Let  $\mathbb{A}^1 \rightarrow \mathbb{A}^2$  be  $t \mapsto (t^2 - 1, t(t^2 - 1))$  whose image is the nodal curve  $X$ . This is an immersed complex manifold. However, the image of the open set  $\overline{B_{1/2}(1)}^C$  is not open in  $X$ . We need to remove the node to get an open set.

*Remark.* I think the definition of algebraic leaves in Campana 2021 (that the Zariski and topological closures coincide) is wrong. For example, it predicts that the dense irrational slope foliation on an abelian surface is algebraic since each leaf is topologically and hence Zariski dense.

**Lemma 41.1.4.** Let  $\iota : L \rightarrow X$  be an immersed submanifold such that  $Z = \overline{\iota(L)}^{\text{Zar}}$  and  $L$  have the same dimension. Then  $L$  is algebraic.

*Proof.* Consider the map  $\iota : \iota^{-1}(Z^{\text{sm}}) \rightarrow Z^{\text{sm}}$  is a local diffeomorphism of smooth manifolds since it is an immersion of manifolds of the same dimension and hence is open.  $\square$

*Remark.* Algebraicity of course implies that  $L$  is a complex (immersed) submanifold.

**Lemma 41.1.5.** Let  $\iota : L \rightarrow X$  be an immersed submanifold with  $L$  connected. Then if  $Z = \overline{\iota(L)}^{\text{Zar}}$  has the same dimension as  $L$  then  $Z$  is irreducible.

*Proof.* We know  $\iota : L \rightarrow Z$  is analytically open away from the singularities. However,  $Z^{\text{sing}} \subset Z$  has codimension at least 2 and thus  $\iota^{-1}(Z^{\text{sing}})$  also has codimension at least 2 so  $L \setminus \iota^{-1}(Z^{\text{sing}})$  is connected. Thus  $\iota(L \setminus \iota^{-1}(Z^{\text{sing}})) \subset Z \setminus Z^{\text{sing}}$  so it must lie in some irreducible component (the irreducible components have become disconnected by removing the singularities). Since  $L \setminus \iota^{-1}(Z^{\text{sing}})$  is dense in  $L$  then  $\iota : L \rightarrow Z$  is contained in some irreducible component.  $\square$

*Remark.* This is false if we don't assume that  $\dim Z = \dim L$ . For example, there are embedded curves  $\mathbb{R} \rightarrow \mathbb{A}_{\mathbb{C}}^3$  whose closure is the union of two planes. Ineed, consider a curve which wanders in the  $xy$ -plane before following the  $x$ -axis then smoothly transitions to wandering in the  $xz$ -plane.

**Definition 41.1.6.** The *singular set* of a coherent subsheaf  $\mathcal{F} \subset T_X$  is,

$$\text{Sing}(\mathcal{F}) = \{x \in X \mid \mathcal{F}_x \rightarrow (T_X)_x \text{ is not a direct factor}\}$$

is union of the points where the rank of  $\mathcal{F}$  jumps up and where the rank of  $\mathcal{F} \rightarrow T_X$  jumps down.

**Definition 41.1.7.** A foliation  $\mathcal{F}$  on  $X$  is *algebraic* if every leaf of the regular foliation  $\mathcal{F}|_{U_{\mathcal{F}}}$  on  $U_{\mathcal{F}} = X \setminus \text{Sing}(\mathcal{F})$  is algebraic.

**Proposition 41.1.8.** If  $X$  is a smooth variety and  $\mathcal{F}$  regular algebraic foliation on  $X$  then every leaf is an *embedded* submanifold which is the analytification of a smooth algebraic subvariety.

*Proof.* It suffices to show that each leaf  $L$  is Zariski closed. Let  $Z$  be the Zariski closure of  $L$ . Choose  $p \in X$  and an open  $U$  such that  $U \cap L$  is a union of slices. Since  $Z$  is closed we may shrink  $U$  so that  $U \cap Z$  is connected. Since  $Z$  is irreducible,  $Z^{\text{sm}}$  is a connected embedded submanifold dense in  $Z$ . Then  $L \cap Z^{\text{sm}} \cap U \subset Z^{\text{sm}} \cap U$  is open and its closure in  $U$  is a union of slices but since  $Z \cap U$  is connected of dimension equal to the dimension of the slices it cannot contain more than one. Hence  $L \cap U$  is a single slice and is closed. Thus taking closures  $L \cap U = Z \cap U$  so  $L = Z$  and hence  $L$  is smooth.  $\square$

*Remark.* Without the algebraicity assumption, the leaves of  $\mathcal{F}$  do not even need to be closed. For example, the irrational slope foliation on an abelian variety.

**Proposition 41.1.9.** Suppose that a foliation  $\mathcal{F} \subset T_X$  is algebric. Then,

- (a) there is a Zariski open  $U \subset X$  and a morphism  $f : U \rightarrow S$  such that  $T_f = \mathcal{F}|_U$ .
- (b) if  $\mathcal{F}$  is a vector bundle then  $f$  is smooth
- (c) if  $\mathcal{F}$  is furthermore a sub-bundle then  $S$  is smooth. (WAIT BUT IF  $X$  IS SMOOTH AND  $f$  IS SMOOTH IT IMPLIES THAT  $S$  IS SMOOTH).

## 41.2 The Case $\kappa = -\infty$

Suppose we assume that  $\mu_{\alpha, \min}(\mathcal{F}^{\vee}) > 0$  (in the notation of Campana) then the Foliation is algebraic.

## 42 The Complex Geometric Picture

**Proposition 42.0.1.** There is a covariant equivalence of categories,

$$\{\mathbb{Z}\text{-Hodge Structures of type } (1, 0) \oplus (0, 1)\} \iff \{\text{complex Tori}\}$$

which specializes to,

$$\{\text{polarized } \mathbb{Z}\text{-Hodge Structures of type } (1, 0) \oplus (0, 1)\} \iff \{\text{abelian varieties}\}$$

*Remark.* If the polarization is required to be principal then the corresponding abelian variety is principally polarized.

**Corollary 42.0.2.** Nonconstant morphisms  $f : X \rightarrow A$  to a simple abelian variety (considered up to translation and isogeny) correspond to irreducible sub- $\mathbb{Q}$ -Hodge structures of  $H^1(X, \mathbb{Z})$ .

*Proof.* Morphisms  $f : X \rightarrow A$  sending a fixed base point  $x_0 \in X$  to  $0 \in A$  are equivalent to homomorphisms  $\text{Alb}_X \rightarrow A$  which correspond to maps of Hodge structures  $H^1(A, \mathbb{Z}) \rightarrow H^1(\text{Alb}_X, \mathbb{Z}) = H^1(X, \mathbb{Z})$ . Considered in the isogeny category, these correspond to maps  $H^1(A, \mathbb{Q}) \rightarrow H^1(X, \mathbb{Q})$  and since  $A$  is simple  $H^1(A, \mathbb{Q})$  is irreducible so the map is either zero or injective.  $\square$

The smoothness of the morphism  $f : X \rightarrow A$  is not in question. Smoothness is equivalent to  $f^*\Omega_A \rightarrow \Omega_X$  being a subbundle (of the correct dimension) i.e. if  $\omega_1, \dots, \omega_g \in H^0(A, \Omega_A)$  is a basis of holomorphic 1-forms then  $f^*\omega_1, \dots, f^*\omega_g \in H^0(X, \Omega_X)$  should form a partial frame (meaning they are everywhere independent).

### 42.1 Maps to Circles

**Proposition 42.1.1.** Let  $M$  be a compact smooth manifold and  $\omega \in M$  a closed nonvanishing 1-form. For any  $\epsilon > 0$  there exists a submersive map  $f_\epsilon : M \rightarrow S^1$  and an integer  $n_\epsilon$  such that,

$$\|\omega - n_\epsilon^{-1} f_\epsilon^* dt\| < \epsilon$$

in the  $L^\infty$  norm.

*Proof.* Because  $S^1$  is a  $K(\mathbb{Z}, 1)$  there are continuous maps  $f_i : M \rightarrow S^1$  such that  $f_i^*[S_1]$  form a basis of  $H^1(X, \mathbb{Z})$ . By [Prop. 17.8 of Bott, Tu, Differential Forms in Algebraic Topology] up to homotopy, we may choose the  $f_i$  smooth. By naturality of the de Rham comparison theorem,

$$\eta_i = f_i^* dt$$

form a basis of  $H_{\text{dR}}^1(X)$ . Thus we can write,

$$\omega = \sum_i \alpha_i \eta_i + dg$$

The idea is to rationally approximate the numbers  $\alpha_i \in \mathbb{R}$ . Indeed, we can choose rational numbers  $\frac{a_i}{n_\epsilon} \in \mathbb{Q}$  such that if let

$$\tilde{\omega} = \sum_i \frac{a_i}{n_\epsilon} \eta_i + dg$$

then we get,

$$\|\omega - \tilde{\omega}\| = \left\| \sum_i \left( \alpha_i - \frac{a_i}{n_\epsilon} \right) \eta_i \right\| < \epsilon$$

this requires choosing the rational approximation on the order of  $\frac{\epsilon}{\text{Vol}(M)}$ . The let,

$$f_\epsilon = \left( \prod_i f_i^{a_i} \right) \cdot (\Pi \circ g)^{n_\epsilon}$$

where  $\Pi : \mathbb{R} \rightarrow S^1$  is the universal cover. Therefore,

$$f_\epsilon^* dt = \sum_i a_i f_i^* dt + n_\epsilon dg = \sum_i a_i \eta_i + n_\epsilon dg = n_\epsilon \tilde{\omega}$$

proving the required inequality. Finally, for sufficiently small  $\epsilon$ , since  $\omega$  is nonvanishing we see that  $\tilde{\omega}$  is also nonvanishing so  $f_\epsilon$  is smooth.  $\square$

*Remark.* Let's see what happens when we try to do this for a holomorphic 1-form  $\omega \in H^0(X, \Omega_X)$ . Write  $\omega = \omega_1 + i\omega_2$  into its real and imaginary parts. Note that because  $\omega$  is holomorphic  $d\omega = 0$  (this requires  $X$  compact Kähler). Therefore, we can rationally approximate,  $\tilde{\omega}_1$  and  $\tilde{\omega}_2$  to get a submersive (because for small enough  $\epsilon$  we can ensure that  $\tilde{\omega}_1$  and  $\tilde{\omega}_2$  are everywhere independent) map  $f : X \rightarrow S^2$  with  $f^*dz = n\tilde{\omega}$  with  $\tilde{\omega} = \tilde{\omega}_1 + i\tilde{\omega}_2$  but there is no reason that  $\tilde{\omega}_2$  should be holomorphic. Suppose we could approximate  $\omega$  by a rational form which is holomorphic. This is exactly a  $\mathbb{Q}$ -Hodge submodule of  $H^1(X, \mathbb{Q})$  of rank 1. Therefore, we are in the business of showing that  $H^1(X, \mathbb{Q})$  is a reducible Hodge structure.

## 42.2 An Idea

Consider the canonical fibration  $f : X \rightarrow S$ . If  $X$  has a nonvanishing 1-form  $\omega$  then Popa-Schnell shows that the map  $X \rightarrow \text{Alb}_X$  cannot fully contract the fiber  $F$  of  $X \rightarrow S$ . If we can show that  $\dim \text{Alb}_F < \dim \text{Alb}_X$  then this means that  $\text{Alb}_X$  is reducible.

## 43 Reflexivity

**Proposition 43.0.1.** Let  $\mathcal{F}$  be a coherent sheaf on an integral scheme  $X$ . Then  $\mathcal{F}^*$  is reflexive.

*Proof.* Let  $\psi : \text{id} \rightarrow (-)^{**}$  be the double dual natural transformation. There are maps,

$$\mathcal{F}^* \xrightarrow{\psi_{\mathcal{F}^*}} \mathcal{F}^{***} \xrightarrow{\psi_{\mathcal{F}^{***}}} \mathcal{F}^*$$

which acts as follows, for  $\varphi \in \mathcal{F}^*$  and  $\ell \in \mathcal{F}^{**}$ ,

$$\varphi \mapsto (\ell \mapsto \ell(\varphi)) \mapsto [(\ell \mapsto \ell(\varphi)) \circ \psi_{\mathcal{F}}]$$

the result of which is the function,

$$x \mapsto \psi_{\mathcal{F}}(\varphi' \mapsto \varphi'(x)) \mapsto \varphi(x)$$

which is just  $\varphi$  so indeed this is the identity. Thus it suffices to show that the second map is injective. However,  $\mathcal{F}^{***}$  is torsion-free and hence it suffices to check this at the generic point where it becomes finite-dimensional linear algebra.  $\square$

Consider a separable map  $f : X \rightarrow C$  where  $X$  is a smooth proper (integral) surface over  $k$  and  $C$  a smooth proper (integral) curve over  $k$ . Then there is a sequence,

$$0 \longrightarrow f^*\Omega_C \longrightarrow \Omega_X \longrightarrow \Omega_{X/C} \longrightarrow 0$$

which is injective on the left by generic smoothness and the fact that  $\Omega_C$  and  $\Omega_X$  are vector bundles. Then we get an exact sequence,

$$0 \longrightarrow \mathcal{F} \longrightarrow \mathcal{T}_X \longrightarrow f^*\mathcal{T}_C \longrightarrow \mathcal{E}xt_{\mathcal{O}_X}^1(\Omega_{X/C}, \mathcal{O}_X) \longrightarrow 0$$

again using that  $\Omega_X$  is a vector bundle where we set  $\mathcal{F} = \Omega_{X/C}^\vee$ . Since  $\mathcal{F}$  is reflexive and  $X$  is a regular surface we see that  $\mathcal{F}$  is a vector bundle. By generic smoothness,  $\mathcal{F}$  has rank 1. I claim that  $\mathcal{F}$  is closed under Lie bracket.

Indeed, this is a local question so we reduce to  $R \rightarrow A \rightarrow B$  are ring maps then  $\text{Hom}_B(\Omega_{B/A}, B) \rightarrow \text{Hom}_B(\Omega_{A/R}, B)$  is closed under Lie bracket. Indeed, for  $X, Y \in \text{Hom}_B(\Omega_{B/A}, B)$  we need to show that  $(X \circ d)(Y \circ d) - (Y \circ d)(X \circ d)$  is a  $A$ -derivation not just an  $R$ -derivation. This is basically obvious because  $X \circ d$  and  $Y \circ d$  kill  $A$ .

Now we get an exact sequence,

$$0 \longrightarrow \mathcal{T}_X/\mathcal{F} \longrightarrow f^*\mathcal{T}_C \longrightarrow \mathcal{E}xt_{\mathcal{O}_X}^1(\Omega_{X/C}, \mathcal{O}_X) \longrightarrow 0$$

Since  $f^*\mathcal{T}_C$  is torsion-free we see that so is  $\mathcal{T}_X/\mathcal{F}$  so  $\mathcal{F}$  is automatically saturated.