

- 1 An object is spun around in circular motion such that it completes 100 cycles in 25 s.
 - a. What is the period of its rotation?
 - b. What is the frequency of its rotation?

- 2 An object completes 2500 cycles in 25 s.
 - a. What is the period of its rotation?
 - b. What is the frequency of its rotation?

- 3 An object is spun around in circular motion such that its period is 12s.
 - a. What is the frequency of its rotation?
 - b. How much time will be required to complete 86 rotations?

- 4 An object completes 10 cycles in 50 s.
 - a. What is the period of its rotation?
 - b. What is the frequency of its rotation?

- 5 An object is spun around in circular motion such that its frequency is 12 Hz.
- a. What is the period of its rotation?
 - b. How much time will be required to complete 86 rotations?

- 6 An object is spun around in circular motion such that its frequency is 500 Hz.
 - a. What is the period of its rotation?
 - b. How much time will be required to complete 7 rotations?

- 7 A 5.0 kg object is spun around in a circle of radius 1.0 m with a period of 4.0s.
- a. What is the frequency of its rotation?
 - b. *What is its velocity?
 - c. *What is its acceleration?

- 8 A 15.0 kg mass is spun in a circle of radius 5.0 m with a frequency of 25 Hz.
- a. What is the period of its rotation?
 - b. *What is its velocity?
 - c. *What is its acceleration?

- 9 A 0.5 kg object is spun around in a circle of radius 2.0 m with a period of 10.0s.
- a. What is the period of its rotation?
 - b. *What is its velocity?
 - c. *What is its acceleration?

- 10 A 500 kg mass is spun in a circle of radius 25 m with a velocity of 250 m/s.
- a. *What is the period of its rotation?
 - b. *What is its frequency?
 - c. What is its acceleration?

- 11 What is the acceleration of an object that has a velocity of 25 m/s and is moving in a circle of radius 10m?

- 12 An object is experiencing an acceleration of 12 m/s^2 while traveling in a circle at a velocity of 3.1 m/s . What is the radius of its motion?

- 13 A 61 kg object is experiencing a net force of 25 N while traveling in a circle of radius 35 m. What is its velocity?

- 14 A 0.25 kg object is experiencing a net force of 15 N while traveling in a circle at a velocity of 21 m/s. What is the radius of its motion?

- 15 An object is experiencing a centripetal acceleration of 36 m/s^2 while traveling in a circle of radius 15 m. What is its velocity?

- 16 A 61 kg object is experiencing a net force of 250 N while traveling in a circle of radius 1.5 m. What is its velocity?

- 17 An object is experiencing an acceleration of 12 m/s^2 while traveling in a circle of radius 5.0 m . What is its velocity?

- 18 What is the net force acting on a 5.0 kg object that has a velocity of 15 m/s and is moving in a circle of radius 1.6m?

- 19 What is the acceleration of an object that has a velocity of 37 m/s and is moving in a circle of radius 45m?

- 20 An object is experiencing a centripetal acceleration of 2.0 m/s^2 while traveling in a circle at a velocity of 0.35 m/s . What is the radius of its motion?

- 21 What is the net force acting on a 52 kg object that has a velocity of 17 m/s and is moving in a circle of radius 1.6m?

- 22 A 6.8 kg object is experiencing a net force of 135 N while traveling in a circle at a velocity of 45 m/s. What is the radius of its motion?

- 23 *A 0.65 kg ball is attached to the end of a string. It is swung in a vertical circle of radius 0.50 m. At the top of the circle its velocity is 2.8 m/s.
- a. Draw a free body diagram for the ball when it is at the top of the circle. Next to that diagram indicate the direction of its acceleration.
 - b. Use that free body diagram to set up the equations needed to determine the Tension in the string.
 - c. Solve those equations for the Tension in the string.

- 24 *A 0.65 kg ball is attached to the end of a string. It is swung in a vertical circle of radius 0.50 m. At the bottom of the circle its velocity is 2.8 m/s.
- Draw a free body diagram for the ball when it is at the bottom of the circle. Next to that diagram indicate the direction of its acceleration.
 - Use that free body diagram to set up the equations needed to determine the Tension in the string.
 - Solve those equations for the Tension in the string.

- 25 *A 0.25 kg ball is attached to the end of a string. It is swung in a vertical circle of radius 0.6 m. At the top of the circle its velocity is 3 m/s.
- a. Draw a free body diagram for the ball when it is at the top of the circle. Next to that diagram indicate the direction of its acceleration.
 - b. Use that free body diagram to set up the equations needed to determine the Tension in the string.
 - c. Solve those equations for the Tension in the string.

- 26 *A 0.25 kg ball is attached to the end of a string. It is swung in a vertical circle of radius 0.6 m. At the bottom of the circle its velocity is 3 m/s.
- a. Draw a free body diagram for the ball when it is at the top of the circle. Next to that diagram indicate the direction of its acceleration.
 - b. Use that free body diagram to set up the equations needed to determine the Tension in the string.
 - c. Solve those equations for the Tension in the string.

- 27 *A ball is attached to the end of a string. It is swung in a vertical circle of radius 1.5 m. What is the minimum velocity that the ball must have to make it around the circle?

- 28 *A ball is attached to the end of a string. It is swung in a vertical circle of radius 0.75 m. What is the minimum velocity that the ball must have to make it around the circle?

- 29 **A car is going over the top of a hill whose curvature approximates a circle of radius 200 m. At what velocity will the occupants of the car appear to weigh 20% less than their normal weight (or their normal weight times 0.8)?

- 30 **A car is going through a dip in the road whose curvature approximates a circle of radius 200 m. At what velocity will the occupants of the car appear to weigh 20% more than their normal weight (or their normal weight times 1.2)?

- 31 **The occupants of a car traveling at a speed of 30 m/s note that on a particular part of a road their apparent weight is 15% higher than their weight when driving on a flat road.
- a. Is that part of the road a hill or a dip?
 - b. What is the vertical curvature of the road?

- 32 *A ball is attached to the end of a string. It is swung in a vertical circle of radius 0.33 m. What is the minimum velocity that the ball must have to make it around the circle?

- 33 *A ball is attached to the end of a string. It is swung in a vertical circle of radius 2.5 m. What is the minimum velocity that the ball must have to make it around the circle?

- 34 **A car is going over the top of a hill whose curvature approximates a circle of radius 350m. At what velocity will the occupants of the car appear to weigh 10% less than their normal weight?

- 35 **A car is going through a dip in the road whose curvature approximates a circle of radius 150m. At what velocity will the occupants of the car appear to weigh 15% more than their normal weight?

- 36 **The occupants of a car traveling at a speed of 40 m/s note that on a particular part of a road their apparent weight is 30% lower than their weight when driving on a flat road.
- a. Is that part of the road a hill or a dip?
 - b. What is the vertical curvature of the road?

- 37 **A car, traveling at a speed of 25 m/s, rounds a flat curve whose radius is 125 m.
- a. Draw a side view free body diagram for the car. Indicate the direction of acceleration.
 - b. Use that free body diagram to set up the equations needed to determine the frictional force acting on the car.
 - c. Solve those equations for the coefficient of friction between the tires and the road.

- 38 **A car, traveling at a speed of 32 m/s, rounds a flat curve whose radius is 250 m.
- Draw a side view free body diagram for the car. Indicate the direction of acceleration.
 - Use that free body diagram to set up the equations needed to determine the frictional force acting on the car.
 - Solve those equations for the coefficient of friction between the tires and the road.