



## **New Jersey Center for Teaching and Learning**

### **Progressive Science Initiative**

This material is made freely available at [www.njctl.org](http://www.njctl.org) and is intended for the non-commercial use of students and teachers. These materials may not be used for any commercial purpose without the written permission of the owners. NJCTL maintains its website for the convenience of teachers who wish to make their work available to other teachers, participate in a virtual professional learning community, and/or provide access to course materials to parents, students and others.

**Click to go to website:**  
**[www.njctl.org](http://www.njctl.org)**



# Fluids

[www.njctl.org](http://www.njctl.org)


# Table of Contents

[\*Click on the topic to go to that section\*](#)

- **Density**
- **Specific Gravity**
- **Pressure in Fluids**
- **Atmospheric Pressure and Gauge Pressure**
- **Pascal's Principal**
- **Buoyancy and Archimedes' Principle**
- **Fluids in Motion & Bernoulli's Principle**
- **Torricelli's Theorem**

# Density


[Return to Table  
of Contents](#)



<http://njc.tl/ua>

# Density

You may recall that the three common phases, or states, of matter are gas, liquid, and solid. Solids maintain a fixed volume and shape, liquids maintain a fixed volume but not shape, and gases can change both. Since gasses and liquids both flow, they are collectively called fluids.



<http://njc.tl/ua>


# Density

What weighs more a pound of feathers or a pound of bricks?

This is a silly questions since they are both a pound.

Sometimes people say that iron is "heavier" than wood. But if you have a log of wood it would be heavier than one small iron nail.

What we should really say is that iron is more dense than wood.



# Density

The density of an object is its mass per unit volume:


$$\rho = \frac{m}{V}$$

$\rho$  (rho) is density.

$m$  is mass.

$V$  is volume.

The SI unit for density is  $\text{kg/m}^3$  but sometimes it is measured in  $\text{g/cm}^3$ . To convert from  $\text{g/cm}^3$  to  $\text{kg/m}^3$  multiply by 1000.



1 The density of a substance is  $\rho$  , its mass is  $m$  and its volume is  $V$ . If the volume is tripled, what is the new mass?

- ☐ A  $m/3$
- ☐ B  $3m$
- ☐ C  $m$
- ☐ D  $m/6$
- ☐ E  $6m$


Answer






2 Liquid A has twice the density of liquid B. A certain experiment needs samples of A and B that have the same mass. What needs to be true about their volumes?

- ☐ A  $V_A = V_B$
- ☐ B  $2V_A = V_B$
- ☐ C  $V_A = 2V_B$
- ☐ D  $V_A/2 = V_B$
- ☐ E  $V_A = 4V_B$

**Answer**

3 What is the density (in  $\text{kg/m}^3$ ) of an object that has a mass of 2kg and a volume of  $4\text{m}^3$ ?


Answer



<http://njc.tl/ud>

- 4 A container of water has a mass of 5kg. What is the volume of this container (in  $\text{m}^3$ )? The density of water is  $1000 \text{ kg/m}^3$ . (Neglect the mass of the container.)


Answer



<http://njc.tl/ue>

# Specific Gravity

[Return to Table  
of Contents](#)




<http://njc.tl/uf>

# Specific Gravity

The specific gravity of a substance is the ratio of its density to the density of water.

$$SG = \frac{\rho}{\rho_{H_2O}}$$

The density of water at 4° C is 1 g/cm<sup>3</sup> or 1000 kg/m<sup>3</sup>.





<http://njc.tl/uf>

# Specific Gravity

Specific gravity is a ratio so it has no units.

A substance with a specific gravity less than one means that it is less dense than water and will float on water and a substance with a specific gravity greater than one means that it is more dense than water and will sink in water.




<http://njc.tl/uf>

5 The following are specific gravities of various objects. Which would float on water?

- ☐ A Copper - 8.96
- ☐ B Gold - 19.3
- ☐ C Aluminum - 2.7
- ☐ D Oak - 0.78
- ☐ E Table Salt - 2.17


Answer



<http://njc.tl/ug>

6 What is the specific gravity of a substance whose density is  $450 \text{ kg/m}^3$ ?

Answer




<http://njc.tl/uh>



7 Mercury's specific gravity is about 13.5. What is its density in  $\text{kg/m}^3$ ?


**Answer**



<http://njc.tl/ui>

# Pressure in Fluids

[Return to Table  
of Contents](#)



<http://njc.tl/uj>

# Pressure in Fluids


Pressure is defined as the force per unit area.

$$P = \frac{F}{A}$$


Pressure is a scalar and its units are in Pascals.

$$1\text{Pa} = \text{N/m}^2.$$

This definition of pressure is true in any situation, not just fluids. You can see from the equation that pressure is related to force and area. Think about what it would mean to get your foot stepped on by a sneaker or a high heel. Which would hurt more? Why?





<http://njc.tl/uj>



# Pressure in Fluids

Fluids can exert a pressure normal to any contact surface.


Pressure is the same in every direction in a fluid at a given depth. If it were not, the fluid would flow.



<http://njc.tl/uj>


8 A perpendicular force is applied to a certain area and produces a pressure  $P$ . If the same force is applied to half the area, the new pressure on the surface is:

- ☐ A  $2P$
- ☐ B  $4P$
- ☐ C  $P$
- ☐ D  $P/2$
- ☐ E  $P/4$

**Answer**

9 A 50kg person stands on a square board with sides of 2m. What is the pressure (in Pa) exerted on the ground by the board?

Answer



<http://njc.tl/ul>

# Pressure in Fluids

$$P = \frac{F}{A}$$

The pressure at a depth of  $h$  below the surface of the fluid is due to the weight ( $mg$ ) of the fluid above it.

$$P = \frac{mg}{A}$$

$$P = \frac{mgh}{Ah}$$


Multiply top and bottom by  $h$ .

$$P = \frac{mgh}{V}$$

$$V = Ah$$


$$P = \rho gh$$

$$\rho = m/V$$




# Pressure in Fluids

$$P = \rho gh$$



The pressure at a given point depends on only the density of the fluid and the depth. (Not the shape of the container.)




<http://njc.tl/um>

This is valid for liquids whose density does not change with depth.



10 There are five containers of the same fluid in a physics lab. Which has the greatest pressure at the bottom of the container?



☐ A


☐ B

☐ C

☐ D

☐ E


Answer



<http://njc.tl/un>

11 What is the pressure (in Pa) at the bottom of a swimming pool whose depth is 2m?


Answer



<http://njc.tl/uo>

# Atmospheric Pressure and Gauge Pressure

[Return to Table of Contents](#)



<http://njc.tl/up>

# Atmospheric Pressure and Gauge Pressure

At sea level, the atmospheric pressure is about  $1.013 \times 10^5 \text{ Pa}$ .  
This is called **1 atm**.


Another unit of pressure is the bar.

$$1 \text{ bar} = 1.00 \times 10^5 \text{ Pa}.$$

Most pressure gauges measure the pressure above atmospheric pressure. This is called gauge pressure.

Absolute pressure is atmospheric pressure plus gauge pressure.


$$P = P_A + P_G$$



# Atmospheric Pressure and Gauge Pressure

Torricelli invented a mercury barometer to measure atmospheric pressure. Sometimes air pressure is described in millimeters or inches of mercury.



A glass tube is filled with mercury. This glass tube sits upside down in a container, called the reservoir, which also contains mercury. The mercury level in the glass tube falls, creating a vacuum at the top.



# Atmospheric Pressure and Gauge Pressure

The barometer works by balancing the weight of mercury in the glass tube against the atmospheric pressure. If the weight of mercury is less than the atmospheric pressure, the mercury level in the glass tube rises. If the weight of mercury is more than the atmospheric pressure, the mercury level falls.

Atmospheric pressure is basically the weight of air in the atmosphere above the reservoir, so the level of mercury continues to change until the weight of mercury in the glass tube is exactly equal to the weight of air above the reservoir.




<http://njc.tl/up>

12 A diver in the ocean measures guage pressure to be 515kPa.  
What is the absolute pressure?

- ☐ A 101kPa
- ☐ B 313kPa
- ☐ C 515kPa
- ☐ D 616kPa
- ☐ E 5150kPa


Answer



<http://njc.tl/uq>

13 What is the absolute pressure (in Pa) at the bottom of a swimming pool whose depth is 2m?

**Answer**




<http://njc.tl/ur>




# Pascal's Principal

[Return to Table  
of Contents](#)




<http://njc.tl/ur>

# Pascal's Principle



Pascal's principle states that if an external pressure is applied to a confined and incompressible fluid, the pressure everywhere in the fluid increases by that amount.

Pascal's Barrel is an experiment attributed to Pascal but it is unclear if it was ever performed by him. In this experiment, a 10 meter long tube was inserted into a barrel filled with water. When water was poured into the tube, the increase in pressure caused the barrel to burst.





<http://njc.tl/ur>

# Pascal's Principle

$$P_{in} = P_{out}$$

$$\frac{F_{in}}{A_{in}} = \frac{F_{out}}{A_{out}}$$




<http://njc.tl/ur>

14 In a hydraulic lift, the large piston has five times the area as the small piston. How much extra force can the large piston exert?

- ☐ A One tenth as much as the small piston
- ☐ B One fifth as much as the small piston
- ☐ C The same as the small piston
- ☐ D Five times as much as the small piston
- ☐ E Fifty times as much as the small piston


Answer



<http://njc.tl/us>

15 The small piston of a hydraulic lift has an area of  $10\text{cm}^2$  and its large piston has an area of  $100\text{ cm}^2$ . A  $40\text{ N}$  force is applied to the small piston. What is the weight of the load can be lifted by the large piston?


Answer



<http://njc.tl/ut>

# Buoyancy and Archimedes' Principle



[Return to Table  
of Contents](#)



<http://njc.tl/uu>

# Buoyancy and Archimedes' Principle


The upward buoyant force on an object immersed in a fluid, partially or completely, is equal to the weight of the displaced fluid.



<http://njc.tl/uu>

# Buoyancy and Archimedes' Principle

If an object is submerged in a fluid, there is a net force on the object because the pressure is greater at the bottom than at the top of the object. The buoyant force is upward because the force is greater at bottom than at the top of the object.



$$F_B = F_2 - F_1$$


$$F_B = P_2 A - P_1 A$$

$$F_B = \rho_F g h_2 A - \rho_F g h_1 A$$

$$F_B = \rho_F g A (h_2 - h_1)$$

$$F_B = \rho_F g V$$


$$F_B = m_F g$$



<http://njc.tl/uu>



# Buoyancy and Archimedes' Principle



$$F_B = \rho_F g V$$


$$F_B = m_F g$$

Where:

$\rho_F$  is the density of the fluid.


$m_F$  is the mass of the displaced fluid.

$V$  is the volume of the displaced fluid.




# Buoyancy and Archimedes' Principle


The net force on a object is the difference between the buoyant force and the gravitational force.



$$F_{\text{scale}} = mg$$




$$F_{\text{scale}} + F_B = mg$$



16 Three objects of the same volume but different materials are completely submerged in water. They are zinc with a density of  $7000 \text{ kg/m}^3$ , nickel with a density of  $8900 \text{ kg/m}^3$ , and silver with a density of  $10500 \text{ kg/m}^3$ . Which has the greatest buoyant force exerted on it?

- ☐ A Zinc
- ☐ B Nickel
- ☐ C Silver
- ☐ D They all have the same buoyant force.
- ☐ E It is impossible to tell without knowing the volume.

Answer




<http://njc.tl/uv>

17 A metal sphere weights 5N in air and 3N when it is submerged in water. What is the buoyant force on the sphere when it is submerged in water?


- ☐ A 0.2N
- ☐ B 2N
- ☐ C 3N
- ☐ D 5N
- ☐ E 8N

Answer



18 An object has a volume of  $2.0 \text{ m}^3$ . What is the buoyant force on the object when it is completely submerged into water (density  $1000 \text{ kg/m}^3$ )?

Answer



<http://njc.tl/ux>


# Buoyancy and Archimedes' Principle

$$F_B = m_{\text{fluid}}g$$

Any floating object  
displaces its own weight of  
fluid.



$$m_{\text{boat}}g$$




<http://njc.tl/uy>

# Buoyancy and Archimedes' Principle

For an object whose density is less than that of the fluid, there will be a net force upward and it will rise until it is partially out of the fluid.

For a floating object, the fraction that is submerged is given by the ratio of the object's density to that of the fluid.



$$F_B = mg$$

$$\rho_F g V_{displ} = \rho_O g V_O$$


$$\rho_F V_{displ} = \rho_O V_O$$

$$\frac{V_{displ}}{V_O} = \frac{\rho_O}{\rho_F}$$



19 A 1500 N object floats in water. What is the weight of displaced water?

Answer




<http://njc.tl/uz>



20 A small empty row boat with a mass of 48kg floats on water.  
What is the volume of the water it displaces?


**Answer**



<http://njc.tl/v0>

# Fluids in Motion & Bernoulli's Principle

[Return to Table  
of Contents](#)



<http://njc.tl/v1>

# Fluids in Motion & Bernoulli's Principle

If the flow of a fluid is smooth, it is called streamline or laminar flow. This is what we will deal with.


The mass flow rate is the mass that passes a given point per unit time. The flow rates of a fluid must be equal, as long as no fluid is added or taken away.

This gives us the equation of continuity:

$$\rho_1 A_1 v_1 = \rho_2 A_2 v_2$$

If the density of the fluid doesn't change, it can be simplified to:


$$A_1 v_1 = A_2 v_2$$



# Fluids in Motion & Bernoulli's Principle


$$A_1 v_1 = A_2 v_2$$

Density does not typically change in liquids. This means that where a pipe is wider the flow is slower.



# Fluids in Motion & Bernoulli's Principle

You can see this happening in a river when the water flow is slow when it is wide and fast when it is narrow.




<http://njc.tl/v1>

21 Water flows at a constant speed through one section of a pipe, when it enters another section that is half the cross sectional area what happens to the speed of the water?

- ☐ A The speed is reduced to one fourth the original.
- ☐ B The speed is reduced to one half the original.
- ☐ C The speed stays the same.
- ☐ D The speed is doubled.
- ☐ E The speed is quadrupled.


Answer



<http://njc.tl/v2>

22 Water flows through a pipe of cross-sectional area  $10 \text{ cm}^2$  at a rate of  $15 \text{ m/s}$ . The cross-sectional area of the pipe is decreased to  $5 \text{ cm}^2$ . What is the water rate in the narrow section of the pipe?

Answer





<http://njc.tl/v3>

# Fluids in Motion & Bernoulli's Principle

A fluid can also change height. If we look at the work done...

$$P + \frac{1}{2} \rho v^2 + \rho gh = \text{constant}$$








# Fluids in Motion & Bernoulli's Principle

One thing this tells us is that as the speed of the water flow goes up, the pressure goes down.

$$P + \frac{1}{2} \rho v^2 + \rho gh = \text{constant}$$




23 A pipe has three different sections with three different cross-sectional area. Where is the pressure the least?




- ☐ A  $P_1$
- ☐ B  $P_2$
- ☐ C  $P_3$
- ☐ D The pressure is the same in all three sections.
- ☐ E The pressures cannot be determined.

Answer



24 Water flows through a horizontal pipe at a speed of 10 m/s and pressure  $2.5 \times 10^5$  Pa. The pipe narrows and the water speed goes up to a 20 m/s. What is the pressure in the narrow section of the pipe?


**Answer**



<http://njc.tl/v6>


# Torricelli's Theorem

[Return to Table  
of Contents](#)




<http://njc.tl/v7>

# Torricelli's Theorem



We can use Bernoulli's Principle, to find the speed of a fluid coming out to spigot of an open tank. This is called Torricelli's Theorem.


$$v_1 = \sqrt{2g(y_2 - y_1)}$$



25 A container of water has spigot at its bottom. What happens to the water speed out of the spigot as the container empties?


- ☐ A The water speed decreases.
- ☐ B The water speed increases.
- ☐ C The water speed stays the same.

Answer



26 A container holds water at a depth of 5 m. There is a hole in the bottom of the container. At what speed will water flow out of the hole?

Answer



<http://njc.tl/v9>

# Summary

## Density

$$\rho = \frac{m}{V}$$

## Specific Gravity

$$SG = \frac{\rho}{\rho_{H_2O}}$$

## Pressure

$$P = \frac{F}{A}$$

## Pressure in Fluids

$$P = \rho gh$$

## Buoyant Force

$$F_B = \rho_F gV$$

$$F_B = m_F g$$

## Equation of Continuity

$$\rho_1 A_1 v_1 = \rho_2 A_2 v_2$$

## Bernoulli's Principle

$$P + \frac{1}{2} \rho v^2 + \rho gh = \text{constant}$$