

# Study of Bifurcations on Purkinje Fibers


Lucas Berg

11 de maio de 2017

# Summary

- 1 Purkinje Fibers
- 2 Source-Sink Problem
- 3 Under Development

# Purkinje Fibers



# Source-Sink Problem

- When an action potential (AP) propagates in cardiac tissue, the wavefront acts as a source of depolarizing current for the adjacent repolarized tissue (the sink).
- The source current density must be sufficient to bring the sink to its activation threshold, and propagation will fail if the source-sink mismatch is too large. This topic has been extensively studied in both normal and diseased heart for propagating wavefronts
- In order to elicit an action potential in a cell, it must receive a depolarizing current from an adjacent, activated cell. The activated cell acts as current source, while the non-activated cell is a current sink with the voltage difference being the driving force for this current. The current transfer is mainly realized via gap junction channels and, to some extent, via the extracellular space.

# Source-Sink Problem

- Whether enough current can be transferred to activate a cell is a complex and geometric problem: if a small source (e.g., a tiny strand of activated cardiomyocytes) meets a large sink (e.g., a large area of non-activated site to many non-activated cardiomyocytes) the current will flow radially from the activated site to many non-activated sites.
- Hence, the source current is distributed to many neighboring cells and in each of these the accumulated charge may be too low to trigger an action potential. This will cause conduction failure.

# Source-Sink Problem

- Situations with source-sink problems generally occur when the curvature of the propagating wave front is high. Accordingly, they may be found at the end of Purkinje Fibers, during propagation through small isthmuses, around obstacles, and during spiral wave re-entry.
- Furthermore, source-sink mismatch may occur at the border between normal cardiac tissue and an ischemic zone, when depolarizing current flows into the ischemic region. Since this region is usually non-excitable, it will act as a current sink, consequently, reduce conduction velocity.

# Source-Sink Problem

COLOCAR UMA FIGURA

# Under Development

- **Objective:** study how bifurcations on Purkinje Fibers affect the source-sink mismatch.
- **Equation:** Monodomain
- **Models:** Noble (1962) and Li & Rudy (2011)
- **Methods to use:** Finite Volumes and Finite Elements
- **Parameters to analyze:**
  - Number of bifurcations.
  - Propagation velocity.
  - Size of the fibers.
  - Size of the discretization in space.



COLOCAR UMA FIGURA (Bif 4 e Bif 40)