



Department of Computer Engineering

CENG 384 Homework Exam 1 Key

22 April 2022

METU Honor Code and Pledge

METU Honor Code: “The members of the METU community are reliable, responsible and honourable people who embrace only the success and recognition they deserve, and act with integrity in their use, evaluation and presentation of facts, data and documents.”

*“I have read and understood the implications of the METU Honor Code. To be precise, I understand that this is a **open** formula-sheet exam, and I am forbidden to access any other source of information other than provided within the exam. I will **TURN OFF** all my electronic equipment (phones, smart watches, etc.) and put it off the table along with other notes and materials that I may have with me. I understand that leaving electronic devices on during the exam is strictly forbidden. I understand and accept to obey all the rules announced by the course staff, and that failure to obey these will result in disciplinary action.”*

Name, SURNAME:
ID:

Signature:

Specifications About the Exam

- **Duration:** X minutes.
- **Notes:**
 - One A4-size double-sided formula-sheet is allowed.
 - Each answer should contain a justifying explanation. However, give shortest possible answers. Unnecessary or irrelevant wordings will reduce your grades.
 - If you use the back of a page for extra space, please mark down a proper notice on the relevant pages.

Grade

Q1:

Q2:

Q3:

Q4:

Q5:

Total:

1 (10 pts, 2 pts each) Determine whether the following statements are *True* or *False*. Circle the correct answer for each question.

- a) [T / F] $x[n] * h[n] * x[n] = h[n] * (x[n])^2$.
- b) [T / F] $y(t) = x(2t)$ is not linear.
- c) [T / F] Linearity is a necessary condition for time invariance.
- d) [T / F] $x[n] = \text{Re}\{(1+j)^n\}$ is not periodic.
- e) [T / F] Let $z = \frac{(1+j)(5-5j)(\sqrt{3}+j)}{10j(5+j5\sqrt{3})}$; z evaluates to $\frac{1}{5}e^{\frac{-j2\pi}{3}}$.

- a) **False**
- b) **False**
- c) **False**
- d) **True**
- e) **True**

2 (24 pts, 12 pts each) Which of the following properties do the systems below have?

a) $y(t) = x(t + 2)$

b) $y(t) = \int_{-\infty}^{2t} x(3\tau) d\tau$

Memory, stability, causality, linearity, invertibility and time-invariance. Clearly explain your reasoning to get full credit.

a) $y(t) = x(t + 2)$

- **Memory:** Has memory, $y(0) = x(2)$
- **Stability:** Stable, all bounded inputs result in bounded outputs
- **Causality:** Not Causal $y(2) = x(4)$, y depends on inputs in the future
- **Linearity:** Superposition property holds, therefore linear
- **Invertibility:** Invertible, $x(t) = y(t - 2)$
- **Time Invariance:** Time invariant, shift in time results in identical time shift

b) $y(t) = \int_{-\infty}^{2t} x(3\tau) d\tau$

- **Memory:** Has memory, output depends on past values of input
- **Stability:** Not stable, even we bound input, we cannot bound the response.
- **Causality:** Not Causal, system uses inputs from the future
- **Linearity:** Superposition property holds, therefore linear
- **Invertibility:** Invertible, we can invert by differentiating
- **Time Invariance:** Time varying, output is not shifted with same value if we shift time in the input.


- 3 (10 pts) Given that $x_1[n]$ is an even signal and $x_2[n]$ is an odd signal, show whether $x_1[n] \cdot x_2[n]$ is an odd signal or an even signal. Show your work.

Since $x_1[n]$ is even, $x_1[n] = x_1[-n]$.

Similarly, since $x_2[n]$ is odd, $x_2[n] = -x_2[-n]$.

$$\begin{aligned} y[n] &= x_1[n] \cdot x_2[n] \\ y[-n] &= x_1[-n]x_2[-n] \\ &= x_1[n](-x_2[n]) \\ &= -y[n] \implies y[n] = x_1[n] \cdot x_2[n] \text{ is odd.} \end{aligned}$$

- 4 (30 pts, 5-15-10 pts respectively) Consider an LTI system given by the following block diagram:



- a) Find the differential equation which represents this system.
b) Find the output $y(t)$, when the input $x(t) = (e^{-t} + e^{-3t})u(t)$. Assume that the system is initially at rest.
c) Find the impulse response of this system. Show your work.

a)

$$y'(t) + 5y(t) = x(t)$$

- b) char eqn. : $r + 5 = 0 \Rightarrow r = -5 \Rightarrow y_h(t) = K \cdot e^{-5t}$
 $y_p(t) = Ae^{-t} + Be^{-3t}$
 $y'_p(t) = -Ae^{-t} - 3Be^{-3t}$

$$\begin{aligned} -Ae^{-t} - 3Be^{-3t} + 5(Ae^{-t} + Be^{-3t}) &= e^{-t} + e^{-3t} \\ 4Ae^{-t} + 2Be^{-3t} &= e^{-t} + e^{-3t} \end{aligned}$$

$$4A = 1 \Rightarrow A = \frac{1}{4}, 2B = 1 \Rightarrow B = \frac{1}{2}$$

$$y(t) = y_h(t) + y_p(t) = (Ke^{-5t} + \frac{1}{4}e^{-t} + \frac{1}{2}e^{-3t})u(t)$$

$$y(0) = K + \frac{1}{4} + \frac{1}{2} = 0 \Rightarrow K = -\frac{3}{4}$$

$$y(t) = (\frac{1}{4}e^{-t} + \frac{1}{2}e^{-3t} - \frac{3}{4}e^{-5t})u(t)$$

c)

$$y'(t) + 5y(t) = x(t)$$

$$h'(t) + 5h(t) = \delta(t) \quad (*)$$

$$h_h(t) = Ce^{\alpha t}$$

$$h'_h(t) = \alpha Ce^{\alpha t}$$

$$\alpha Ce^{\alpha t} + 5Ce^{\alpha t} = 0 \Rightarrow \alpha = -5 \Rightarrow h_h(t) = Ce^{-5t}$$

For $h_p(t)$, multiply both sides with dt and integrate between 0^- and 0^+ .

$$\int_{0^-}^{0^+} dh(t) + 5 \int_{0^-}^{0^+} h(t) dt = \int_{0^-}^{0^+} \delta(t) dt = 1$$

0 (continuity)

$h(0^-) = 0$ since the system is initially at rest. $\Rightarrow h(0^+) = 1$. Plug this into (*)

$$h(0^+) = Ce^{-5 \cdot 0} = C = 1 \Rightarrow h(t) = e^{-5t}u(t).$$

5 (26 pts, 13 pts each) Consider the following linear system:

$$y[n] = \sum_{k=-\infty}^{\infty} x[k]g[n-3k],$$

where $g[n] = u[n-1] - u[n-3]$.


- a) Find and plot $y[n]$ if $x[n] = \delta[n-1]$.
- b) Find and plot $y[n]$ if $x[n] = u[n-1]$.

a) Find $y[n]$ if $x[n] = \delta[n-1]$.

When you plug $x[n]$ into the equation above you will get a result which is in terms of $g[n]$. This comes from the sifting property.

$$y[n] = \sum_{k=-\infty}^{\infty} g[n-3k]\delta[k-1]$$

As you can see this equation is defined only when $k = 1$. Therefore we get the result $g[n-3]$ which is equal to $u[n-4] - u[n-6]$.




b) Find $y[n]$ if $x[n] = u[n-1]$.

When you plug $x[n]$ into the equation above you will get the following equation:

$$y[n] = \sum_{k=-\infty}^{\infty} g[n-3k]u[k-1]$$

Since $u[n-1]$ is defined on the interval of $[1, \infty)$ the formula simply becomes the following:

$$y[n] = \sum_{k=1}^{\infty} g[n-3k]$$



In the figure above you can see the values of $y[n]$ for the first 15 n values. From the figure we see that $y[n]$ will be $u[n-4] - \sum_{k=2}^{\infty} \delta[n-3k]$.