



Department of Computer Engineering

CENG 384 Homework Exam 2 Key

10 June 2022

METU Honor Code and Pledge

METU Honor Code: “The members of the METU community are reliable, responsible and honourable people who embrace only the success and recognition they deserve, and act with integrity in their use, evaluation and presentation of facts, data and documents.”

*“I have read and understood the implications of the METU Honor Code. To be precise, I understand that this is a **open** formula-sheet exam, and I am forbidden to access any other source of information other than provided within the exam. I will **TURN OFF** all my electronic equipment (phones, smart watches, etc.) and put it off the table along with other notes and materials that I may have with me. I understand that leaving electronic devices on during the exam is strictly forbidden. I understand and accept to obey all the rules announced by the course staff, and that failure to obey these will result in disciplinary action.”*

Name, SURNAME:
ID:

Signature:

Specifications About the Exam

- **Duration:** 110 minutes.
- **Notes:**
 - One A4-size double-sided formula-sheet is allowed.
 - Each answer should contain a justifying explanation. However, give shortest possible answers. Unnecessary or irrelevant wordings will reduce your grades.
 - If you use the back of a page for extra space, please mark down a proper notice on the relevant pages.

Grade

Q1:


Q2:

Q3:

Q4:

Total:

- 1 (25 pts, 5-5-7-8 pts respectively) Consider the following block diagram representation of a discrete time LTI system, where D represents the unit delay operator.



- Find the difference equation which represents this system.
- Find the frequency response, $H(e^{j\omega})$.
- Find and plot the impulse response, $h[n]$, using $H(e^{j\omega})$ you found in part b.
- Using Fourier transform, find and plot the output, $y[n]$, when the input is

$$x[n] = \delta[n] + \frac{1}{2}\delta[n-1].$$

a)

$$\frac{1}{2}x[n] - x[n-2] = y[n].$$

b)

$$\frac{1}{2}X(e^{j\omega}) - e^{-2j\omega}X(e^{j\omega}) = Y(e^{j\omega}),$$

$$H(e^{j\omega}) = \frac{Y(e^{j\omega})}{X(e^{j\omega})} = \frac{1}{2} - e^{-2j\omega}.$$

- c) Using table 5.2 we can take the IFT of $H(e^{j\omega})$ and reach $h[n]$:

$$h[n] = \frac{1}{2}\delta[n] - \delta[n-2].$$




Figure 1: n vs. $h[n]$.

d) Using table 5.2 we can take the FT of $x[n]$ and reach $X(e^{j\omega})$:

$$X(e^{j\omega}) = 1 + \frac{1}{2}e^{-j\omega},$$

$$Y(e^{j\omega}) = X(e^{j\omega})H(e^{j\omega}) = \frac{1}{2} - e^{-2j\omega} + \frac{1}{4}e^{-j\omega} - \frac{1}{2}e^{-3j\omega},$$

Using table 5.2 we can take the IFT of $Y(e^{j\omega})$ and reach $y[n]$:

$$y[n] = \frac{1}{2}\delta[n] + \frac{1}{4}\delta[n-1] - \delta[n-2] - \frac{1}{2}\delta[n-3].$$




Figure 2: n vs. $y[n]$.

- 2 (25 pts, 5-10-10 pts respectively) Consider a continuous time LTI system represented by the following frequency response,


$$H(j\omega) = \frac{j\omega}{2 + 3j\omega - \omega^2}.$$

- Find the differential equation which represents this system.
- Find a block diagram representation of this system using minimum number of adders, integrators and differentiators.
- Find and plot the impulse response, $h(t)$.

a)

$$\begin{aligned} \frac{Y(j\omega)}{X(j\omega)} &= \frac{j\omega}{2 + 3j\omega - \omega^2}, \\ ((j\omega)^2 + 3j\omega + 2)Y(j\omega) &= j\omega X(j\omega), \\ y''(t) + 3y'(t) + 2y(t) &= x'(t). \end{aligned}$$

- b) There are many alternatives. One such alternative is as follows:



- c) By partial fraction we get the following representation for $H(j\omega)$:

$$H(j\omega) = \frac{-1}{j\omega + 1} + \frac{2}{j\omega + 2},$$

Using table 4.2 we can take the IFT of $H(j\omega)$ and reach $h(t)$:

$$h(t) = (2e^{-2t} - e^{-t})u(t).$$




Figure 3: t vs. $h(t)$.

- 3** (30 pts, 10-5-5-10 pts respectively) Consider a discrete-time LTI system, represented by the following impulse response:

$$h[n] = (n-1)(0.3)^n u[n]$$

- a) Find the frequency response, $H(e^{j\omega})$, of this system.
b) Find the spectral coefficients, a_k , of the input

$$x[n] = \sin(\omega_0 n) + \sin(4\omega_0 n).$$

- c) Find the spectral coefficients, b_k , of the output for the input given in part (b).
d) Find the difference equation which represents this system.

a)

$$h[n] = (n+1)(0.3)^n u[n] - 2(0.3)^n u[n]$$

Using table 5.2 we can take the FT of $h[n]$ and reach $H(e^{j\omega})$:

$$H(e^{j\omega}) = \frac{1}{(1-0.3e^{-j\omega})^2} - \frac{2}{1-0.3e^{-j\omega}}.$$

- b) From Euler's Formula we get the following expansion for $x[n]$:

$$x[n] = \frac{e^{j\omega_0 n} - e^{-j\omega_0 n}}{2j} + \frac{e^{4j\omega_0 n} - e^{-4j\omega_0 n}}{2j},$$

$$a_1 = \frac{1}{2j} = \frac{-j}{2}, \quad a_{-1} = -\frac{1}{2j} = \frac{j}{2}, \quad a_4 = \frac{1}{2j} = \frac{-j}{2}, \quad a_{-4} = -\frac{1}{2j} = \frac{j}{2}.$$

- c) $b_1 = a_1 H(e^{j\omega_0}) = \frac{1}{2j} H(e^{j\omega_0}), \quad b_{-1} = a_{-1} H(e^{-j\omega_0}) = \frac{-1}{2j} H(e^{-j\omega_0}),$

$$b_4 = a_4 H(e^{4j\omega_0}) = \frac{1}{2j} H(e^{4j\omega_0}), \quad b_{-4} = a_{-4} H(e^{-4j\omega_0}) = \frac{-1}{2j} H(e^{-4j\omega_0}).$$

d)

$$H(e^{j\omega}) = \frac{Y(e^{j\omega})}{X(e^{j\omega})} = \frac{0.6e^{-j\omega} - 1}{0.09e^{-2j\omega} - 0.6e^{-j\omega} + 1},$$

$$(0.09e^{-2j\omega} - 0.6e^{-j\omega} + 1)Y(e^{j\omega}) = (0.6e^{-j\omega} - 1)X(e^{j\omega}),$$

$$0.09y[n-2] - 0.6y[n-1] + y[n] = 0.6x[n-1] - x[n].$$

- 4 (20 pts, 10 pts each) The output $y(t)$ of a causal LTI system is related to the input $x(t)$ by the equation:

$$\frac{dy(t)}{dt} + 9y(t) = \int_{-\infty}^{+\infty} x(\tau)z(t-\tau)d\tau - x(t)$$

where $z(t) = e^{-t}u(t) + 2\delta(t)$.

- a) Find the frequency response, $H(j\omega)$, of this system.
b) Determine the impulse response, $h(t)$, of the system.

a)

$$\begin{aligned}y'(t) + 9y(t) &= x(t) * z(t) - x(t), \\j\omega Y(j\omega) + 9Y(j\omega) &= X(j\omega)Z(j\omega) - X(j\omega), \\Z(j\omega) &= \frac{1}{1+j\omega} + 2 = \frac{3+2j\omega}{1+j\omega},\end{aligned}$$

Now plug $Z(j\omega)$ into the equation above:

$$\begin{aligned}(j\omega + 9)Y(j\omega) &= \left(\frac{3+2j\omega}{1+j\omega} - 1\right) X(j\omega), \\(j\omega + 9)Y(j\omega) &= \left(\frac{2+j\omega}{1+j\omega}\right) X(j\omega), \\H(j\omega) &= \frac{Y(j\omega)}{X(j\omega)} = \frac{2+j\omega}{(1+j\omega)(9+j\omega)}.\end{aligned}$$

- b) By partial fraction we get the following representation for $H(j\omega)$:

$$H(j\omega) = \frac{1/8}{j\omega + 1} + \frac{7/8}{j\omega + 9},$$

Using table 4.2 we can take the IFT of $H(j\omega)$ and reach $h(t)$:

$$h(t) = \left(\frac{1}{8}e^{-t} + \frac{7}{8}e^{-9t}\right)u(t).$$