

Winter-run refuge flows with reintroduction: modeled outcome levels across five hydroclimates, and how much of the difference from the baseline is the hydroclimate

A results brief for a general reader — prototype (v0_3), built by the brief builder from the locked 2026-08-29 API snapshot — reporting modeled outcome levels for one management strategy read across the five hydroclimates, each difference from the baseline split into the two effects that produce it. The build stamp, the run identifiers and the verification status are in references.

TL;DR. Compared with the baseline — current operations under the historical hydroclimate — this strategy’s modeled outcome levels differ from the baseline in all nine sectors of the water system under at least one hydroclimate. Reading each sector under each hydroclimate as one result, 45 in all, the hydroclimate effect is the larger of the two in 22, and the operations effect in 23. The strategy’s value differs from current operations at 98 of the 279 locations reported here under the historical hydroclimate and at 206 under extreme climate stress, and across the five hydroclimates the count ranges from 98 to 206; where it does not differ, the difference from the baseline is the hydroclimate alone. Freshwater for Delta exports is the clearest case of the two effects running opposite ways: under high climate stress the hydroclimate contributes +1.1 and the strategy −1.0, leaving a difference from the baseline of +0.1.

What this is

This is a *results brief*: it reports the modeled outcome levels, and nothing else, for the winter-run refuge flows with reintroduction strategy — one of 29 management strategies in the COEQWAL scenario library — simulated once under each of five hydroclimates and read as a set. The selection rule is that there is none: this brief reports every sector of the water system these model runs carry an outcome level for, across all five modeled hydroclimates, and no filter is applied to either. All nine sectors of the water system are reported under every hydroclimate. A scenario is a strategy paired with one hydroclimate; the five hydroclimates are a sample of stress levels, not a forecast of which will occur. For how the five were constructed, see the hydroclimates reference brief.

Against current operations, this strategy differs in one of the eight operational settings the scenario library records: Flows — Flows on the Sacramento River assigned to improve salmon survival; balanced against

allocation reductions for Shasta storage. For the full configuration, see the winter-run refuge flows with reintroduction strategy brief.

Winter-run refuge flows with reintroduction specifies the same water management actions as Winter-run refuge flows. The two strategies share the same CalSim 3 simulation of the water system — the same model run, not a duplicate. They differ in one action that is not a water management action: reintroduction of winter-run Chinook salmon to cold-water habitat upstream of Shasta Dam. Reintroduction enters the modeling through the winter-run life cycle model, not through CalSim 3, so modeled results differ from Winter-run refuge flows only for the winter-run salmon key outcome. Results for the other eight key outcomes are identical by construction.

What the model shows

Across the nine sectors these runs report, the modeled level values under the baseline — current operations under the historical hydroclimate — range from 1.7 to 4.2. Read across the five hydroclimates, the values this brief reports range from 1.3 to 4.1. Of the 45 results — one sector under one hydroclimate — 17 sit in a different level band from that sector's baseline value and 28 sit in the same band; in eight of the nine sectors the five values do not all sit in one band.

The figure below draws every one of those results beside the baseline, with the two effects that produce the difference shown as separate marks, so a reader can see how much of each difference the hydroclimate carries and how much the strategy does. The section that follows says what the two effects are before the figure draws them; what the level values themselves mean is set out in how to read the values.

The baseline, and the two effects

The baseline is current operations under the historical hydroclimate. It is the one reference this brief asks a reader to learn, and it is the same value in every column, because it does not depend on which hydroclimate a scenario pairs with.

Any difference between a scenario and the baseline comes from two places, and they add exactly:

The hydroclimate effect is current operations under that hydroclimate minus the baseline — the same operations, a different hydroclimate. **The operations effect** is this strategy minus current operations *under that same hydroclimate* — the only comparison that holds the hydroclimate fixed, and therefore the only one that isolates what the strategy does. Together they equal the difference from the baseline.

Outcome levels are reported as a value on a continuous axis from Level 1 to Level 4.99, with the four level bands as shaded zones; the whole number below any value is its band. A value that lands exactly on a whole number takes that band: 4.0 is Level 4, at the start of its zone rather than the top of Level 3. The level scale is ordered by definition, Level 1 through Level 4; a band is a category and a position on the axis is a modeled magnitude, neither is a judgment. The scale, the two thresholds and the two fields a level value comes in are set out in how to read the values.

A sector’s level value is the mean of the level values at its locations. The nine sectors differ in how many locations that is. Delta estuary ecology is reported at one location, Delta-Eastside Water. Winter-run salmon is reported at one location, Sacramento River at Keswick. Freshwater for Delta exports is reported at two locations, Harvey O. Banks Pumping Plant and C.W. "Bill" Jones Pumping Plant (formerly Tracy). Both carry the same level value in every scenario in this ensemble. Freshwater for in-Delta uses is reported at two locations, Emmaton and Jersey Point. Both carry the same level value in every scenario in this ensemble. The remaining five sectors are reported across 8 to 132 locations. On the figure, a square mark identifies the sectors reported at one or two locations. This page reports sectors, not individual locations; where a place brief has been published for one of a sector’s locations, that brief reports the location’s own values across the same scenarios.

All nine sectors, all five hydroclimates




Figure 1. Modeled level values for winter-run refuge flows with reintroduction and the two effects that produce their difference from the baseline, nine sectors of the water system × five hydroclimates. Lanes within each sector run left to right: Historical (H), Moderate (M), Moderate-high (MH), High (Hi), Extreme (E) climate stress. Vertical axis: continuous level value, Level 1 at top, level bands as shaded zones. Dashed level: the baseline. Solid level: this scenario. A short flat tick in place of an arrow is a change smaller than 0.1 on the level scale. Square marks are the sectors reported at one or two locations. Every value is in the table that follows, and the scale is set out in how to read the values.

The same 45 results as numbers

Each cell reads top to bottom: the hydroclimate effect, the operations effect, and the resulting value with its band. The two effects add to the difference between the scenario and the baseline. Outlined cells are those where the two run opposite ways. Each hydroclimate column carries its own address for citation, and its own slice of the companion data file.

Sector	The baseline	Historical	Moderate	Moderate-high	High	Extreme	Operations effect across the five
Community surface water	L1 1.7	L1 1.7	L1 1.8	L1 1.7	L1 1.9	L2 2.2	one direction -0.3 to 0.0
Agricultural revenue	L2 2.3	L2 2.5	L3 3.0	L3 3.0	L3 3.1	L3 3.5	one direction +0.2 to +0.6
Environmental flows	L3 3.7	L3 3.5	L3 3.5	L3 3.5	L3 3.5	L3 3.9	one direction -0.4 to -0.2
Delta estuary ecology SINGLE LEVEL VALUE	L3 3.2	L3 3.1	L2 2.9	L3 3.0	L3 3.1	L3 3.5	one direction -0.2 to 0.0
Winter-run salmon SINGLE LEVEL VALUE	L4 4.2	L3 3.5	L3 3.8	L3 3.6	L3 3.7	L4 4.1	one direction -1.1 to -0.1
Freshwater for Delta exports SINGLE LEVEL VALUE	L2 2.7	L1 1.3	L2 2.7	L2 2.3	L2 2.8	L3 3.3	one direction -1.4 to -0.4
Freshwater for in-Delta uses SINGLE LEVEL VALUE	L2 2.8	L2 2.3	L3 3.0	L2 2.6	L2 2.9	L3 3.4	one direction -0.8 to -0.1
Reservoir storage	L2 2.4	L2 2.4	L2 2.7	L2 2.2	L2 2.7	L3 3.4	one direction 0.0 to +0.3
Groundwater storage	L2 2.7	L2 2.8	L2 2.8	L2 2.9	L3 3.1	L3 3.2	one direction 0.0 to +0.2

How many places a strategy reaches

A strategy's rules apply in some places and not others, and a sector's value is the unweighted mean of its locations. This strategy's value differs from current operations by at least 0.1 on the level scale at 21–45 of the 74 Community surface water locations, 39–117 of the 132 Agricultural revenue locations, 6–10 of the 17 Environmental flows locations, the one Delta estuary ecology location under some hydroclimates and not others, the one Winter-run salmon location, all 2 Freshwater for Delta exports locations, all 2 Freshwater for in-Delta uses locations, as few as 4 and as many as all 8 Reservoir storage locations and 22–28 of the 42 Groundwater storage locations, depending on the hydroclimate — 98 to 206 of the 279 locations reported here. A sector where few locations differ therefore shows a small sector-level difference even when the locations that do differ move by a full level or more. Read across the five hydroclimates together, 739 of the 1,395 location-and-hydroclimate results reported here differ from current operations: 311 toward Level 1, 428 toward Level 4. The median difference over the locations counted runs from 0.1 on the level scale at Delta estuary ecology to 1.3 at Reservoir storage. The locations that differ do not all differ in the same direction: in 25 of the 45 sector-and-hydroclimate combinations reported here, some of a sector's locations differ from current operations toward Level 1 and others toward Level 4, so a sector value can sit close to the baseline while its locations differ in both directions. This page reports sectors, not individual locations; where a place brief has been published for one of a sector's locations, that brief reports the location's own values across the same scenarios.

Related briefs

What does this strategy do? It is described, rather than measured, by a reference brief that carries no modeled results: Winter-run refuge flows with reintroduction strategy brief.

What happens under a different strategy? The same nine sectors and the same five hydroclimates are reported once per strategy, and once with the strategy held fixed:

Current operations	Groundwater pumping limits in the Central Valley
Current operations with historical land use	Groundwater pumping limits in the San Joaquin Valley
Current operations with reintroduction	Groundwater pumping limits via reduced crop acreage in the Central Valley
Current operations without TUCPs	Groundwater pumping limits via reduced crop acreage in the San Joaquin Valley
Current USBR operations	Increase Delta outflows (55% unimpaired flow target)
Current USBR operations without TUCPs	Increase Delta outflows (65% unimpaired flow target)
Delta Conveyance Project	Increase Shasta carry-over storage
Functional environmental flows	Maintain Delta outflows (45% unimpaired flow target)
Functional environmental flows with reduced crop acreage	
Functional environmental flows with reduced crop acreage and reintroduction	
Functional environmental flows with reintroduction	

No flow requirements	The water system under current operations
Prioritizing full demands of community water systems	Winter-run refuge flows
Prioritizing human health water deliveries to community water systems	Winter-run refuge flows with reduced crop acreage
Reduce Delta outflows (35% unimpaired flow target)	Winter-run refuge flows with reduced crop acreage and reintroduction
Relax Delta salinity standards	Winter-run refuge flows with reintroduction — <i>you are here</i>

What happens to one sector under every strategy? A sector brief transposes this grid — one sector of the water system, every strategy, the same five hydroclimates:

Agricultural revenue	Freshwater for in-Delta uses
Community surface water	Groundwater storage
Delta estuary ecology	Reservoir storage
Environmental flows	Winter-run salmon
Freshwater for Delta exports	

How were the five hydroclimates constructed? The hydroclimates reference brief defines them. It reports no results of its own; every brief above is read against the same five.

What this doesn’t say

This brief reports modeled results from the COEQWAL ensemble. It does not predict what will happen, evaluate whether those results are acceptable, or recommend any course of action. The outcome-level framework provides definitional categories; the meaning of those categories for any individual reader, community, or decision is not specified by COEQWAL.

In particular, for this brief: the five hydroclimates are a sample of stress levels, carry no probabilities, and are not a timeline, so no average over them is reported here. Splitting a difference into two effects is arithmetic on modeled values, not a causal claim about the water system, and the two are separable only because each is defined against its own reference. A step or a spread on the level axis is a modeled magnitude, not a benefit or a cost. A sector’s value is the unweighted mean of its locations, so locations count equally regardless of the people, acres, or acre-feet each represents, and a location already at either end of the scale cannot move further along it. Locations are modeled nodes, not measured conditions at a place.

Extreme climate stress holds the single highest of a sector’s five values in all nine sectors reported here.

References, run identifiers and the build stamp

Model runs (this strategy / current operations): Historical `s0031-R` / `s0020`; Moderate `s0144-R` / `s0134`; Moderate-high `s0072-R` / `s0047`; High `s0092-R` / `s0056`; Extreme `s0118-R` / `s0108`. The baseline is current operations under the historical hydroclimate, `s0020`.

Build stamp. Snapshot `2026-08-29`; builder version `0.1.0`; built 2026-08-29T16:19:04Z; snapshot data read 2026-08-29; verification: pass. The verification status is written by the independent check (`briefbuilder.verifier`), which recomputes every value in the companion data file from the per-location data and reads every arrow on this page, after both files are written. It is not stamped by whatever produced them.

Companion data file: `results_winter-run-refuge-flows-with-reintroduction_schema1.json`, holding every value in this brief plus the per-location counts, the spread measures, the derivation of every number in the prose and the figure's declared attribute contract. Per-hydroclimate slices: Historical `#/rows/*/by_hydroclimate/BL`; Moderate `#/rows/*/by_hydroclimate/HC4`; Moderate-high `#/rows/*/by_hydroclimate/HC1`; High `#/rows/*/by_hydroclimate/HC2`; Extreme `#/rows/*/by_hydroclimate/HC3`. Built from the locked 2026-08-29 API snapshot (api.coeqwal.org v2.1.1 (locked dataset snapshot)). Decomposition per `Decomposition_Math_Quickref` (coeqwal-viz): the hydroclimate effect is current operations under that hydroclimate minus current operations under the historical hydroclimate; the operations effect is this strategy minus current operations under that same hydroclimate. Axis convention [1, 4.99] per `Ordinal_vs_continuous_tiers.md`; the visible band is `floor(value)`. A difference of 0.1 on the level scale is required before a difference is reported as movement, before a location is counted as differing from its comparison, and before an effect prints as anything other than 0.0. Color semantics (warm = hydroclimate, blue = operations) follow the ops-vs-climate stepper. Strategy identity is its descriptive name; run IDs appear only in this note and the data file.

Explore further

Every value on this page comes from the outcome-level data, and the data explorer holds the same data without this brief's selection of it. The link below opens winter-run refuge flows with reintroduction across all five hydroclimates and all nine sectors, so a reader can take the comparison apart, add a hydroclimate or a sector this brief reports as one number, and read the locations behind a sector value directly.

Open winter-run refuge flows with reintroduction in the data explorer.

How to read the values

The level scale runs from 1 to 4.99. Level 1 is the favorable end of the scale: the outcome-level framework names Level 1 Optimal and Level 4 Critical, so a smaller value sits nearer Optimal and a larger value nearer Critical. The API serves this number in a field named `weighted_score`; that name describes how the number is computed, and does not mean that a larger number is a more favorable result.

Two numbers describe each result. The integer `tier_level` is the level label. `tier_continuous` is the position within the band of that label, and always falls inside it: `tier_continuous` is at least `tier_level` and less than `tier_level` plus one. Labels, counts and prose use `tier_level`; figure positions and statistics use `tier_continuous`.

Level 1 is Optimal, Level 2 is Acceptable, Level 3 is At-risk, and Level 4 is Critical. These are definitional category names supplied by the outcome-level framework; what a category means for any reader, community, or decision is not specified by COEQWAL.

A sector value is the unweighted mean over the locations in that sector. Every location counts equally, regardless of the people, acres, or acre-feet it represents.

Any difference between two values under different hydroclimates contains both a hydroclimate effect and an operations effect; the two are separable only through the decomposition, because each is defined against its own reference. A strategy is compared with current operations under the same hydroclimate. Comparing one strategy with itself across hydroclimates is a different quantity: the hydroclimate effect plus the change in the operations effect.

One threshold governs every claim and every count here. A difference smaller than 0.1 on the level scale is not reported as movement, and a location is not counted as differing from its comparison. An outcome level is a summary built so that results in nine sectors with no shared unit can be compared, and 0.1 on the level scale is the finest reading it carries; a smaller difference is left unstated rather than described.

The five hydroclimates are a purposive sample of climate stress levels. They are not a sequence, a trend, or a forecast, they carry no probabilities, and no mean is taken over them.

A run ID identifies one modeling run of one strategy under one hydroclimate. It is an internal identifier, used for citation and for tracing a value back to the data platform, not a name for the strategy; the public identity of a strategy is its descriptive name.

Level values are reported to one decimal place throughout this brief — in the prose, in the figure and in the table. The level scale does not carry meaning finer than that. Each effect shown is the difference between the floored values, except that an effect smaller than 0.1 on the level scale, which is the threshold this brief reports movement at, is shown as 0.0, so a row containing one does not add up. Values are floored rather than rounded, so a printed value stays inside the band its own label names; full-precision values are in the machine-readable sidecar this brief is built from.