

# CS-UY 2413: Design & Analysis of Algorithms

Prof. Lisa Hellerstein

Fall 2024  
New York University

## Homework 2

Due 11:59pm Monday, Sep 30, New York time. By handing in the homework you are agreeing to the Homework Rules; see EdStem.

Our Master Theorem: The version of the Master Theorem that we covered in class is on the last page of this homework. We won't be covering the version of the Master Theorem in the textbook and you're not responsible for knowing it. (But you may find it interesting!)

Reminder: For  $r \neq 1$ ,  $r^0 + r^1 + \dots + r^k = \frac{r^{k+1}-1}{r-1}$ .

1. For each example, indicate whether  $f = o(g)$  (little-oh),  $f = \omega(g)$  (little-omega), or  $f = \Theta(g)$  (big-Theta). No justification is necessary.
  - (a)  $f(n) = n^2 \log n$ ,  $g(n) = n^3$
  - (b)  $f(n) = 2^n$ ,  $g(n) = n^2$
  - (c)  $f(n) = n \log n$ ,  $g(n) = n$
  - (d)  $f(n) = 7 \sum_{i=0}^n 2^i$ ,  $g(n) = 2^n$
  - (e)  $f(n) = \log_2 n$ ,  $g(n) = \log_{10} n$
2. Give a formal proof of the following statement: If  $f(n) \geq 1$  for all  $n \in \mathbb{N}$ ,  $g(n) \geq 1$  for all  $n \in \mathbb{N}$ ,  $f(n) = O(g(n))$ , and  $g(n)$  is unbounded (meaning  $\lim_{n \rightarrow \infty} g(n) = \infty$ ) then  $\sqrt{f(n)} = O(\sqrt{g(n)})$ . Use the formal definition of big-Oh in your answer. In your proof, you can use the fact that the value of  $\sqrt{n}$  increases as  $n$  increases.
3. For each of the following recurrences, determine whether Our Master Theorem (on the last page of this HW) can be applied to the recurrence. If it can, use it to give the solution to the recurrence in  $\Theta$  notation; no need to give any details. If not, write "Our Master Theorem does not apply."
  - (a)  $T(n) = 2T(n/2) + n \log n$
  - (b)  $T(n) = 9T(n/3) + n^2$
  - (c)  $T(n) = T(n/2) + 1$

4. Our Master Theorem can be applied to a recurrence of the form  $T(n) = aT(n/b) + n^d$ , where  $a, b, d$  are constants with  $a > 0, b > 1, d > 0$ . Consider instead a recurrence of the form  $T_{new}(n) = aT_{new}(n/b) + n \log_d n$  where  $a > 0, b > 1, d > 1$  (and  $T(1) = 1$ ). For each of the following, state whether the given property of  $T_{new}$  is true. If so, explain why it is true. If not, explain why it is not true. (Even if you know the version of the Master Theorem in the textbook, don't use it in your explanation.)
- (a)  $T_{new}(n) = O(n^2)$  if  $\log_b a < 1$
  - (b)  $T_{new}(n) = \Omega(n \log^2 n)$
5. Consider the recurrence  $T(n) = 2T(n/2) + n$  for  $n > 1$ , and  $T(1) = 1$ .
- (a) Compute the value of  $T(8)$ , using the recurrence. Show your work.
  - (b) Use a recursion tree to solve the recurrence and get a closed-form expression for  $T(n)$ , when  $n$  is a power of 2. (Check that your expression is correct by plugging in  $n = 8$  and comparing with your answer to (a).)
  - (c) Suppose that the base case is  $T(2) = 3$ , instead of  $T(1) = 1$ . What is the solution to the recurrence in this case, for  $n \geq 2$ ?
6. Consider a variation of mergesort that works as follows: If the array has size 1, return. Otherwise, divide the array into quarters, rather than in half. Recursively sort each quarter using this variation of mergesort. Then merge the first (leftmost) quarter with the second quarter. Then merge the result with the third quarter, and finally merge that result with the last quarter.
- (a) Write a recurrence for the running time of this variation of mergesort. It should be similar to the recurrence for ordinary mergesort. Assume  $n$  is a power of 4.
  - (b) Apply Our Master Theorem to the recurrence to get the running time of the algorithm, in theta notation. Show your work.
7. Consider the following recursive algorithm. Assume  $n$  is a power of 2. The algorithm takes as input an array  $A[1..n]$  of numbers.
- If  $n = 1$ , return.
  - Recursively call the algorithm on  $A[1..n/2]$ .
  - Recursively call the algorithm on  $A[n/2+1..n]$ .
  - Recursively call the algorithm on  $A[1..n/2]$  again.
- (a) Suppose the algorithm performs an operation that takes  $\Theta(n)$  time in each recursive call. Write a recurrence expressing the running time of the algorithm.

- (b) Apply Our Master Theorem to your recurrence. What is the running time of the algorithm, in theta notation?
  - (c) If the initial array A contains the sequence [1, 2, ..., n], describe the array after the first 3 recursive calls.
8. Which of the following statements is **FALSE**?
- (a) If  $f(n) = O(g(n))$ , then  $f(n) = \Theta(g(n))$ .
  - (b) If  $f(n) = O(g(n))$  and  $g(n) = O(h(n))$ , then  $f(n) = O(h(n))$ .
  - (c) If  $f(n) = \Theta(g(n))$ , then  $f(n) = O(g(n))$ .
  - (d) If  $f(n) = \Omega(g(n))$ , then  $g(n) = O(f(n))$ .

### Theorem 0.1 (Our Master Theorem)

Let  $a, b, d, n_0$  be constants such that  $a > 0$ ,  $b > 1$ ,  $d \geq 0$  and  $n_0 > 0$ . Let  $T(n) = aT(n/b) + \Theta(n^d)$  for when  $n \geq n_0$ , and  $T(n) = \Theta(1)$  when  $0 \leq n < n_0$ . Then,

$$T(n) = \begin{cases} \Theta(n^d \log n) & \text{if } d = \log_b a \\ \Theta(n^{\log_b a}) & \text{if } d < \log_b a \\ \Theta(n^d) & \text{if } d > \log_b a \end{cases}$$

We assume here that  $T(n)$  is a function defined on the natural numbers. We use  $aT(n/b)$  to mean  $a'T(\lfloor n/b \rfloor) + a''T(\lceil n/b \rceil)$  where  $a', a'' > 0$  such that  $a' + a'' = a$ .