
B cell
Cancer cell
Endothelial cell
Fibroblast
Myeloid cell
T cell
Epithelial
PVL
Mast cell
A Bassez B · E
Union = 1,249 cell type–task pairs
766
374
109
B Bassez B · NE
Union = 998 cell type–task pairs
533
289
176
C Shiao · R
Union = 1,417 cell type–task pairs
863
439
115
D Shiao · NR
Union = 1,289 cell type–task pairs
984
267
38
Intersection size
Set membership
scKNIFE
scCellFie
Shared
scKNIFE only
scCellFie only
E T cell · R/NR
ERAD / recognition of misfolded protein
Patient median normalized Method 5 activity (%)
R
NR
δ=+0.48, p=0.093, q=1
F Myeloid · E/NE
Glycogen degradation
BA: δ=-0.49, p=0.091, q=0.67
BB: δ=-0.50, p=0.67, q=1
G Bassez A · Myeloid
Pre→On · Glutathione synthesis
q=8.4e−15
q=5.7e−12
H Bassez B · T cell
Pre→On · Gluconeogenesis from glutamine
q=1.5e−65
q=4.8e−130
I Shiao · Fibroblast
Base→RTPD1 · Alanine synthesis from glutamine
q=3.4e−100
q=1.7e−75
J Shiao · Tumor
Base→PD1 · Bilirubin synthesis
q=0.296 (ns)
q=<1e−300
Patient medians are the biological replicate · pooled-cell Wilcoxon/BH values are descriptive · Bassez E/NE denotes clonotype expansion, not clinical response
Exact between-group Δ p=0.018
Exact between-group Δ p=0.1
Exact between-group Δ p=0.048
Exact between-group Δ p=0.042