

Proof of Individual Agent Opinion Convergence in a Weakly Connected Influence Graph Using Classic Update Function

Bernardo T. Amorim

bernardoamorim@dcc.ufmg.br

April 2020

Definition 1. The *classic update-function*, is defined as:

$$Bel_p^{t+1}(a_i|a_j) = Bel_p^t(a_i) + In(a_j, a_i)(Bel_p^t(a_j) - Bel_p^t(a_i)) \quad (1)$$

Definition 2. While the *overall classic update*, is defined as:

$$Bel_p^{t+1}(a_i) = \frac{1}{|A|} \sum_{a_j \in A} Bel_p^{t+1}(a_i|a_j) \quad (2)$$

Definition 3. We say a influence graph In over agents A is *weakly connected* if for all $a_i, a_j \in A$, there exist $a_{k_1}, a_{k_2}, \dots, a_{k_l} \subseteq A$ such that $In(a_i, a_{k_1}) > 0$, $In(a_{k_l}, a_j) > 0$, and for $m = 1, \dots, l-1$, $In(a_{k_m}, a_{k_{m+1}}) > 0$.

Definition 4. max_t and min_t are the maximum and minimum belief values in a given instant t , respectively.

To prove our conjecture, let's do some simplifications:

$$\begin{aligned} Bel_p^{t+1}(a_i) &= \frac{1}{|A|} \sum_{a_j \in A} Bel_p^{t+1}(a_i|a_j). \\ &= \frac{1}{|A|} \sum_{a_j \in A} (Bel_p^t(a_i) + In(a_j, a_i)(Bel_p^t(a_j) - Bel_p^t(a_i))) \\ &= Bel_p^t(a_i) + \frac{1}{|A|} \sum_{a_j \in A} In(a_j, a_i)(Bel_p^t(a_j) - Bel_p^t(a_i)) \end{aligned} \quad (3)$$

Since we have a finite number of beliefs and $\forall a_i \in A : Bel_p^t(a_i) \in [0, 1]$, there are always min_t and a max_t . We shall also note that, by the Squeeze Theorem, individual agent opinion converges to the same value if and only if $\lim_{t \rightarrow \infty} min_t = \lim_{t \rightarrow \infty} max_t$.

Thus, since we want to prove that it always converges, if $min_t = max_t$ we have nothing to prove, so assume $min_t \neq max_t$.

Lemma 1. *In a weakly connected graph and under classic belief update, if $\max_t \neq \min_t$:*

$$\forall a_i \in A : \min_t \leq Bel_p^{t+1}(a_i) \leq \max_t$$

Proof. By the equation 3:

$$Bel_p^{t+1}(a_i) = Bel_p^t(a_i) + \frac{1}{|A|} \sum_{a_j \in A} In(a_j, a_i)(Bel_p^t(a_j) - Bel_p^t(a_i))$$

Trying to maximize the right side, we can substitute $Bel_p^t(a_j)$ by \max_t , this turns our equation into an inequity, since $\forall a_j \in A, Bel_p^t(a_j) \leq \max_t$, by the definition of \max_t . That makes the terms inside the summation either equal or smaller than 0, thus:

$$\begin{aligned} Bel_p^{t+1}(a_i) &\leq Bel_p^t(a_i) + \frac{1}{|A|} \sum_{a_j \in A} In(a_j, a_i)(\max_t - Bel_p^t(a_i)) \\ &= Bel_p^t(a_i) + \frac{1}{|A|} \sum_{a_j \in A} (\max_t - Bel_p^t(a_i)) \quad (\text{since } In(a_j, a_i) \leq 1) \\ &= Bel_p^t(a_i) + \frac{|A|}{|A|} (\max_t - Bel_p^t(a_i)) \\ &= Bel_p^t(a_i) + \max_t - Bel_p^t(a_i) \\ Bel_p^{t+1}(a_i) &\leq \max_t \end{aligned} \tag{4}$$

Since a_i was arbitrary, the Lemma is true for all agents. The same reasoning can be used to show the equivalent property for \min_t \square

Corollary 1. *In a weakly connected influence graph and a classic update function, if $\min_t \neq \max_t$, then $\max_{t+1} \leq \max_t$ and $\min_{t+1} \geq \min_t$.*

Proof. The result of Lemma 1 tells us that all beliefs in the time $t + 1$ are either smaller than \max_t or equal to \max_t , thus, since \max_{t+1} must be one of those elements, $\max_{t+1} \leq \max_t$. And the same reasoning can be used for \min_t . \square

Corollary 2. $\lim_{t \rightarrow \infty} \max_t = U$ and $\lim_{t \rightarrow \infty} \min_t = L$ for some $U, L \in [0, 1]$.

Proof. Since both \max_t and \min_t are bounded between 0 and 1 by the definition of belief; and Lemma 1 showed us that they are monotonic, according to the Monotonic Convergence Theorem, the limits exist. \square

Now that we have those properties, our proof will follow by showing that an agent a_i that holds some belief $Bel_p^t(a_i)$ will influence every other agent by the time $t + |A| - 1$. To see this, we must open the definition of belief throughout time. But before we do this, let's jump into some small definitions and corollaries that will help us on the way.

Definition 5. Let's call the sequence $P(a_i|a_j) = (a_i, a_k, \dots, a_{k+l}, a_j)$ a *path* from a_i to a_j , if:

- All elements on the sequence are different.

- The first element in the sequence is a_i .
- The last element in the sequence is a_j .
- If a_n is the n 'th element in the sequence, if it has a successor a_{n+1} , $In(a_n, a_{n+1}) > 0$.

Note that many paths from a_i to a_j can exist, although our notation isn't enough to differentiate them. But in subsequent steps we will only need one of those paths, so the notation shouldn't be a problem.

Definition 6. Let's denote by $|P(a_i|a_j)|$ the size of a path from a_i to a_j , which we define as the number of elements in the sequence $P(a_i|a_j) - 1$.

Corollary 3. $|P(a_j|a_j)| \leq |A| - 1$.

Proof. This follows directly from the definition of path. Since it doesn't have repeated elements and we have $|A|$ agents, the path can't have more than $|A|$ elements, since the size of a path is defined as the number of elements minus one, the maximum size is $|A| - 1$. \square

Lemma 2. $\forall a_i, a_k \in A$ and $\forall n \geq 1$:

$$Bel_p^{t+n}(a_i) \leq max_t + \frac{1}{|A|} (In(a_k, a_i)(Bel_p^{t+n-1}(a_k) - max_t)) \quad (5)$$

Proof. By the Definitions 1 and 2:

$$\begin{aligned} Bel_p^{t+n}(a_i) &= \frac{1}{|A|} \sum_{a_j \in A} Bel_p^{t+n}(a_i|a_j) \\ Bel_p^{t+n}(a_i) &= \frac{1}{|A|} \sum_{a_j \in A} (Bel_p^{t+n-1}(a_i) + In(a_j, a_i)(Bel_p^{t+n-1}(a_j) - Bel_p^{t+n-1}(a_i))) \end{aligned}$$

Now note that the belief in the next time step is maximized when the belief of a_i itself is the maximum one possible. Given this property, we can replace $Bel_p^{t+n-1}(a_i)$ by max_{t+n-1} and thus turn our equation into an inequity:

$$\begin{aligned} Bel_p^{t+n}(a_i) &\leq \frac{1}{|A|} \sum_{a_j \in A} (max_{t+n-1} + In(a_j, a_i)(Bel_p^{t+n-1}(a_j) - max_{t+n-1})) \\ &= max_{t+n-1} + \frac{1}{|A|} \sum_{a_j \in A} In(a_j, a_i)(Bel_p^{t+n-1}(a_j) - max_{t+n-1}) \end{aligned}$$

To make our Lemma useful in future manipulations, we will take an arbitrary element a_k out of the summation:

$$\begin{aligned} Bel_p^{t+n}(a_i) &\leq max_{t+n-1} + \frac{1}{|A|} \sum_{a_j \in A \setminus \{a_k\}} (In(a_j, a_i)(Bel_p^{t+n-1}(a_j) - max_{t+n-1})) \\ &\quad + \frac{1}{|A|} (In(a_k, a_i)(Bel_p^{t+n-1}(a_k) - max_{t+n-1})) \end{aligned}$$

Since max_{t+n-1} is the greatest belief possible in that time step, the summation can be at most 0, thus:

$$Bel_p^{t+n}(a_i) \leq max_{t+n-1} + \frac{1}{|A|} (In(a_k, a_i) (Bel_p^{t+n-1}(a_k) - max_{t+n-1}))$$

Since max doesn't increase throughout time, $max_t \leq max_{t+n-1}$, thus:

$$Bel_p^{t+n}(a_i) \leq max_t + \frac{1}{|A|} (In(a_k, a_i) (Bel_p^{t+n-1}(a_k) - max_t))$$

□

Definition 7. Let's denote by In_{min} the smallest influence that's different from 0 in the influence graph.

Using the same notation we used in Corollary 2, let's call $\lim_{t \rightarrow \infty} max_t = U$ and $\lim_{t \rightarrow \infty} min_t = L$.

Now that we have all of these tools, let's jump to Lemma 3 which will be a tool in the most important part of the proof:

Calling a_k the agent who holds the belief min_t in the time t :

Lemma 3. $\forall a_i \in A : max_t - Bel_p^{t+|P(a_k|a_i)|}(a_i) \geq \epsilon$, with $\epsilon = \left(\frac{In_{min}}{|A|}\right)^{|P(a_k|a_i)|} \cdot (U - L)$.

Proof. By equation 3:

$$Bel_p^{t+|P(a_k|a_i)|}(a_i) = Bel_p^{t+|P(a_k|a_i)|-1}(a_i) + \frac{1}{|A|} \sum_{a_j \in A} Bel_p^{t+|P(a_k|a_i)|-1}(a_i|a_j)$$

What we will do now is separate, at each step, one element of the summation and apply Lemma 2 to modify our inequity. But we will be careful when choosing the elements we separate from the summation.

We will separate from the summation the elements in $P(a_k|a_i)$, starting from the end of the path until we get to a_k .

To simplify our notation, let's index the elements in the path from a_k to a_i , starting from the end of the path (since we are backtracking it will make more sense) by calling a_n the n^{th} element from the end to the beginning of the sequence.

Thus, by Lemma 2:

$$Bel_p^{t+|P(a_k|a_i)|}(a_i) \leq max_t + \frac{1}{|A|} In(a_1, a_i) (Bel_p^{t+|P(a_k|a_i)|-1}(a_1) - max_t)$$

Note, now, that if $|P(a_k, a_i)| = 1$, we could prove our result. Instead of showing it I will expand this two more times, show the general version and then prove the Lemma for all cases.

Using Lemma 2 again:

$$\begin{aligned}
& Bel_p^{t+|P(a_k|a_i)|}(a_i) \\
& \leq max_t + \frac{1}{|A|} In(a_1, a_i) (Bel_p^{t+|P(a_k|a_i)|-1}(a_1) - max_t) \\
& \leq max_t + \frac{1}{|A|} In(a_1, a_i) \left(\left(max_t + \frac{1}{|A|} In(a_2, a_1) (Bel_p^{t+|P(a_k|a_i)|-2}(a_2) - max_t) \right) - max_t \right) \\
& = max_t + \frac{1}{|A|} In(a_1, a_i) \left(\frac{1}{|A|} In(a_2, a_1) (Bel_p^{t+|P(a_k|a_i)|-2}(a_2) - max_t) \right) \\
& = max_t + \frac{1}{|A|^2} In(a_2, a_1) In(a_1, a_i) (Bel_p^{t+|P(a_k|a_i)|-2}(a_2) - max_t) \\
& \leq max_t + \frac{1}{|A|^2} In(a_2, a_1) In(a_1, a_i) \times \\
& \quad \left(\left(max_t + \frac{1}{|A|} In(a_3, a_2) (Bel_p^{t+|P(a_k|a_i)|-3}(a_3) - max_t) \right) - max_t \right) \\
& = max_t + \frac{1}{|A|^2} In(a_2, a_1) In(a_1, a_i) \left(\frac{1}{|A|} In(a_3, a_2) (Bel_p^{t+|P(a_k|a_i)|-3}(a_3) - max_t) \right) \\
& = max_t + \frac{1}{|A|^3} In(a_3, a_2) In(a_2, a_1) In(a_1, a_i) (Bel_p^{t+|P(a_k|a_i)|-3}(a_3) - max_t)
\end{aligned}$$

We can see a patten forming since the equation above has the same form of the one before it, and this pattern will continue throughout time. Now, denoting P_{In} the product of the influences in the path ($P_{In} = In(a_k, a_{|P(a_k|a_i)|}) \times \dots \times In(a_1, a_i)$), we can write the generalized version of the inequity above as:

$$\begin{aligned}
Bel_p^{t+|P(a_k|a_i)|}(a_i) & \leq max_t + \frac{P_{In}}{|A|^{|P(a_k|a_i)|}} \cdot (Bel_p^t(a_k) - max_t) \\
& = max_t + \frac{P_{In}}{|A|^{|P(a_k|a_i)|}} \cdot (min_t - max_t)
\end{aligned}$$

This inequity comes from the fact that the path ends after $|P(a_k|a_i)|$ steps with a_k as the start of the path, and, by definition, the belief of a_k in the time t is min_t .

Since the rightmost term in the equation is either equal to or smaller than 0, to make the inequality hold for all a_i 's, we shall substitute P_{In} by the smallest value possible. According to Definition 7, In_{min} is the smallest positive influence in the graph. By the definition of path (5) all influences are positive, thus the smallest possible value of P_{In} is $In_{min}^{|P(a_k|a_i)|}$. Thus:

$$Bel_p^{t+|P(a_k|a_i)|}(a_i) \leq max_t + \left(\frac{In_{min}}{|A|} \right)^{|P(a_k|a_i)|} \cdot (min_t - max_t)$$

According to Corollary 2, the maximum value of min_t is L and the minimum value of max_t is U , those are the values we should plug to maintain the inequity:

$$\begin{aligned}
Bel_p^{t+|P(a_k|a_i)|}(a_i) &\leq max_t + \left(\frac{In_{min}}{|A|}\right)^{|P(a_k|a_i)|} \cdot (L - U) \\
Bel_p^{t+|P(a_k|a_i)|}(a_i) - max_t &\leq \left(\frac{In_{min}}{|A|}\right)^{|P(a_k|a_i)|} \cdot (L - U) \\
max_t - Bel_p^{t+|P(a_k|a_i)|}(a_i) &\geq \left(\frac{In_{min}}{|A|}\right)^{|P(a_k|a_i)|} \cdot (U - L) \\
max_t - Bel_p^{t+|P(a_k|a_i)|}(a_i) &\geq \epsilon
\end{aligned} \tag{6}$$

□

Theorem 1. $\forall a_i \in A : max_t - Bel_p^{t+|A|-1}(a_i) \geq \epsilon$, with $\epsilon = \left(\frac{In_{min}}{|A|}\right)^{|A|-1} \cdot (U - L)$.

Proof. Let's keep the notation of the previous Lemma and call a_k the agent that holds the belief min_t in the time t .

First we should note that, if $|P(a_k|a_i)| = |A| - 1$, our theorem is true by Lemma 3 and we nothing to prove.

Else if $|P(a_k|a_i)| \neq |A| - 1$, then $|P(a_k|a_i)| < |A| - 1$ according to Corollary 3. What we will do then is manipulate the formula of belief until we can use Lemma 3 and then end our proof.

Before getting into the proof, I will introduce a new notation. Given the fact that the equations get pretty big, I will introduce a new notation. Let's call:

$$S^t(a_i) = \sum_{x_j \in A} In(a_j, a_i)(Bel_p^t(a_j) - Bel_p^t(a_i)) \tag{7}$$

Although it is a strange notation I believe it will make it easier to understand.

Now let's open the definition of belief throughout time and get the general form:

$$\begin{aligned}
Bel_p^{t+|A|-1}(a_i) &= Bel_p^{t+|A|-2}(a_i) + \frac{1}{|A|} \sum_{x_j \in A} In(a_j, a_i)(Bel_p^{t+|A|-2}(a_j) - Bel_p^{t+|A|-2}(a_i)) \\
&= Bel_p^{t+|A|-2}(a_i) + \frac{1}{|A|} \cdot S^{t+|A|-2}(a_i) \\
&= Bel_p^{t+|A|-3}(a_i) + \frac{1}{|A|} \cdot (S^{t+|A|-3}(a_i) + S^{t+|A|-2}(a_i)) \\
&= Bel_p^{t+|A|-4}(a_i) + \frac{1}{|A|} \cdot (S^{t+|A|-4}(a_i) + S^{t+|A|-3}(a_i) + S^{t+|A|-2}(a_i))
\end{aligned}$$

Generalizing:

$$Bel_p^{t+|A|-1}(a_i) = Bel_p^{t+|A|-1-n}(a_i) + \frac{1}{|A|} \sum_{l=t+|A|-1-x}^{t+|A|-2} S^l$$

Now, if we choose $n = |A| - |P(a_k|a_i)| - 1$ then we will be able to use Lemma 3, but before doing so let's separate $S^{t+|P(a_k|a_i)|}$ from the summation (note that the

first two terms in the following inequity are actually $Bel_p^{t+|P(a_k|a_i)|+1}(a_i)$, this will be important in the future):

$$\begin{aligned}
& Bel_p^{t+|A|-1}(a_i) \\
&= Bel_p^{t+|P(a_k|a_i)|}(a_i) + \frac{1}{|A|} \cdot S^{t+|P(a_k|a_i)|}(a_i) + \frac{1}{|A|} \sum_{l=t+|P(a_k|a_i)|+1}^{t+|A|-2} S^l(a_i) \\
&= Bel_p^{t+|P(a_k|a_i)|}(a_i) + \frac{1}{|A|} \sum_{a_j \in A} In(a_j, a_i) (Bel_p^{t+|P(a_k|a_i)|}(a_j) - Bel_p^{t+|P(a_k|a_i)|}(a_i)) + (8) \\
&\quad \frac{1}{|A|} \sum_{l=t+|P(a_k|a_i)|+1}^{t+|A|-2} S^l(a_i)
\end{aligned}$$

Note that, if we separate a_i from the first summation in the equality above, what comes out $(In(a_i, a_i)(Bel_p^{t+|P(a_k|a_i)|}(a_j) - Bel_p^{t+|P(a_k|a_i)|}(a_j)))$ equals 0, thus we can take it out of the summation without changing anything more:

$$\begin{aligned}
Bel_p^{t+|A|-1}(a_i) &= Bel_p^{t+|P(a_k|a_i)|}(a_i) + \frac{1}{|A|} \sum_{a_j \in A \setminus \{a_i\}} In(a_j, a_i) (Bel_p^{t+|P(a_k|a_i)|}(a_j) - Bel_p^{t+|P(a_k|a_i)|}(a_i)) \\
&\quad + \frac{1}{|A|} \sum_{n=t+|P(a_k|a_i)|+1}^{t+|A|-2} S^n(a_i)
\end{aligned}$$

Since $Bel_p^{t+|P(a_k|a_i)|}(a_j)$ is multiplied by positive things, we can replace it by max_t keeping the inequity:

$$\begin{aligned}
Bel_p^{t+|A|-1}(a_i) &\leq Bel_p^{t+|P(a_k|a_i)|}(a_i) + \frac{1}{|A|} \sum_{a_j \in A \setminus \{a_i\}} In(a_j, a_i) (max_t - Bel_p^{t+|P(a_k|a_i)|}(a_i)) \\
&\quad + \frac{1}{|A|} \sum_{n=t+|P(a_k|a_i)|+1}^{t+|A|-2} S^n(a_i)
\end{aligned}$$

According to the Equation 6 in Lemma 3:

$$Bel_p^{t+|P(a_k|a_i)|}(a_i) \leq max_t + \left(\frac{In_{min}}{|A|} \right)^{|P(a_k|a_i)|} \cdot (L - U)$$

To keep things simple, let's call $\delta = \left(\frac{In_{min}}{|A|} \right)^{|P(a_k|a_i)|} \cdot (L - U)$.

Using the argument that, the way to maximize the belief is to replace it by its maximum value possible (this is clearer looking at definition 1). We will replace $Bel_p^{t+|P(a_k|a_i)|}(a_i)$ by its upper bound, which is given by Lemma 3. Thus:

$$\begin{aligned}
Bel_p^{t+|A|-1}(a_i) &\leq max_t + \delta + \frac{1}{|A|} \sum_{a_j \in A \setminus \{a_i\}} In(a_j, a_i)(max_t - max_t - \delta) + \frac{1}{|A|} \sum_{n=t+|P(a_k|a_i)|+1}^{t+|A|-2} S^n(a_i) \\
&= max_t + \delta + \frac{1}{|A|} \sum_{a_j \in A \setminus \{a_i\}} In(a_j, a_i)(-\delta) + \frac{1}{|A|} \sum_{n=t+|P(a_k|a_i)|+1}^{t+|A|-2} S^n(a_i) \\
&\leq max_t + \delta + \frac{(|A| - 1)(In_{min})(-\delta)}{|A|} + \frac{1}{|A|} \sum_{n=t+|P(a_k|a_i)|+1}^{t+|A|-2} S^n(a_i) \\
&= max_t + \frac{\delta (|A| - In_{min}(|A| - 1))}{|A|} + \frac{1}{|A|} \sum_{n=t+|P(a_k|a_i)|+1}^{t+|A|-2} S^n(a_i) \\
&\leq max_t + \frac{\delta (In_{min})}{|A|} + \frac{1}{|A|} \sum_{n=t+|P(a_k|a_i)|+1}^{t+|A|-2} S^n(a_i)
\end{aligned}$$

Now you must remember that the terms we were manipulating in the last steps where actually $Bel_p^{t+|P(a_k|a_i)|+1}$, thus $max_t + \frac{\delta (|A|(1-In_{min})-In_{min})}{|A|}$ is an upper bound of $Bel_p^{t+|P(a_k|a_i)|+1}$. Having this in mind, I will repeat the step of separating one term from the summation so we can see the pattern:

$$\begin{aligned}
Bel_p^{t+|A|-1}(a_i) &\leq max_t + \frac{\delta (In_{min})}{|A|} + \frac{1}{|A|} \sum_{n=t+|P(a_k|a_i)|+1}^{t+|A|-2} S^n(a_i) \\
&= max_t + \frac{\delta (In_{min})}{|A|} + \frac{1}{|A|} S^{t+|P(a_k|a_i)|+1}(a_i) + \frac{1}{|A|} \sum_{n=t+|P(a_k|a_i)|+2}^{t+|A|-2} S^n(a_i) \\
&= max_t + \frac{\delta (In_{min})}{|A|} \\
&\quad + \frac{1}{|A|} \sum_{a_j \in A} (In(a_j, a_i) (Bel_p^{t+|P(a_k|a_i)|+1}(a_j) - Bel_p^{t+|P(a_k|a_i)|+1}(a_i)) \\
&\quad + \frac{1}{|A|} \sum_{n=t+|P(a_k|a_i)|+2}^{t+|A|-2} S^n(a_i) \\
&= max_t + \frac{\delta (In_{min})}{|A|} \\
&\quad + \frac{1}{|A|} \sum_{a_j \in A \setminus \{a_i\}} In(a_j, a_i) (Bel_p^{t+|P(a_k|a_i)|+1}(a_j) - Bel_p^{t+|P(a_k|a_i)|+1}(a_i)) \\
&\quad + \frac{1}{|A|} \sum_{n=t+|P(a_k|a_i)|+2}^{t+|A|-2} S^n(a_i) \\
&\leq max_t + \frac{\delta (In_{min})}{|A|} \\
&\quad + \frac{1}{|A|} \sum_{a_j \in A \setminus \{a_i\}} In(a_j, a_i) (max_t - Bel_p^{t+|P(a_k|a_i)|+1}(a_i)) + \frac{1}{|A|} \sum_{n=t+|P(a_k|a_i)|+2}^{t+|A|-2} S^n(a_i)
\end{aligned}$$

Since $Bel_p^{t+|P(a_k|a_i)|+1} \leq max_t + \frac{\delta (In_{min})}{|A|}$:

$$\begin{aligned}
Bel_p^{t+|A|-1}(a_i) &\leq max_t + \frac{\delta (In_{min})}{|A|} \\
&\quad + \frac{1}{|A|} \sum_{a_j \in A \setminus \{a_i\}} \left(In(a_j, a_i) \left(max_t - max_t - \frac{\delta (In_{min})}{|A|} \right) \right) \\
&\quad + \frac{1}{|A|} \sum_{n=t+|P(a_k|a_i)|+2}^{t+|A|-2} S^n(a_i) \\
&= max_t + \frac{\delta (In_{min})}{|A|} \\
&\quad + \frac{1}{|A|} \sum_{a_j \in A \setminus \{a_i\}} \left(In(a_j, a_i) \left(-\frac{\delta (In_{min})}{|A|} \right) \right) \\
&\quad + \frac{1}{|A|} \sum_{n=t+|P(a_k|a_i)|+2}^{t+|A|-2} S^n(a_i) \\
&= max_t + \frac{\delta (In_{min})}{|A|} - \frac{(|A|-1)(In_{min}) (\delta (In_{min}))}{|A|^2} \\
&\quad + \frac{1}{|A|} \sum_{n=t+|P(a_k|a_i)|+2}^{t+|A|-2} S^n(a_i) \\
&= max_t + \frac{\delta (|A|)(In_{min}) - \delta (|A|-1)(In_{min})^2}{|A|^2} + \frac{1}{|A|} \sum_{n=t+|P(a_k|a_i)|+2}^{t+|A|-2} S^n \\
&\leq max_t + \frac{\delta (|A|)(In_{min})^2 - \delta (|A|-1)(In_{min})^2}{|A|^2} + \frac{1}{|A|} \sum_{n=t+|P(a_k|a_i)|+2}^{t+|A|-2} S^n(a_i) \\
&= max_t + \delta \left(\frac{In_{min}}{|A|} \right)^2 + \frac{1}{|A|} \sum_{n=t+|P(a_k|a_i)|+2}^{t+|A|-2} S^n(a_i)
\end{aligned} \tag{9}$$

Although I know it is not convincing to show a pattern and claim that it continues, I don't know how to show it in a better way. Anyway, my claim is that we can repeat the procedure of separating one S from the summation, replace the beliefs inside the summation by max_t and by the upper bound of the last belief we analyzed and then we will end up with:

$$\begin{aligned}
Bel_p^{t+|A|-1}(a_i) &\leq max_t + \delta \left(\frac{In_{min}}{|A|} \right)^{|A|-|P(a_k|a_i)|-1} \\
&= max_t + \left(\frac{In_{min}}{|A|} \right)^{|P(a_k|a_i)|} \left(\frac{In_{min}}{|A|} \right)^{|A|-|P(a_k|a_i)|-1} .(L - U) \\
&= max_t + \left(\frac{In_{min}}{|A|} \right)^{|A|-1} .(L - U) \\
Bel_p^{t+|A|-1}(a_i) - max_t &\leq \left(\frac{In_{min}}{|A|} \right)^{|A|-1} .(L - U) \\
max_t - Bel_p^{t+|A|-1}(a_i) &\geq \left(\frac{In_{min}}{|A|} \right)^{|A|-1} .(U - L) \\
max_t - Bel_p^{t+|A|-1}(a_i) &\geq \epsilon
\end{aligned}$$

□

Corollary 4. $max_t - max_{t+|A|-1} \geq \epsilon$

Proof. Since $max_{t+|A|-1}$ must be one of the beliefs in the time $t+|A|-1$ and, according to Theorem 1 all of them are smaller than max_t by a factor of at least ϵ , $max_{t+|A|-1}$ must also be smaller than max_t by a factor of at least ϵ . □

Theorem 2. $\lim_{t \rightarrow \infty} max_t = U = \lim_{t \rightarrow \infty} min_t = L$

Proof. Suppose, by contradiction, that $U \neq L$. Plugging this values into the ϵ formula show us that $\epsilon \neq 0$. Since, according to Theorem 1, $max_{t+|A|-1}$ is smaller than max_t by a factor of ϵ , we can finally reach to a contradiction and end our proof.

To see this contradiction, let's assume we did $v = (|A| - 1) \left(\lceil \frac{1}{\epsilon} \rceil + 1 \right)$ time steps after $t = 0$. Since max diminishes by at least ϵ at each $|A| - 1$ steps:

$$\begin{aligned}
max_0 &\geq max_v + \epsilon \left(\left\lceil \frac{1}{\epsilon} \right\rceil + 1 \right) \\
max_0 - \epsilon \left(\left\lceil \frac{1}{\epsilon} \right\rceil + 1 \right) &\geq max_v
\end{aligned}$$

But $\epsilon \cdot \left(\left\lceil \frac{1}{\epsilon} \right\rceil + 1 \right) > 1$, thus $max_0 < \epsilon \cdot \left(\left\lceil \frac{1}{\epsilon} \right\rceil + 1 \right)$. And this would imply that $max_v < 0$, which contradicts the definition of belief!

Since assuming that $U \neq L$ led us to a contradiction we can conclude that $U = L$. This result implies that all agents belief converge to the same value, as we wanted to prove. □

TEMPORARY

Lemma 4. If $Bel_p^t(a_i) \leq x$:

$$Bel_p^t(a_i) \leq x + \frac{1}{|A|} \sum_{a_j \in A} (In(a_j, a_i) (Bel_p^t(a_j) - x))$$