

Limbic system

Bernd Porr*, Ailsa Millen, Graeme Hattan, Maria Thomson

Version

1 Introduction

This page is a hybrid of an open ended circuit diagram with hyperlinks which point to the wiki underneath which in turn points to references which back up the connections. Think of google maps meets wikipedia.

The main aim is to have a circuit diagram which is supported by references which then informs computer models which simulate behavioural experiments.

1.1 Review articles

- This article by Castro and Berridge promotes their long standing differentiation between 'wanting' and 'liking' [Berridge, 2009] now updated by 'hedonic hot spots' [Castro et al., 2015].
- Berthoud's article features fantastic connection diagrams between the different nuclei of the limbic system [Berthoud, 2004].
- Zahm again provides very detailed anatomical connection diagrams and is a must read [Zahm, 2000].
- The review by Abbas Khani and Gregor Rainer focusses on the roles of the different nuclei, how they interact, how they implement reinforcement guided decision making and which forms are out there (reversal, go no go etc). A very balanced article which, for example, not just cites the prediction error paradigm of DA but reports also the other roles DA [Khani and Rainer, 2016].

*BerndPorr@glasgow.ac.uk

Contents

1	Introduction	1
1.1	Review articles	1
2	Amygdala	4
2.1	Central / Medial	4
2.2	Basolateral / lateral	4
2.3	Efferents	4
2.4	Afferents	4
3	Dorsal Raphe Nucleus (DRN)	5
3.1	Afferents	5
3.1.1	OFC to DRN	5
3.1.2	mPFC to DRN	5
3.1.3	LHb to DRN	5
3.1.4	LH to DRN	6
3.1.5	Amygdala to DRN	6
3.1.6	Basal Ganglia to DRN	6
3.1.7	Anterior cingulate cortex (ACA)	6
3.2	Efferents	6
3.2.1	Dorsal pathway of the DRN	7
3.2.2	Medial pathway of the DRN	7
3.2.3	Ventral pathway of the DRN	7
3.3	Receptors	7
3.3.1	5HTR1	7
3.3.2	5HTR2	8
3.3.3	5HTR3	8
3.3.4	5HTR4	9
3.4	Activity	9
3.5	Behavioural studies	9
4	Entopeduncular Nucleus (EP)	10
4.1	Efferents	10
4.2	Function	10
5	Lateral Habenula	10
5.1	Afferents	10
5.2	Efferents	11
5.3	Function	11

6	Lateral hypothalamus (LH)	11
6.1	Efferents	11
6.2	Afferents	12
6.3	Behavioural experiments	12
7	Nucleus Accumbens Core	12
7.1	Afferents	12
7.2	Efferents	12
7.2.1	Direct pathway	12
7.2.2	Indirect pathway	13
8	Nucleus Accumbens Shell	13
8.1	Medial Shell	13
8.2	Lateral Shell	13
8.3	Behavioural experiments	13
8.4	Function	13
9	Nucleus Accumbens (NAcc)	14
9.1	Behavioural experiments	14
9.2	Signals	14
9.3	Plasticity	14
10	Orbitofrontal Cortex (OFC)	15
10.1	Afferents	15
10.2	Efferents	15
10.3	Neuronal activity	16
10.4	Behavioural experiments	16
10.4.1	Medial Prefrontal Cortex (mPFC)	16
10.5	Efferents	16
10.6	Neuromodulation	16
10.7	Behaviour	17
11	Rostral Medial Tegmental Nucleus (RMTg)	17
11.1	Afferents	17
11.2	Efferents	17
12	Ventral Pallidum (VP)	17
12.1	Afferents	18
12.2	Efferents	18

13 Ventral Tegmental Area (VTA)	18
13.1 Afferents	18
13.2 Efferents	18
13.3 Function	19
13.3.1 Phasic activity	19
13.3.2 Tonic activity (DA)	20
13.3.3 Tonic activity (GABA)	20
A License	20

2 Amygdala

The connectivity of the amygdala has been outlined in these reviews [Alheid, 2003], [Sah et al., 2003] and [Swanson and Petrovich, 1998].

2.1 Central / Medial

For the the central and medial Amygdala [Swanson, 2003][Swanson and Petrovich, 1998] have pointed out that these parts contain cells which are similar to those in the striatum (i.e. GABAergic) and thus can be seen as an extension of the striatum dealing with lower executive or autonomic functions.

2.2 Basolateral / lateral

The basolateral part of the Amygdala which is related to reward processing and has glutamatergic neurons which have similar roles to that of the cortex and then projects into the striatal parts of the amygdala. See Fig 5 and 6 in [Swanson and Petrovich, 1998] for very informative connection diagrams.

2.3 Efferents

The anterior basolateral amygdala projects to the NAcc core and the posterior part to the extended amygdala structures [Alheid, 2003].

A projection from the Amygdala to the DRN [Pollak Dorocic et al., 2014] has been reported.

2.4 Afferents

The basolateral Amygdala has abundant projections from somatosensory cortices [Swanson and Petrovich, 1998], limbic cortices [Ottersen, 1982] and (ventral) hippocampus [Pitkänen et al., 2000].

mPFC inputs are inhibitory in nature [Rosenkranz and Grace, 2002] but this inhibition can be suppressed by DA release so that in the presence of DA the cortex can drive the BLA.

3 Dorsal Raphe Nucleus (DRN)

Review papers by Michelsen and Schmitz [Michelsen et al., 2007] and [Nakamura, 2013].

The DRN seems to generate 5HT to wait to obtain a reward behaviour [Nakamura, 2013].

3.1 Afferents

A detailed tracing / optogenetic study can be found here [Pollak Dorocic et al., 2014].

LHb, mPFC and LH appear to be the main afferents to the DRN [Vertes et al., 2010] [Sparta and Stuber, 2014] [Lee et al., 2003].

3.1.1 OFC to DRN

The OFC has strong reciprocal connections to with DRN [Zhou et al., 2015] where the OFC is probably the main nucleus of the DRN's ability to track the long term anticipated reward and reversal learning [Roberts, 2011].

3.1.2 mPFC to DRN

The mPFC (in particular its ventral part) has GLU projections to the DRN [Gonçalves et al., 2009] [Lee et al., 2003].

The conventional view is that the mPFCs glutamergic projections to the DRN connect to locally inhibitory neurons that then target 5HT neurons [Celada et al., 2001]. Stimulation of mPFC neurons usually inhibit DRN neurons which makes a strong case for these scenario.

However, [Pollak Dorocic et al., 2014] found that the mPFC has direct excitatory control of 5HT which they consider to be potentially critical for the correct function of the serotonergic system.

3.1.3 LHb to DRN

Similarly with regards to LHb projections to the DRN the conventional view is that the LHb neurons target local GABA neurons that then inhibit 5HT. However, [Pollak Dorocic et al., 2014] found a direct connection to the

DRN. Contrary to [Pollak Dorocic et al., 2014], [Ogawa et al., 2014] found few monosynaptic connections from the LHb to the DRN and instead posits that the LHb inhibits DRN 5HT via the rostral medial tegmental nucleus (RMTg). This has also been confirmed by [Sego et al., 2014].

Overall the picture emerges that the LHb exerts its influence via the RMTg and that direct connections from the LHb to the DRN are rare.

3.1.4 LH to DRN

Strong monosynaptic glutamatergic projections have been shown by [Lee et al., 2003] and [Aghajanian et al., 1990]. It's interesting that of these most prominent projections mentioned above it appears that this is the only excitatory one.

3.1.5 Amygdala to DRN

Tracing studies have shown robust projections from the Amygdala (CEA) to the DRN [Pollak Dorocic et al., 2014] which are monosynaptic and are most likely glutamatergic [Swanson and Petrovich, 1998]. In contrast to the GABAergic inputs above this seems to be one of the few excitatory inputs.

3.1.6 Basal Ganglia to DRN

Several nuclei from the basal ganglia project to the DRN including SNr and globus pallidus which is inhibitory in nature [Pollak Dorocic et al., 2014].

3.1.7 Anterior cingulate cortex (ACA)

The paper by [Pollak Dorocic et al., 2014] has also shown an excitatory pathway from the ACA to the DR.

3.2 Efferents

Most recently the projections to the limbic cortices have been identified as the strongest [Linley et al., 2013] [Roberts, 2011] whereas in this older publication [Reisine et al., 1984] states that the DRN projects to the striatum and caudate nucleus.

The paper by [Vertes et al., 2010] claims the DRN efferents include the VTA, SNc, LH and NAcc core. [Nakamura, 2013] stresses the projections from the DRN to the SNr, the VTA (inhibitory), amygdala, cortex (with inhibition of the mPFC) and to the NAcc where 5HT has at least partially a disinhibitory effect by targeting interneurons.

The actual effect of 5HT depends on the prominent receptor type in the target area. Some 5HT receptors are excitatory, some inhibitory and some ramp up plasticity [Frazer and Hensler, 1999].

The review paper by [Michelsen et al., 2007] makes a distinction between dorsal, medial and ventral pathways:

3.2.1 Dorsal pathway of the DRN

* all parts of the striatum ranging from the Nacc shell over core to the dorsal striatum * to a lesser extent the globus pallidus (GP)

3.2.2 Medial pathway of the DRN

* mainly the SNr.

3.2.3 Ventral pathway of the DRN

This pathway targets a large number of different limbic nuclei. In order of density:

* Septum (dense) * Amygdala (dense) * Habenula (dense) * Piriform, insular and frontal cortices (dense) * Occipital, entorhinal, perirhinal, frontal orbital, anterior cingulate and infralimbic cortices (moderate) * Thalamic and hypothalamic nuclei (dense to moderate) * Olfactory bulb. [Lottem et al., 2016] has shown that the spontaneous activity of the olfactory cortex is suppressed by 5HT release but that odor evoked activity is unaffected by 5HT.
* Hippocampus * Interpeduncular nucleus * Geniculate body

3.3 Receptors

In contrast to dopamine 5HT has a vast array of different receptors. While the DRN and MRN generate a global 5HT signal the effects on different brain areas can be vastly different because every brain area has their own 5HT receptor distribution [Palacios et al., 1990] [Carhart-Harris and Nutt, 2017]. Before we go into the efferents we present the different receptors and where they are located. If not otherwise cited it's based on the review by [Mengod et al., 2010] and the classic [Palacios et al., 1990].

3.3.1 5HTR1

The 5HTR1 has numerous subclasses: * 5HTR1A receptors are found on excitatory/pyramidal neurons and inhibit those. This receptor has been called the "limbic" receptor because it is prominent in the limbic areas of

the brain: hippocampus, lateral septum, cortex (cingulate/entorhinal) and raphe nucleus. These receptors are often co-expressed with the excitatory 5HT_{2A} receptor. They are located on the soma or dendrite and thus can inhibit the firing of neurons [Riad et al., 2000]. * 5HTR_{1B} are found mainly on inhibitory neurons and inhibit those but occasionally also on pyramidal neurons. They are very prominent in the basal ganglia, in particular in the GP, SNr, VP and EP (which are the output nuclei of the BG). They are both auto and heterosynaptic receptors and are located at the axon terminals [Riad et al., 2000] and control rather the release of transmitter in contrast to 5HTR_{1A} which control spiking. * 5HTR_{1D} are located in the caudate putamen, Nacc, olfactory cortex, dorsal raphe nucleus und locus coeruleus. It's predominantly located on axon terminals of both 5HT and non 5HT neurons and inhibit release of neurotransmitters. * 5HTR_{1E} is prominent in the (entorhinal) cortex, caudate putamen and claustrum. * 5HTR_{1F} has its highest levels in the cortical regions, olfactory bulb, Nacc, parascicular nucleus, thalamus, medial mamillary nucleus, hippocampus, subiculum and amygdala.

Having both inhibitory 5HT receptors on both excitatory and inhibitory neurons means that this can cancel out in average but will possibly change the dynamics of the network.

3.3.2 5HTR₂

* 5HTR_{2A}: These receptors are excitatory and enhancing inputs when activated meaning they are located on dendrites and on the soma. These receptors are very prominent in the cortex and have been localised on GABAergic interneurons but also glutamatergic projection neurons. * 5HTR_{2B}: it's function and localisation is still poorly understood * 5HTR_{2C}: has only been found in the CNS and there in the choroid plexus, cortex, NAcc, hippocampus, amygdala, caudate and SNr. They are also postsynaptic but might be also presynaptic.

3.3.3 5HTR₃

It's highest concentration is in the dorsal vagal complex of the brainstem.

Is a fast excitatory receptor which is mainly located in the hippocampus (and possibly amygdala) [Palacios et al., 1990] and is co-expressed with GLU receptors in the hippocampus.

Some evidence points to it controlling DA release [Mengod et al., 2010].

3.3.4 5HTR4

This receptor seems to control primarily plasticity, for both LTP and LTD [Peñas-Cazorla and Vilaró, 2015]. In an experiment by [Mlinar et al., 2006] stimulation of this receptor causes LTP in hippocampal slices which were very long lasting for over 2hrs.

Even more impressive are the findings by [Hagena and Manahan-Vaughan, 2017] who show that 5HTR4 activation shifts the frequency threshold between LTD and LTP: it is generally accepted that under LFS LTD is induced whereas under HFS LTP is induced. The frequency threshold where LTD turns into LTP can be shifted by the 5HTR4 receptor. If this receptor is stimulated even lower frequencies can cause LTP which otherwise would have caused LTD!

In the rat it is located mainly in the limbic system: hippocampus, striatum, inferior colliculus, SNr, ventral pallidum, fundus striatae, olfactory tubercle, septum and amygdala. It has also high concentrations in the parietal cortex.

3.4 Activity

As shown by [Li et al., 2016]: the activity increases when "when a mouse voluntarily seeks and acquires sucrose, food, sex and social interaction". 5HT neurons are activated by surprising reward events (such as VTA neurons do) and reward predicting cues is presented. The activity only drops off after the reward has been experienced. In particular the DRN activity stays active while the animal is waiting for a reward.

3.5 Behavioural studies

Numerous studies have reported that 5HT is required for delayed reward scenarios, for example where a rat has to wait in front of a dispenser to retrieve a reward [Khani and Rainer, 2016] as already mentioned above where the activity was measured during a delayed reward scenario [Li et al., 2016].

This has been combined into the proposal by [?] that 5HT controls patience and reward.

Premature responding is increased after [Fletcher et al., 2007] application of a 5HT(2A) receptor antagonist and decreased after 5-HT(2C) application. In earlier studies impulsivity could also be increased by 5HT depletion [Harrison et al., 1997].

4 Entopeduncular Nucleus (EP)

The entopeduncular nucleus, also called the GPi in primates, actions are strongly regulated by expected reward outcomes and it is an important output centre of the basal ganglia [Rajakumar et al., 1993]

4.1 Efferents

Different parts of the EP appear to perform different tasks.

* The caudal part of the EP has efferents to the (vm-)thalamus and brainstem nuclei and therefore controls motor activity [Rajakumar et al., 1993] [Wallace et al., 2017]. * The rostral part of the EP has strong projections to the lateral subnuclei of the LHb (including the oval nucleus) LHb [Rajakumar et al., 1993][Hong and Hikosaka, 2008] which are glutamatergic and excitatory [Shabel et al., 2012] [Wallace et al., 2017]. The LHb then projects primarily to the rostral medial tegmental area (rMTg) which is a GABAergic nucleus that innervates the VTA.

4.2 Function

The reward prediction error is considered to be instigated from the LHb and Barrot et al, 2012 theorized that the EP could play a major role in this activity as it has such an important input to the LHb [Barrot et al., 2012] and in turn to the VTA as mentioned above. This relates to the rostral part of the EP.

5 Lateral Habenula

5.1 Afferents

The 2 strongest forebrain afferents to the LHb are the EP and LH and other connections include the lateral preoptic area and the VP [Parent et al., 1981].

Araki, 1984, states that the connection from EP to LHb is GABAergic [Araki et al., 1984] but Shabel 2012 qualifies it as excitatory and glutamergic [Shabel et al., 2012].

Mok proposed that the actual connection was mainly excitatory [Mok and Mogenson, 1974]. A view supported by Poller who found a strong glutamergic projection that targeted VTA and RMTg projecting neurons [Poller et al., 2013].

The LHb also receives DA input from the midbrain VTA and SNc [Kowski et al., 2009] and reciprocally its main, inhibitory projections are to the VTA, SNc

and the DRN [Ji and Shepard, 2007][Christoph et al., 1986][Rajakumar et al., 1993].

5.2 Efferents

The main targets of the LHb are the VTA, Midbrain reticular nucleus (MRN), periaqueductal gray (PAG) and the superior central nucleus raphe (CS) [Quina et al., 2015].

Slightly puzzling is that the effect on the VTA by the LHb is known to be inhibitory. Hong et al. 2011 theorize that as LHb neurons are largely glutamergic their inhibitory function must be through an intermediary, the RMTg [Hong et al., 2011].

5.3 Function

Bromberg-Martin et al. 2010 research viewed LHb as the most important source of reward memory in DA neurons. They cited 3 potential pathways for transmission; 1) prefrontal cortex to the striatum to the global pallidus to LHb 2) For rats, the mPFC to LHb 3) common sources project reward memory signals to LHb and DA neurons. They found that many LHb and DA neurons signalled past reward results in their tonic activity, this was surprising as previous studies had reported this was purely the case for phasic activity [Bromberg-Martin et al., 2010].

It is considered to be the main source of the negative reward signal that facilitates the DA reward prediction error as LHb innervation inhibits midbrain DA. [Shen et al., 2012][Shabel et al., 2012][Matsumoto and Hikosaka, 2007][Barrot et al., 2012]

6 Lateral hypothalamus (LH)

The lateral hypothalamus has both Glutamatergic and GABAergic neurons [Stanley et al., 2011].

6.1 Efferents

The strongest outputs from the LH are to the VTA and the lateral habenula (LHb) [Stuber and Wise, 2016]. The projection from the LH to the Habenula is excitatory [Poller et al., 2013] which in turn then projects to the RMTg which has GABA-ergic neurons.

The LH contains both Glu and GABA neurons where the GLU neurons project to the DA neurons in the VTA whereas the GABA neurons in the LH project to the GABA neurons in the VTA which in turn then disinhibit DA neurons in the VTA.

6.2 Afferents

The LH receives many different inputs from different cortical and subcortical areas which are both excitatory and inhibitory. The PFC seems to be an important source of excitatory information, in particular the mPFC. It also receives inputs from the extended Amygdala and the hippocampus [Stuber and Wise, 2016].

6.3 Behavioural experiments

Feeding is stimulated when LH cells are activated by glutamate agonists [Stanley et al., 1993] and that stimulation of GABAergic cells in the LH inhibits feeding [Stanley et al., 2011].

7 Nucleus Accumbens Core

7.1 Afferents

The NAc core receives inputs from the dorsal-medial prefrontal cortex and the hippocampus [Brog et al., 1993a].

7.2 Efferents

There are two distinct output pathways from the NAcc core which have its origins from the two sub-populations of neurons in the NAcc core. The one sub-population carries mainly D1 receptors and the other one carries mainly D2 receptors [Kelley, 2004] [Humphries and Prescott, 2010].

7.2.1 Direct pathway

The D1 receptor carrying neurons feed directly into the SNr and are able to inhibit tonically active SNr neurons, thus the NAcc core is able to disinhibit motor programs.

7.2.2 Indirect pathway

There is an indirect pathway via the VP to the SNr originating from the NAcc core. In contrast to the direct pathway these neurons in the indirect pathway carry mainly D2 receptors which are inhibitory in nature. D2 receptors are very sensitive to low DA concentrations and will react to the tonic DA concentrations.

8 Nucleus Accumbens Shell

The shell can be further divided into the medial and lateral shell [Ikemoto, 2007] [Usuda et al., 1998] used in the model by [Humphries and Prescott, 2010].

8.1 Medial Shell

The medial Shell projects to the medial Ventral Pallidum (VP) [Ikemoto, 2007].

8.2 Lateral Shell

The lateral shall projects to the ventrolateral Ventral Pallium (VP) [Ikemoto, 2007].

8.3 Behavioural experiments

The shell seems to be responsible for behavioural flexibility meaning that it controls the animal's ability to shift to another target when the reward is lost [Aquili et al., 2014]

8.4 Function

Overall the Shell seems to learn the stimuli which are associated with a reward, and thus enhances the future salience of those stimuli [Cassidy and Tong, 2017].

In this context it is also interesting that shell DA also tracks rather the inventive value than the reward prediction error [Sackett et al., 2017] which probably means that the VTA has regions which work with the shell and that they are distinct from the core.

9 Nucleus Accumbens (NAcc)

The Nacc is the ventral extension of the striatum, and therefore also called the ventral striatum. The ventral striatum contains neurons known as medium spiny neurons (MSN's).

The accumbens has two major subterritories: the shell and the core [Heimer et al., 1991] where the shell can be further subdivided [Usuda et al., 1998] in lateral and medial parts and the core rather depending on its D1 or D2 receptors.

[[nacc-shell]]

[[nacc-core]]

9.1 Behavioural experiments

The Nacc core and shell have distinct roles controlling reward based learning:

In reversal learning [Dalton et al., 2014] the shell seems to control switching contingencies (i.e. the reward is moved from one site to another) whereas the core controls the actual approach behaviour towards the rewarding site.

Impulsivity is altered by 5HT antagonists which point towards an important role of 5HT in the retrieval of delayed rewards. 5-HT(2A) antagonists reduced decreased impulsive responding and the 5-HT(2C) antagonist increased impulsivity [Robinson et al., 2008].

9.2 Signals

By measuring the DA concentration in both the core and shell [Saddoris et al., 2015] it turns out that Core DA follows the classical prediction error signals where it spikes most to the predicting cue whereas the shell responds to all reward related events during the experiment.

9.3 Plasticity

It's well known the bursts of dopamine cause LTP in conjunction with pre- and (possible) postsynaptic activity (so called 3 factor Hebbian rule) or heterosynaptic LTP. However [Goto and Grace, 2005] showed that D1 receptors cause LTP on hippocampal fibres whereas D2 receptors control the cortical inputs in the opposite way. It appears to be that 5HT causes retrograde cannabinoid (CB) release which inhibits pre-synaptic GLU release [Burattini et al., 2014]. However, also postsynaptic HFS causes CB1 mediated depression. [Mathur et al., 2011] tested this more thoroughly in that they conclude that 5HT actually causes presynaptic inhibition via the 5HT1B

receptor, with that LTD and that HFS can have a similar effect by releasing CB.

10 Orbitofrontal Cortex (OFC)

The OFC associates sensory stimuli with reward related information [Schoenbaum et al., 2009] or in other words it computes the (potential) reward value of a sensor cue [?], [Bari and Robbins, 2013].

10.1 Afferents

The OFC receives inputs from a wide range of brain areas which allows it associate sensor cues (and also actions) to rewards. [?] provides an overview of these inputs which are from the:

- hippocampus
- subiculum
- PFC
- perihirnal cortex, and
- nucleus reuniens

Serotonin seems to have a strong effect on the OFC which has been shown by [Zhou et al., 2015]. Stimulation of the DRN results in both excitatory activity and inhibitory activity in the OFC. In addition the release of 5HT has a strong impact on plasticity: after pairing an odour stimulus with the release of 5HT the odour stimulus creates long lasting activity in the OFC which starts at the presentation of the stimulus and ends after reward delivery [Zhou et al., 2015].

10.2 Efferents

Its major subcortical targets include the dorsal raphe nucleus [Luo et al., 2015], medial striatum, NAcc, lateral preoptic area, amygdala and the hypothalamus [Vertes et al., 2012].

* The l-OFC innervates the l-shell and amygdala and * the mOFC the m-shell, hippocampus and amygdala [Brog et al., 1993b] [Noonan et al., 2012]

10.3 Neuronal activity

The paper by [Tremblay and Schultz, 1999] proposed that OFC neurons code the motivational value of rewards. The activity of OFC neurons increased in response to reward-predicting stimuli, during the expectation of rewards, and after the receipt of rewards. Also actions, associated with rewards, increase the firing rates of OFC neurons [?].

10.4 Behavioural experiments

The review by [?] presents behavioural experiments involving the OFC which confirm that the OFC computes behavioural reward value computed from sensory cues, such as odour, and actions. They contrast this to the hippocampus which computes reward value in relation to place fields.

Spatial reversal learning is improved by injecting the 5-HT(2C) receptor antagonist into the OFC [Boulougouris and Robbins, 2010].

Reversal learning is impaired if 5HT processing is disrupted in the OFC [Bari and Robbins, 2013].

10.4.1 Medial Prefrontal Cortex (mPFC)

The mPFC is considered by Homberg to be the most important location for top level cognitive functions.

The ventral portion of the mPFC is called Infralimbic cortex (IL) [Tsutsui-Kimura et al., 2016].

10.5 Efferents

A well known target of the mPFC is the Nacc core where the Nacc seems to be taking the role for response inhibition and waiting [Neufang et al., 2016] [Feja et al., 2014].

Projections from the mPFC to the DRN, allowing the mPFC to regulate 5HT firing and therefore control its own 5HT innervation [Homberg, 2012][Juckel et al., 1999]. This projection is inhibitory (see also the DRN page).

10.6 Neuromodulation

5HT plays a significant role in the mPFC which has been shown in great detail in [Santana and Artigas, 2017] and concluded that 5-HT1A, 5-HT2A, 5-HT2C, and 5-HT3, dopamine D1 and D2 are widely expressed in the mPFC.

The receptor 5-HT₃ was only expressed on GABAergic interneurons while all the other ones were expressed on both pyramidal and interneurons.

10.7 Behaviour

Lesions to the IL causes more impulsive behaviour [Tsutsui-Kimura et al., 2016] and reduction in 5HT is also associated with less impulse control [Neufang et al., 2016].

11 Rostral Medial Tegmental Nucleus (RMTg)

The newly discovered rostral medial tegmental nucleus, also called the tail of the VTA, is partially embedded in the VTA [Bourdy and Barrot, 2012]. It has been suggested that it has an ideal location to function as a switch between opposing aversion and reward responding areas and to direct information to DA neurons [Barrot et al., 2012].

11.1 Afferents

The main afferent to the RMTg is the glutamergic connection from the LHB, which is 7 times stronger than the LHB projection to the VTA [Barrot et al., 2012] and other afferents include the VTA and SNc [Lavezzi and Zahm, 2011].

RMTg GABA neurons differ in their targets to the VTA GABA neurons which, for example, target the forebrain in large numbers [Barrot et al., 2012].

11.2 Efferents

The RMTgs GABA efferents are the principal inhibitory connection to the VTA and SNc and play a critical role in RPE and aversive signalling [Bourdy and Barrot, 2012]. The RMTg also sends projections to other neuromodulatory systems including the raphe nucleus and the locus ceruleus [Barrot et al., 2012] [Hong et al., 2011].

12 Ventral Pallidum (VP)

The ventral pallidum is described as the limbic area of the pallidal complex as many reward circuits converge on this region. It encodes reward and motivation information engendered by rewarding stimuli [Smith et al., 2009]. The VP is divided into medial and lateral sections [Sesack and Grace, 2010].

12.1 Afferents

The VP is innervated by inhibitory GABA connections from the NAcc [Basar et al., 2010].

See Nacc core/shell for the exact projections.

12.2 Efferents

VP efferents project to the SNr, EP, prefrontal cortex, thalamus, LHb and the VTA [Groenewegen et al., 1993] [Ikemoto, 2007].

The ventral pallidum projects to the mediodorsal thalamic nucleus which in turn then projects to the infralimbic, prelimbic, agranular insular and cingulate cortex [Ikemoto, 2007].

The m-VP is the main source of GABAergic innervation to the VTA [Sesack and Grace, 2010].

13 Ventral Tegmental Area (VTA)

Dopamine neurons make up 60-65

13.1 Afferents

Excitatory afferents to the VTA include the LH, PFC and pedunclopontine nucleus (PPTg). Inhibitory, modulatory projections include the NAcc and the VP [Sesack and Grace, 2010].

Electrophysiology suggest that the NAcc projects mainly on the GABAergic neurons in the VTA.

13.2 Efferents

The VTA projects to numerous targets which include [Beckstead et al., 1979]:
* Nucleus Accumbens (NAcc) * Lateral habenula (LHb), nuclei reuniens and centralis medius, and the most medial zone of the mediodorsal nucleus * posterior hypothalamic nucleus * Amygdala (central, lateral and medial) * Bed nucleus of the stria terminalis, * Nucleus of the diagonal band, and the medial half of the lateral septal nucleus * Anteromedial (frontocingulate) cortex * Entorhinal cortex

Although it is principally cited for its DA output GABA also plays a major role in the activity of the VTA. GABA projections from the VTA to the NAcc are reciprocated with GABA projections to the VTA. There is also a large projection of GABA neurons from the VTA to the PFC

[Carr and Sesack, 2000]. Local GABA neurons can also inhibit their neighbouring dopamine neurons.[Sesack and Grace, 2010] and are a strong candidate to calculate the reward prediction error [Eshel et al., 2015].

13.3 Function

It is very well known that rats self stimulate the VTA indefinitely [Stuber and Wise, 2016] and that phasic optogenetic activation of the VTA drives behavioural conditioning [Tsai et al., 2009].

In particular the pathway from the VTA to the NAc core (NAcc) and NAc shell (NAcSh) is instrumental here. VTA dopamine release in response to a rewarding stimulus induces goal-directed behaviour to acquire and consume it [Morales and Margolis, 2017].

In an experiment where DA release was artificially triggered via an optogenetic stimulation caused robust reward seeking behaviour [Steinberg et al., 2013].

13.3.1 Phasic activity

Looking at single cell recordings some neurons were excited by the reward (US), some by the reward predicting CS and some reacted to both stimuli [Cohen et al., 2012]. The response to the CS became gradually stronger and the ones to the US smaller.

DA in the VTA signals a reward prediction error resembling that of TD learning which has been first suggested by [Schultz et al., 1997] and then matched quantitatively by [Bayer and Glimcher, 2005].

DA VTA neurons react strongly to unexpected rewards, these responses diminish after repeated presentation of the reward but then rather spike when a CS is presented which predicts the reward.

During omission of the reward the DA activity supposed to experience a 'dip' in activity [Takahashi et al., 2017]. However, except of a few examples it is usually a reduction of the DA response after omission.

Also the DA activity won't vanish completely after a reward is expected but is diminished. This behaviour can still be matched on TD learning when using long-lasting eligibility traces [Pan et al., 2005].

However, [Sadacca et al., 2016] has recently challenged this view that DA neurons code simply a reward prediction error about an experienced reward but that they also respond to putative cached values of cues which have been previously paired with a reward.

13.3.2 Tonic activity (DA)

On the other hand the VTA generates tonic activity which can be seen as a motivational value signal which is principally sent to the NAcc [Sesack and Grace, 2010][Bromberg-Martin et al., 2010] .

13.3.3 Tonic activity (GABA)

Cohen et als 2012 research found that VTA GABA neurons signalled expected reward [Cohen et al., 2012] so that this can be used to calculate the reward prediction error locally in the VTA.

A License

This work is licensed under the Creative Commons Attribution 4.0 International License. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/> or send a letter to Creative Commons, PO Box 1866, Mountain View, CA 94042, USA.



References

- G. K. Aghajanian, J. S. Sprouse, P. Sheldon, and K. Rasmussen. Electrophysiology of the central serotonin system: receptor subtypes and transducer mechanisms. *Annals of the New York Academy of Sciences*, 600:93–103; discussion 103, 1990. ISSN 0077-8923. URL <http://www.ncbi.nlm.nih.gov/pubmed/2123618>.
- G. F. Alheid. Extended amygdala and basal forebrain. *Annals of the New York Academy of Sciences*, 985:185–205, Apr 2003. ISSN 0077-8923. URL <http://www.ncbi.nlm.nih.gov/pubmed/12724159>.
- L. Aquili, A. W. Liu, M. Shindou, T. Shindou, and J. R. Wickens. Behavioral flexibility is increased by optogenetic inhibition of neurons in the nucleus accumbens shell during specific time segments. *Learning & memory (Cold Spring Harbor, N.Y.)*, 21(4):223–231, Apr 2014. ISSN 1549-5485. doi: 10.1101/lm.034199.113. URL <http://www.ncbi.nlm.nih.gov/pubmed/24639489>.

- M. Araki, P. L. McGeer, and E. G. McGeer. Retrograde hrp tracing combined with a pharmacohistochemical method for gaba transaminase for the identification of presumptive gabaergic projections to the habenula. *Brain research*, 304(2):271–277, Jun 1984. ISSN 0006-8993. URL <http://www.ncbi.nlm.nih.gov/pubmed/6331587>.
- A. Bari and T. W. Robbins. Inhibition and impulsivity: behavioral and neural basis of response control. *Progress in neurobiology*, 108:44–79, Sep 2013. ISSN 1873-5118. doi: 10.1016/j.pneurobio.2013.06.005. URL <http://www.ncbi.nlm.nih.gov/pubmed/23856628>.
- M. Barrot, S. R. Sesack, F. Georges, M. Pistis, S. Hong, and T. C. Jhou. Braking dopamine systems: a new gaba master structure for mesolimbic and nigrostriatal functions. *The Journal of neuroscience : the official journal of the Society for Neuroscience*, 32(41):14094–14101, Oct 2012. ISSN 1529-2401. doi: 10.1523/JNEUROSCI.3370-12.2012. URL <http://www.ncbi.nlm.nih.gov/pubmed/23055478>.
- K. Basar, T. Sesia, H. Groenewegen, H. W. M. Steinbusch, V. Visser-Vandewalle, and Y. Temel. Nucleus accumbens and impulsivity. *Progress in neurobiology*, 92(4):533–557, Dec 2010. ISSN 1873-5118. doi: 10.1016/j.pneurobio.2010.08.007. URL <http://www.ncbi.nlm.nih.gov/pubmed/20831892>.
- H. M. Bayer and P. W. Glimcher. Midbrain dopamine neurons encode a quantitative reward prediction error signal. *Neuron*, 47(1):129–141, Jul 2005. ISSN 0896-6273. doi: 10.1016/j.neuron.2005.05.020. URL <http://www.ncbi.nlm.nih.gov/pubmed/15996553>.
- R. M. Beckstead, V. B. Domesick, and W. J. Nauta. Efferent connections of the substantia nigra and ventral tegmental area in the rat. *Brain research*, 175(2):191–217, Oct 1979. ISSN 0006-8993. URL <http://www.ncbi.nlm.nih.gov/pubmed/314832>.
- K. C. Berridge. 'liking' and 'wanting' food rewards: brain substrates and roles in eating disorders. *Physiology & behavior*, 97(5):537–550, Jul 2009. ISSN 1873-507X. doi: 10.1016/j.physbeh.2009.02.044. URL <http://www.ncbi.nlm.nih.gov/pubmed/19336238>.
- H. Berthoud. Mind versus metabolism in the control of food intake and energy balance. *Physiol Behav*, 81(5):781–793, Jul 2004.

- V. Boulougouris and T. W. Robbins. Enhancement of spatial reversal learning by 5-HT_{2C} receptor antagonism is neuroanatomically specific. *The Journal of neuroscience : the official journal of the Society for Neuroscience*, 30(3):930–938, Jan 2010. ISSN 1529-2401. doi: 10.1523/JNEUROSCI.4312-09.2010. URL <http://www.ncbi.nlm.nih.gov/pubmed/20089901>.
- R. Bourdy and M. Barrot. A new control center for dopaminergic systems: pulling the vta by the tail. *Trends in neurosciences*, 35(11):681–690, Nov 2012. ISSN 1878-108X. doi: 10.1016/j.tins.2012.06.007. URL <http://www.ncbi.nlm.nih.gov/pubmed/22824232>.
- J. Brog, A. Salyapongse, A. Deutch, and D. Zahm. The patterns of afferent innervation of the core and shell in the "accumbens" part of the rat ventral striatum: immunohistochemical detection of retrogradely transported fluoro-gold. *J Comp Neurol*, 338(2):255–278, Dec 1993a.
- J. S. Brog, A. Salyapongse, A. Y. Deutch, and D. S. Zahm. The patterns of afferent innervation of the core and shell in the "accumbens" part of the rat ventral striatum: immunohistochemical detection of retrogradely transported fluoro-gold. *The Journal of comparative neurology*, 338(2):255–278, Dec 1993b. ISSN 0021-9967. doi: 10.1002/cne.903380209. URL <http://www.ncbi.nlm.nih.gov/pubmed/8308171>.
- E. S. Bromberg-Martin, M. Matsumoto, and O. Hikosaka. Dopamine in motivational control: rewarding, aversive, and alerting. *Neuron*, 68(5):815–834, Dec 2010. ISSN 1097-4199. doi: 10.1016/j.neuron.2010.11.022. URL <http://www.ncbi.nlm.nih.gov/pubmed/21144997>.
- C. Burattini, G. Battistini, F. Tamagnini, and G. Aicardi. Low-frequency stimulation evokes serotonin release in the nucleus accumbens and induces long-term depression via production of endocannabinoid. *Journal of neurophysiology*, 111(5):1046–1055, Mar 2014. ISSN 1522-1598. doi: 10.1152/jn.00498.2013. URL <http://www.ncbi.nlm.nih.gov/pubmed/24335217>.
- R. L. Carhart-Harris and D. J. Nutt. Serotonin and brain function: a tale of two receptors. *Journal of psychopharmacology (Oxford, England)*, 31(9):1091–1120, Sep 2017. ISSN 1461-7285. doi: 10.1177/0269881117725915. URL <http://www.ncbi.nlm.nih.gov/pubmed/28858536>.
- D. B. Carr and S. R. Sesack. Gaba-containing neurons in the rat ventral tegmental area project to the prefrontal cortex. *Synapse (New York, N.Y.)*, 38(2):114–123, Nov 2000. ISSN 0887-4476.

- doi: 10.1002/1098-2396(200011)38:2<114::AID-SYN2>3.0.CO;2-R. URL <http://www.ncbi.nlm.nih.gov/pubmed/11018785>.
- R. M. Cassidy and Q. Tong. Hunger and satiety gauge reward sensitivity. *Frontiers in endocrinology*, 8:104, May 2017. ISSN 1664-2392. doi: 10.3389/fendo.2017.00104. URL <http://www.ncbi.nlm.nih.gov/pubmed/28572791>.
- D. C. Castro, S. L. Cole, and K. C. Berridge. Lateral hypothalamus, nucleus accumbens, and ventral pallidum roles in eating and hunger: interactions between homeostatic and reward circuitry. *Frontiers in systems neuroscience*, 9:90, Jun 2015. ISSN 1662-5137. doi: 10.3389/fnsys.2015.00090. URL <http://www.ncbi.nlm.nih.gov/pubmed/26124708>.
- P. Celada, M. V. Puig, J. M. Casanovas, G. Guillazo, and F. Artigas. Control of dorsal raphe serotonergic neurons by the medial prefrontal cortex: Involvement of serotonin-1a, gaba(a), and glutamate receptors. *The Journal of neuroscience : the official journal of the Society for Neuroscience*, 21(24):9917–9929, Dec 2001. ISSN 1529-2401. URL <http://www.ncbi.nlm.nih.gov/pubmed/11739599>.
- G. R. Christoph, R. J. Leonzio, and K. S. Wilcox. Stimulation of the lateral habenula inhibits dopamine-containing neurons in the substantia nigra and ventral tegmental area of the rat. *The Journal of neuroscience : the official journal of the Society for Neuroscience*, 6(3):613–619, Mar 1986. ISSN 0270-6474. URL <http://www.ncbi.nlm.nih.gov/pubmed/3958786>.
- J. Y. Cohen, S. Haesler, L. Vong, B. B. Lowell, and N. Uchida. Neuron-type-specific signals for reward and punishment in the ventral tegmental area. *Nature*, 482(7383):85–88, Jan 2012. ISSN 1476-4687. doi: 10.1038/nature10754. URL <http://www.ncbi.nlm.nih.gov/pubmed/22258508>.
- G. L. Dalton, A. G. Phillips, and S. B. Floresco. Preferential involvement by nucleus accumbens shell in mediating probabilistic learning and reversal shifts. *The Journal of neuroscience : the official journal of the Society for Neuroscience*, 34(13):4618–4626, Mar 2014. ISSN 1529-2401. doi: 10.1523/JNEUROSCI.5058-13.2014. URL <http://www.ncbi.nlm.nih.gov/pubmed/24672007>.
- N. Eshel, M. Bukwich, V. Rao, V. Hemmelder, J. Tian, and N. Uchida. Arithmetic and local circuitry underlying dopamine prediction errors. *Nature*, 525(7568):243–246, Sep 2015. ISSN 1476-4687. doi: 10.1038/nature14855. URL <http://www.ncbi.nlm.nih.gov/pubmed/26322583>.

- M. Feja, L. Hayn, and M. Koch. Nucleus accumbens core and shell in-activation differentially affects impulsive behaviours in rats. *Progress in neuro-psychopharmacology & biological psychiatry*, 54:31–42, Oct 2014. ISSN 1878-4216. doi: 10.1016/j.pnpbp.2014.04.012. URL <http://www.ncbi.nlm.nih.gov/pubmed/24810333>.
- P. J. Fletcher, M. Tampakeras, J. Sinyard, and G. A. Higgins. Opposing effects of 5-HT(2a) and 5-HT(2c) receptor antagonists in the rat and mouse on premature responding in the five-choice serial reaction time test. *Psychopharmacology*, 195(2):223–234, Dec 2007. ISSN 0033-3158. doi: 10.1007/s00213-007-0891-z. URL <http://www.ncbi.nlm.nih.gov/pubmed/17673981>.
- A. Frazer and J. Hensler. Serotonin receptors. In G. Siegel, B. Agranoff, and R. Albers, editors, *Basic Neurochemistry: Molecular, Cellular and Medical Aspects*. Lippincott-Raven, Philadelphia, 6th edition, 1999.
- L. Gonçalves, M. I. Nogueira, S. J. Shammah-Lagnado, and M. Metzger. Prefrontal afferents to the dorsal raphe nucleus in the rat. *Brain research bulletin*, 78(4-5):240–247, Mar 2009. ISSN 1873-2747. doi: 10.1016/j.brainresbull.2008.11.012. URL <http://www.ncbi.nlm.nih.gov/pubmed/19103268>.
- Y. Goto and A. A. Grace. Dopaminergic modulation of limbic and cortical drive of nucleus accumbens in goal-directed behavior. *Nature neuroscience*, 8(6):805–812, Jun 2005. ISSN 1097-6256. doi: 10.1038/nn1471. URL <http://www.ncbi.nlm.nih.gov/pubmed/15908948>.
- H. J. Groenewegen, H. W. Berendse, and S. N. Haber. Organization of the output of the ventral striatopallidal system in the rat: ventral pallidal efferents. *Neuroscience*, 57(1):113–142, Nov 1993. ISSN 0306-4522. URL <http://www.ncbi.nlm.nih.gov/pubmed/8278047>.
- H. Hagena and D. Manahan-Vaughan. The serotonergic 5-HT₄ receptor: A unique modulator of hippocampal synaptic information processing and cognition. *Neurobiology of learning and memory*, 138:145–153, Feb 2017. ISSN 1095-9564. doi: 10.1016/j.nlm.2016.06.014. URL <http://www.ncbi.nlm.nih.gov/pubmed/27317942>.
- A. A. Harrison, B. J. Everitt, and T. W. Robbins. Central 5-HT depletion enhances impulsive responding without affecting the accuracy of attentional performance: interactions with dopaminergic mechanisms. *Psychopharmacology*, 133(4):329–342, Oct 1997. ISSN 0033-3158. URL <http://www.ncbi.nlm.nih.gov/pubmed/9372531>.

- L. Heimer, D. Zahm, L. Churchill, P. Kalivas, and C. Wohltmann. Specificity in the projection patterns of accumbal core and shell in the rat. *Neuroscience*, 41(1):89–125, 1991.
- J. R. Homberg. Serotonin and decision making processes. *Neuroscience and biobehavioral reviews*, 36(1):218–236, Jan 2012. ISSN 1873-7528. doi: 10.1016/j.neubiorev.2011.06.001. URL <http://www.ncbi.nlm.nih.gov/pubmed/21693132>.
- S. Hong and O. Hikosaka. The globus pallidus sends reward-related signals to the lateral habenula. *Neuron*, 60(4):720–729, Nov 2008. ISSN 1097-4199. doi: 10.1016/j.neuron.2008.09.035. URL <http://www.ncbi.nlm.nih.gov/pubmed/19038227>.
- S. Hong, T. C. Jhou, M. Smith, K. S. Saleem, and O. Hikosaka. Negative reward signals from the lateral habenula to dopamine neurons are mediated by rostromedial tegmental nucleus in primates. *The Journal of neuroscience : the official journal of the Society for Neuroscience*, 31(32):11457–11471, Aug 2011. ISSN 1529-2401. doi: 10.1523/JNEUROSCI.1384-11.2011. URL <http://www.ncbi.nlm.nih.gov/pubmed/21832176>.
- M. D. Humphries and T. J. Prescott. The ventral basal ganglia, a selection mechanism at the crossroads of space, strategy, and reward. *Progress in neurobiology*, 90(4):385–417, Apr 2010. ISSN 1873-5118. doi: 10.1016/j.pneurobio.2009.11.003. URL <http://www.ncbi.nlm.nih.gov/pubmed/19941931>.
- S. Ikemoto. Dopamine reward circuitry: two projection systems from the ventral midbrain to the nucleus accumbens-olfactory tubercle complex. *Brain research reviews*, 56(1):27–78, Nov 2007. ISSN 0165-0173. doi: 10.1016/j.brainresrev.2007.05.004. URL <http://www.ncbi.nlm.nih.gov/pubmed/17574681>.
- H. Ji and P. D. Shepard. Lateral habenula stimulation inhibits rat midbrain dopamine neurons through a gaba(a) receptor-mediated mechanism. *The Journal of neuroscience : the official journal of the Society for Neuroscience*, 27(26):6923–6930, Jun 2007. ISSN 1529-2401. doi: 10.1523/JNEUROSCI.0958-07.2007. URL <http://www.ncbi.nlm.nih.gov/pubmed/17596440>.
- G. Juckel, A. Mendlin, and B. L. Jacobs. Electrical stimulation of rat medial prefrontal cortex enhances forebrain serotonin output: implications for electroconvulsive therapy and transcranial magnetic stimulation in depression. *Neuropsychopharmacology : official publication of*

- the American College of Neuropsychopharmacology*, 21(3):391–398, Sep 1999. ISSN 0893-133X. doi: 10.1016/S0893-133X(98)00097-9. URL <http://www.ncbi.nlm.nih.gov/pubmed/10457536>.
- A. Kelley. Ventral striatal control of appetitive motivation: Role in ingestive behavior and reward-related learning. *Neurosci Biobehav Rev*, 27(8):765–776, Jan 2004.
- A. Khani and G. Rainer. Neural and neurochemical basis of reinforcement-guided decision making. *Journal of neurophysiology*, 116(2):724–741, Aug 2016. ISSN 1522-1598. doi: 10.1152/jn.01113.2015. URL <http://www.ncbi.nlm.nih.gov/pubmed/27226454>.
- A. B. Kowski, R. W. Veh, and T. Weiss. Dopaminergic activation excites rat lateral habenular neurons in vivo. *Neuroscience*, 161(4):1154–1165, Jul 2009. ISSN 1873-7544. doi: 10.1016/j.neuroscience.2009.04.026. URL <http://www.ncbi.nlm.nih.gov/pubmed/19374940>.
- H. N. Lavezzi and D. S. Zahm. The mesopontine rostromedial tegmental nucleus: an integrative modulator of the reward system. *Basal ganglia*, 1(4):191–200, Nov 2011. ISSN 2210-5336. doi: 10.1016/j.baga.2011.08.003. URL <http://www.ncbi.nlm.nih.gov/pubmed/22163100>.
- H. S. Lee, M. A. Kim, R. J. Valentino, and B. D. Waterhouse. Glutamatergic afferent projections to the dorsal raphe nucleus of the rat. *Brain research*, 963(1-2):57–71, Feb 2003. ISSN 0006-8993. URL <http://www.ncbi.nlm.nih.gov/pubmed/12560111>.
- Y. Li, W. Zhong, D. Wang, Q. Feng, Z. Liu, J. Zhou, C. Jia, F. Hu, J. Zeng, Q. Guo, L. Fu, and M. Luo. Serotonin neurons in the dorsal raphe nucleus encode reward signals. *Nature communications*, 7:10503, Jan 2016. ISSN 2041-1723. doi: 10.1038/ncomms10503. URL <http://www.ncbi.nlm.nih.gov/pubmed/26818705>.
- S. B. Linley, W. B. Hoover, and R. P. Vertes. Pattern of distribution of serotonergic fibers to the orbitomedial and insular cortex in the rat. *Journal of chemical neuroanatomy*, 48-49:29–45, Mar 2013. ISSN 1873-6300. doi: 10.1016/j.jchemneu.2012.12.006. URL <http://www.ncbi.nlm.nih.gov/pubmed/23337940>.
- E. Lottem, M. L. Lörincz, and Z. F. Mainen. Optogenetic activation of dorsal raphe serotonin neurons rapidly inhibits spontaneous but not odor-evoked activity in olfactory cortex. *The Journal of neuroscience*

- : *the official journal of the Society for Neuroscience*, 36(1):7–18, Jan 2016. ISSN 1529-2401. doi: 10.1523/JNEUROSCI.3008-15.2016. URL <http://www.ncbi.nlm.nih.gov/pubmed/26740645>.
- M. Luo, J. Zhou, and Z. Liu. Reward processing by the dorsal raphe nucleus: 5-HT and beyond. *Learning & memory (Cold Spring Harbor, N.Y.)*, 22(9): 452–460, Sep 2015. ISSN 1549-5485. doi: 10.1101/lm.037317.114. URL <http://www.ncbi.nlm.nih.gov/pubmed/26286655>.
- B. N. Mathur, N. A. Capik, V. A. Alvarez, and D. M. Lovinger. Serotonin induces long-term depression at corticostriatal synapses. *The Journal of neuroscience : the official journal of the Society for Neuroscience*, 31(20):7402–7411, May 2011. ISSN 1529-2401. doi: 10.1523/JNEUROSCI.6250-10.2011. URL <http://www.ncbi.nlm.nih.gov/pubmed/21593324>.
- M. Matsumoto and O. Hikosaka. Lateral habenula as a source of negative reward signals in dopamine neurons. *Nature*, 447(7148):1111–1115, Jun 2007. ISSN 1476-4687. doi: 10.1038/nature05860. URL <http://www.ncbi.nlm.nih.gov/pubmed/17522629>.
- G. Mengod, R. Corts, M. T. Vilar, and D. Hoyer. Chapter 1.6 - distribution of 5-HT receptors in the central nervous system. In C. P. Mller and B. L. Jacobs, editors, *Handbook of the Behavioral Neurobiology of Serotonin*, volume 21 of *Handbook of Behavioral Neuroscience*, pages 123 – 138. Elsevier, 2010. doi: [https://doi.org/10.1016/S1569-7339\(10\)70074-6](https://doi.org/10.1016/S1569-7339(10)70074-6). URL <http://www.sciencedirect.com/science/article/pii/S1569733910700746>.
- K. A. Michelsen, C. Schmitz, and H. W. M. Steinbusch. The dorsal raphe nucleus—from silver stainings to a role in depression. *Brain research reviews*, 55(2):329–342, Oct 2007. ISSN 0165-0173. doi: 10.1016/j.brainresrev.2007.01.002. URL <http://www.ncbi.nlm.nih.gov/pubmed/17316819>.
- B. Mlinar, S. Mascalchi, G. Mannaioni, R. Morini, and R. Corradetti. 5-HT₄ receptor activation induces long-lasting epsp-spike potentiation in CA1 pyramidal neurons. *The European journal of neuroscience*, 24(3):719–731, Aug 2006. ISSN 0953-816X. doi: 10.1111/j.1460-9568.2006.04949.x. URL <http://www.ncbi.nlm.nih.gov/pubmed/16930402>.
- A. C. Mok and G. J. Mogenson. Effects of electrical stimulation of the lateral hypothalamus, hippocampus, amygdala and olfactory bulb on unit activity of the lateral habenular nucleus in the rat. *Brain research*, 77(3):417–429, Sep 1974. ISSN 0006-8993. URL <http://www.ncbi.nlm.nih.gov/pubmed/4851746>.

- M. Morales and E. B. Margolis. Ventral tegmental area: cellular heterogeneity, connectivity and behaviour. *Nature reviews. Neuroscience*, 18(2):73–85, Feb 2017. ISSN 1471-0048. doi: 10.1038/nrn.2016.165. URL <http://www.ncbi.nlm.nih.gov/pubmed/28053327>.
- K. Nakamura. The role of the dorsal raphé nucleus in reward-seeking behavior. *Frontiers in integrative neuroscience*, 7:60, Aug 2013. ISSN 1662-5145. doi: 10.3389/fnint.2013.00060. URL <http://www.ncbi.nlm.nih.gov/pubmed/23986662>.
- S. Neufang, A. Akhrif, C. G. Herrmann, C. Drepper, G. A. Homola, J. Nowak, J. Waider, A. G. Schmitt, K.-P. Lesch, and M. Romanos. Serotonergic modulation of 'waiting impulsivity' is mediated by the impulsivity phenotype in humans. *Translational psychiatry*, 6(11):e940, Nov 2016. ISSN 2158-3188. doi: 10.1038/tp.2016.210. URL <http://www.ncbi.nlm.nih.gov/pubmed/27824354>.
- M. P. Noonan, N. Kolling, M. E. Walton, and M. F. S. Rushworth. Re-evaluating the role of the orbitofrontal cortex in reward and reinforcement. *The European journal of neuroscience*, 35(7):997–1010, Apr 2012. ISSN 1460-9568. doi: 10.1111/j.1460-9568.2012.08023.x. URL <http://www.ncbi.nlm.nih.gov/pubmed/22487031>.
- S. K. Ogawa, J. Y. Cohen, D. Hwang, N. Uchida, and M. Watabe-Uchida. Organization of monosynaptic inputs to the serotonin and dopamine neuromodulatory systems. *Cell reports*, 8(4):1105–1118, Aug 2014. ISSN 2211-1247. doi: 10.1016/j.celrep.2014.06.042. URL <http://www.ncbi.nlm.nih.gov/pubmed/25108805>.
- O. P. Ottersen. Connections of the amygdala of the rat. iv: Corticoamygdaloid and intraamygdaloid connections as studied with axonal transport of horseradish peroxidase. *The Journal of comparative neurology*, 205(1):30–48, Feb 1982. ISSN 0021-9967. doi: 10.1002/cne.902050104. URL <http://www.ncbi.nlm.nih.gov/pubmed/7068948>.
- J. M. Palacios, C. Waeber, D. Hoyer, and G. Mengod. Distribution of serotonin receptors. *Annals of the New York Academy of Sciences*, 600:36–52, 1990. ISSN 0077-8923. URL <http://www.ncbi.nlm.nih.gov/pubmed/2252320>.
- W.-X. Pan, R. Schmidt, J. R. Wickens, and B. I. Hyland. Dopamine cells respond to predicted events during classical conditioning: evidence for eligibility traces in the reward-learning network. *The Journal of neuroscience* :

- the official journal of the Society for Neuroscience*, 25(26):6235–6242, Jun 2005. ISSN 1529-2401. doi: 10.1523/JNEUROSCI.1478-05.2005. URL <http://www.ncbi.nlm.nih.gov/pubmed/15987953>.
- A. Parent, S. Gravel, and R. Boucher. The origin of forebrain afferents to the habenula in rat, cat and monkey. *Brain research bulletin*, 6(1):23–38, Jan 1981. ISSN 0361-9230. URL <http://www.ncbi.nlm.nih.gov/pubmed/7470948>.
- R. Peñas-Cazorla and M. T. Vilaró. Serotonin 5-HT₄ receptors and forebrain cholinergic system: receptor expression in identified cell populations. *Brain structure & function*, 220(6):3413–3434, Nov 2015. ISSN 1863-2661. doi: 10.1007/s00429-014-0864-z. URL <http://www.ncbi.nlm.nih.gov/pubmed/25183542>.
- A. Pitkänen, M. Pikkarainen, N. Nurminen, and A. Ylinen. Reciprocal connections between the amygdala and the hippocampal formation, perirhinal cortex, and postrhinal cortex in rat. a review. *Annals of the New York Academy of Sciences*, 911:369–391, Jun 2000. ISSN 0077-8923. URL <http://www.ncbi.nlm.nih.gov/pubmed/10911886>.
- I. Pollak Dorocic, D. Fürth, Y. Xuan, Y. Johansson, L. Pozzi, G. Silberberg, M. Carlén, and K. Meletis. A whole-brain atlas of inputs to serotonergic neurons of the dorsal and median raphe nuclei. *Neuron*, 83(3):663–678, Aug 2014. ISSN 1097-4199. doi: 10.1016/j.neuron.2014.07.002. URL <http://www.ncbi.nlm.nih.gov/pubmed/25102561>.
- W. C. Poller, V. I. Madai, R. Bernard, G. Laube, and R. W. Veh. A glutamatergic projection from the lateral hypothalamus targets vta-projecting neurons in the lateral habenula of the rat. *Brain research*, 1507:45–60, Apr 2013. ISSN 1872-6240. doi: 10.1016/j.brainres.2013.01.029. URL <http://www.ncbi.nlm.nih.gov/pubmed/23348378>.
- L. A. Quina, L. Tempest, L. Ng, J. A. Harris, S. Ferguson, T. C. Jhou, and E. E. Turner. Efferent pathways of the mouse lateral habenula. *The Journal of comparative neurology*, 523(1):32–60, Jan 2015. ISSN 1096-9861. doi: 10.1002/cne.23662. URL <http://www.ncbi.nlm.nih.gov/pubmed/25099741>.
- N. Rajakumar, K. Elisevich, and B. A. Flumerfelt. Compartmental origin of the striato-entopeduncular projection in the rat. *The Journal of comparative neurology*, 331(2):286–296, May 1993. ISSN 0021-9967. doi: 10.1002/cne.903310210. URL <http://www.ncbi.nlm.nih.gov/pubmed/8509503>.

- T. D. Reisine, P. Soubri  , A. Ferron, C. Blas, R. Romo, and J. Glowinski. Evidence for a dopaminergic innervation of the cat lateral habenula: its role in controlling serotonin transmission in the basal ganglia. *Brain research*, 308(2):281–288, Aug 1984. ISSN 0006-8993. URL <http://www.ncbi.nlm.nih.gov/pubmed/6478209>.
- M. Riad, S. Garcia, K. C. Watkins, N. Jodoin, E. Doucet, X. Langlois, S. el Mestikawy, M. Hamon, and L. Descarries. Somatodendritic localization of 5-HT_{1A} and preterminal axonal localization of 5-HT_{1B} serotonin receptors in adult rat brain. *The Journal of comparative neurology*, 417(2):181–194, Feb 2000. ISSN 0021-9967. URL <http://www.ncbi.nlm.nih.gov/pubmed/10660896>.
- A. C. Roberts. The importance of serotonin for orbitofrontal function. *Biological psychiatry*, 69(12):1185–1191, Jun 2011. ISSN 1873-2402. doi: 10.1016/j.biopsych.2010.12.037. URL <http://www.ncbi.nlm.nih.gov/pubmed/21353665>.
- E. S. J. Robinson, J. W. Dalley, D. E. H. Theobald, J. C. Glennon, M. A. Pezze, E. R. Murphy, and T. W. Robbins. Opposing roles for 5-HT_{2A} and 5-HT_{2C} receptors in the nucleus accumbens on inhibitory response control in the 5-choice serial reaction time task. *Neuropsychopharmacology : official publication of the American College of Neuropsychopharmacology*, 33(10):2398–2406, Sep 2008. ISSN 1740-634X. doi: 10.1038/sj.npp.1301636. URL <http://www.ncbi.nlm.nih.gov/pubmed/18046307>.
- J. A. Rosenkranz and A. A. Grace. Cellular mechanisms of infralimbic and prelimbic prefrontal cortical inhibition and dopaminergic modulation of basolateral amygdala neurons in vivo. *The Journal of neuroscience : the official journal of the Society for Neuroscience*, 22(1):324–337, Jan 2002. ISSN 1529-2401. URL <http://www.ncbi.nlm.nih.gov/pubmed/11756516>.
- D. A. Sackett, M. P. Saddoris, and R. M. Carelli. Nucleus accumbens shell dopamine preferentially tracks information related to outcome value of reward. *eNeuro*, 4(3), May/Jun 2017. ISSN 2373-2822. doi: 10.1523/ENEURO.0058-17.2017. URL <http://www.ncbi.nlm.nih.gov/pubmed/28593190>.
- B. F. Sadacca, J. L. Jones, and G. Schoenbaum. Midbrain dopamine neurons compute inferred and cached value prediction errors in a common framework. *eLife*, 5, Mar 2016. ISSN 2050-084X. doi: 10.7554/eLife.13665. URL <http://www.ncbi.nlm.nih.gov/pubmed/26949249>.

- M. P. Sadoris, F. Cacciapaglia, R. M. Wightman, and R. M. Carelli. Differential dopamine release dynamics in the nucleus accumbens core and shell reveal complementary signals for error prediction and incentive motivation. *The Journal of neuroscience : the official journal of the Society for Neuroscience*, 35(33):11572–11582, Aug 2015. ISSN 1529-2401. doi: 10.1523/JNEUROSCI.2344-15.2015. URL <http://www.ncbi.nlm.nih.gov/pubmed/26290234>.
- P. Sah, E. S. L. Faber, M. Lopez De Armentia, and J. Power. The amygdaloid complex: anatomy and physiology. *Physiological reviews*, 83(3): 803–834, Jul 2003. ISSN 0031-9333. doi: 10.1152/physrev.00002.2003. URL <http://www.ncbi.nlm.nih.gov/pubmed/12843409>.
- N. Santana and F. Artigas. Laminar and cellular distribution of monoamine receptors in rat medial prefrontal cortex. *Frontiers in neuroanatomy*, 11: 87, Sep 2017. ISSN 1662-5129. doi: 10.3389/fnana.2017.00087. URL <http://www.ncbi.nlm.nih.gov/pubmed/29033796>.
- G. Schoenbaum, M. R. Roesch, T. A. Stalnaker, and Y. K. Takahashi. A new perspective on the role of the orbitofrontal cortex in adaptive behaviour. *Nat. Rev. Neurosci.*, 10(12):885–92, 2009. doi: 10.1038/nrn2753.
- W. Schultz, P. Dayan, and P. R. Montague. A neural substrate of prediction and reward. *Science (New York, N.Y.)*, 275(5306):1593–1599, Mar 1997. ISSN 0036-8075. URL <http://www.ncbi.nlm.nih.gov/pubmed/9054347>.
- C. Sego, L. Gonçalves, L. Lima, I. C. Furigo, J. Donato, and M. Metzger. Lateral habenula and the rostromedial tegmental nucleus innervate neurochemically distinct subdivisions of the dorsal raphe nucleus in the rat. *The Journal of comparative neurology*, 522(7):1454–1484, May 2014. ISSN 1096-9861. doi: 10.1002/cne.23533. URL <http://www.ncbi.nlm.nih.gov/pubmed/24374795>.
- S. R. Sesack and A. A. Grace. Cortico-basal ganglia reward network: microcircuitry. *Neuropsychopharmacology : official publication of the American College of Neuropsychopharmacology*, 35(1):27–47, Jan 2010. ISSN 1740-634X. doi: 10.1038/npp.2009.93. URL <http://www.ncbi.nlm.nih.gov/pubmed/19675534>.
- S. J. Shabel, C. D. Proulx, A. Trias, R. T. Murphy, and R. Malinow. Input to the lateral habenula from the basal ganglia is excitatory, aversive, and suppressed by serotonin. *Neuron*, 74(3):475–481,

- May 2012. ISSN 1097-4199. doi: 10.1016/j.neuron.2012.02.037. URL <http://www.ncbi.nlm.nih.gov/pubmed/22578499>.
- X. Shen, X. Ruan, and H. Zhao. Stimulation of midbrain dopaminergic structures modifies firing rates of rat lateral habenula neurons. *PloS one*, 7(4):e34323, Apr 2012. ISSN 1932-6203. doi: 10.1371/journal.pone.0034323. URL <http://www.ncbi.nlm.nih.gov/pubmed/22485164>.
- K. S. Smith, A. J. Tindell, J. W. Aldridge, and K. C. Berridge. Ventral pallidum roles in reward and motivation. *Behavioural brain research*, 196(2):155–167, Jan 2009. ISSN 1872-7549. doi: 10.1016/j.bbr.2008.09.038. URL <http://www.ncbi.nlm.nih.gov/pubmed/18955088>.
- D. R. Sparta and G. D. Stuber. Cartography of serotonergic circuits. *Neuron*, 83(3):513–515, Aug 2014. ISSN 1097-4199. doi: 10.1016/j.neuron.2014.07.030. URL <http://www.ncbi.nlm.nih.gov/pubmed/25102556>.
- B. Stanley, V. Willett, H. Donias, L. Ha, and L. Spears. The lateral hypothalamus: a primary site mediating excitatory amino acid-elicited eating. *Brain Res*, 630(1-2):41–49, Dec 1993.
- B. G. Stanley, K. R. Urstadt, J. R. Charles, and T. Kee. Glutamate and gaba in lateral hypothalamic mechanisms controlling food intake. *Physiology & behavior*, 104(1):40–46, Jul 2011. ISSN 1873-507X. doi: 10.1016/j.physbeh.2011.04.046. URL <http://www.ncbi.nlm.nih.gov/pubmed/21550353>.
- E. E. Steinberg, R. Keiflin, J. R. Boivin, I. B. Witten, K. Deisseroth, and P. H. Janak. A causal link between prediction errors, dopamine neurons and learning. *Nature neuroscience*, 16(7):966–973, Jul 2013. ISSN 1546-1726. doi: 10.1038/nn.3413. URL <http://www.ncbi.nlm.nih.gov/pubmed/23708143>.
- G. D. Stuber and R. A. Wise. Lateral hypothalamic circuits for feeding and reward. *Nature neuroscience*, 19(2):198–205, Feb 2016. ISSN 1546-1726. doi: 10.1038/nn.4220. URL <http://www.ncbi.nlm.nih.gov/pubmed/26814589>.
- L. W. Swanson. The amygdala and its place in the cerebral hemisphere. *Annals of the New York Academy of Sciences*, 985:174–184, Apr 2003. ISSN 0077-8923. URL <http://www.ncbi.nlm.nih.gov/pubmed/12724158>.
- L. W. Swanson and G. D. Petrovich, Petrovicha. What is the amygdala? *Trends in Neurosci*, 21(8):323–331, 1998. URL <http://www.sciencedirect.com/science/article/pii/S016622369801265X>.

- Y. K. Takahashi, H. M. Batchelor, B. Liu, A. Khanna, M. Morales, and G. Schoenbaum. Dopamine neurons respond to errors in the prediction of sensory features of expected rewards. *Neuron*, 95(6):1395–1405.e3, Sep 2017. ISSN 1097-4199. doi: 10.1016/j.neuron.2017.08.025. URL <http://www.ncbi.nlm.nih.gov/pubmed/28910622>.
- L. Tremblay and W. Schultz. Relative reward preference in primate orbitofrontal cortex. *Nature*, 398(6729):704–708, Apr 1999. ISSN 0028-0836. doi: 10.1038/19525. URL <http://www.ncbi.nlm.nih.gov/pubmed/10227292>.
- H.-C. Tsai, F. Zhang, A. Adamantidis, G. D. Stuber, A. Bonci, L. de Lecea, and K. Deisseroth. Phasic firing in dopaminergic neurons is sufficient for behavioral conditioning. *Science (New York, N.Y.)*, 324(5930):1080–1084, May 2009. ISSN 1095-9203. doi: 10.1126/science.1168878. URL <http://www.ncbi.nlm.nih.gov/pubmed/19389999>.
- I. Tsutsui-Kimura, Y. Ohmura, T. Izumi, T. Matsushima, H. Amita, T. Yamaguchi, T. Yoshida, and M. Yoshioka. Neuronal codes for the inhibitory control of impulsive actions in the rat infralimbic cortex. *Behavioural brain research*, 296:361–372, Jan 2016. ISSN 1872-7549. doi: 10.1016/j.bbr.2015.08.025. URL <http://www.ncbi.nlm.nih.gov/pubmed/26341319>.
- I. Usuda, K. Tanaka, and T. Chiba. Efferent projections of the nucleus accumbens in the rat with special reference to subdivision of the nucleus: biotinylated dextran amine study. *Brain research*, 797(1):73–93, Jun 1998. ISSN 0006-8993. URL <http://www.ncbi.nlm.nih.gov/pubmed/9630528>.
- R. P. Vertes, S. B. Linley, and W. B. Hoover. Pattern of distribution of serotonergic fibers to the thalamus of the rat. *Brain structure & function*, 215(1):1–28, Jul 2010. ISSN 1863-2661. doi: 10.1007/s00429-010-0249-x. URL <http://www.ncbi.nlm.nih.gov/pubmed/20390296>.
- R. P. Vertes, W. B. Hoover, and J. J. Rodriguez. Projections of the central medial nucleus of the thalamus in the rat: node in cortical, striatal and limbic forebrain circuitry. *Neuroscience*, 219:120–136, Sep 2012. ISSN 1873-7544. doi: 10.1016/j.neuroscience.2012.04.067. URL <http://www.ncbi.nlm.nih.gov/pubmed/22575585>.
- M. L. Wallace, A. Saunders, K. W. Huang, A. C. Philson, M. Goldman, E. Z. Macosko, S. A. McCarroll, and B. L. Sabatini. Genetically distinct parallel pathways in the entopeduncular nucleus for limbic and sensorimotor output of the basal ganglia. *Neuron*, 94(1):138–152.e5,

Apr 2017. ISSN 1097-4199. doi: 10.1016/j.neuron.2017.03.017. URL <http://www.ncbi.nlm.nih.gov/pubmed/28384468>.

D. Zahm. An integrative neuroanatomical perspective on some subcortical substrates of adaptive responding with emphasis on the nucleus accumbens. *Neurosci Biobehav Rev*, 24(1):85–105, Jan 2000.

J. Zhou, C. Jia, Q. Feng, J. Bao, and M. Luo. Prospective coding of dorsal raphe reward signals by the orbitofrontal cortex. *The Journal of neuroscience : the official journal of the Society for Neuroscience*, 35(6):2717–2730, Feb 2015. ISSN 1529-2401. doi: 10.1523/JNEUROSCI.4017-14.2015. URL <http://www.ncbi.nlm.nih.gov/pubmed/25673861>.