

tRNA database 2026, extended compilation of tRNA sequences and tRNA genes

Bruno J. Schmidt,^{1,*} Frank Jühling,² Thomas Spicher,³ Ella Cassidy,^{4,9} Maximilian Feussner,⁵ Chase L. Beisel,⁵ Joern Pütz,^{2,6} Mario Mörl,⁷ Peter F. Stadler^{1,3,4,8,9,10} and Stephan H. Bernhart^{8,10}

¹Max Planck Institute for Mathematics in the Sciences, Inselstr 26, 04103, Leipzig, Germany

²Inserm, Institute for Translational Medicine and Liver Disease, University of Strasbourg, Street, Postcode, State, France

³Theoretical biochemistry group, Department for theoretical Chemistry, University of Vienna, Währinger Str. 17, 1090, Vienna, Austria

⁴Bioinformatics Group, Department for Computer Science, Leipzig University, Härtelstr. 16-18, 04107, Saxony, Germany

⁵Botnar Institute of Immune Engineering - BIIE, Klingelbergstr. 48, 4056, Basel, Switzerland

⁶Department, Organization, Street, Postcode, State, Country

⁷Institute of Biochemistry, Leipzig University, Johannisallee 23, 04103, Saxony, Germany

⁸Interdisciplinary Center for Bioinformatics - IZBI, Leipzig University, Härtelstr. 16-18, 04107, Saxony, Germany

⁹Center for Scalable Data Analytics and Artificial Intelligence, Leipzig University, Härtelstr. 16-18, 04107, Saxony, Germany

¹⁰LeiCeM - Leipzig Center of Metabolism, Leipzig University, Ritterstr. 26, 04109, Saxony, Germany

*Corresponding author. email-id.com

Abstract

The tRNAdb database published more than 15 years ago originated as an extension of Mathias Sprinzl's seminal collection of tRNA data dating back to the late 1970s. Throughout its lifespan, the tRNAdb has been used extensively, accumulating more than 1 000 citations. Here, we present tRNAdb 2026, an updated version hosted at Leipzig University. The tRNAdb 2026 features an approximately 10-fold increase in data coverage, containing more than 500 000 new mitochondrial and nuclear tRNA sequences together with the unaltered legacy dataset. Importantly, identifiers and metadata of the legacy dataset remain consistent with the original database, ensuring that previously published references remain valid and findable. The expanded dataset integrates publicly available tRNA annotations and sequences generated using contemporary prediction pipelines followed by semiautomated curation to ensure annotation quality and consistency. In addition, cross-references to external resources such as GtRNAdb facilitate interoperability with existing tRNA annotation frameworks. A modern web-based frontend provides intuitive and efficient access to tRNAdb 2026, while an API enables automated data retrieval and integration into bioinformatics workflows. To support transparent scientific documentation and reproducibility across future updates, we added a versioning system for both current and future database releases.

Keywords tRNA sequence, tRNA structure, database, tRNA modifications

Abbreviations Ala: Alanine, Arg: Arginine, Asn: Asparagine, Asp: Aspartic acid, Cys: Cysteine, Glu: Glutamic acid, Gln: Glutamine, Gly: Glycine, His: Histidine, Ile: Isoleucine, Leu: Leucine, Lys: Lysine, Met: Methionine, Phe: Phenylalanine, Pro: Proline, Ser: Serine, SeC: Selenocysteine, Thr: Threonine, Trp: Tryptophan, Tyr: Tyrosine, Val: Valine, Sup: Suppressor, AC: Anticodon, NCBI: National Center for Biotechnology Information, API: Application Programming Interface, PDB: Protein Data Bank, BV-BRC: Bacterial and Viral Bioinformatics Resource Center

Key Messages

- The tRNAdb has been greatly expanded
- We provide a modern web based front end for tRNAdb
- We provide an API and release versioning

Introduction

There has been growing interest in tRNAs, promoted by an increasingly broad portfolio of roles beyond their primary function of mediating the genetic code. Some 10 000 publications now link tRNAs with various aspects of cancer, with similar numbers concerning the

aspect of evolution of tRNAs and their processing. A solid, well-curated body of data is therefore a key resource for an increasing research community. Since the publication of the first "Compilation of tRNA sequences and sequences of tRNA genes" in 2005 (16), and its extension as the tRNAdb 2009 (8), a number of databases either

dedicated to, or containing tRNA sequences, have been published (3; 1; 15).

When access to the tRNAdb was interrupted due to technical issues in 2024/2025, the data was still widely used. A large number of requests for its continued existence prompted us to re-implemented the tRNAdb on a new, solid technical foundation, and to include an Application Programming Interface (API) for automatic data retrieval. For the present release, we extended its content by leveraging existing nuclear tRNAs from the gtrNA database as well as mitochondrial, archaeal and phage derived tRNAs we identified using MITOS2 (7), tRNAscan-SE 2.0.12 (4), and additional specialized approaches.

Implementation

Infrastructure and Implementation

tRNAdb2026 was completely reimplemented as a three tier application consisting of a MariaDB relational database, a flask API backend, and a CSS, HTML, and javascript based web frontend. This architecture enables independent change of individual components in order to ensure simple maintenance and future extensions. In contrast to the 2009 implementation (exclusively in Java and HTML/CSS), the new release aligns tRNAdb with modern software design philosophies. Notably, the tRNAdb website utilizes the public API endpoint exclusively for searching and retrieving data. To provide reliable download of large datasets, tRNAdb supports asynchronous data compilation and transfer via dedicated export threads as well as progress reporting of ongoing export jobs via polling.

Data model and provenance

For easy retrieval and global searches, sequences and their most relevant data are stored within a single, fully indexed relation enabling efficient filtering across all sequence categories, attributes, and selected metadata. This includes nucleotide sequence, secondary structure, anticodon, amino acid and origin (i.e., genomic DNA, chemically modified RNA, and mitochondrial DNA). Every sequence has a taxonomic identifier and a separate table holds relevant taxonomic information obtained from the National Center for Biotechnology Information (NCBI) taxonomy database (14). To facilitate legacy compatibility, the original tRNA dataset (8) remains unaltered regarding sequence, structure, anticodon, amino acid, references, and tRNAdb identifier. We solely updated the taxonomic assignment, i.e., species names and NCBI taxonomy identifiers, to be consistent with the current NCBI taxonomy database. We used the NCBI taxonomy data dump 2026-06-01, accessed 10 June 2026, to create an updated local taxonomic table and reconciled legacy taxonomic assignments using Bio.entrez (6) to resolve their current taxonomic information.

The updated relational schema supports a release versioning system which separates the legacy dataset and data contained in tRNAdb 2026 update. **Every sequence present in tRNAdb has a version identifier attached which links it to the first release it appeared in tRNAdb together with relevant release information.** For every release, data sources (i.e., origin database and version) and software (with corresponding version) used are documented. Additionally, every tRNAdb release is described with a public version string that is **exposed** on the “Details” page of individual entries, and

is also available as an export field. The public version string contains a short description how the sequences published in this particular release were obtained. A more detailed description, e.g., software and data sources, is available on the “Documentation and Help” page. Sequences in tRNAdb can be filtered by their corresponding release version, allowing users to browse exclusively legacy data, or current releases.

Subsequent releases of the tRNAdb dataset will be published with a corresponding release identifier together with the prior described release metadata. To enable a sustainable update procedure, we implemented a graphical user interface supported pipeline for import, normalization, and duplicate detection for subsequent tRNAdb versions.

API and programmatic access

With this update we provide a new public HTTP interface, accessible by POST JSON requests at <https://tdb.bioinf.uni-leipzig.de/api>. The API implements OpenAPI specifications and a thorough documentation for users is available on the “Documentation” page of the tRNAdb website. Fundamentally, our API supports nearly identical search parameters as the tRNAdb website and allows filtering by any combination of origin, species, clade, NCBI taxonomy ID, amino acid, anticodon, sequence and structural motifs, reference (genome accession, external source, publication), and release version. For many fields, multiple values can be supplied and entries matching at least one are returned. Individual records can be precisely retrieved by using the stable identifier **trnadb_id** which also remained unaltered for the 2009 legacy dataset.

No prior registration, and no personal information (e.g., email address, API key) is required to access the tRNAdb search API at <https://tdb.bioinf.uni-leipzig.de/api> but user requests are limited to one request per second and at most 1000 entries are returned per request to ensure fair use. Every API response, given the request was valid, contains a **total_results** field that states the total number of sequences matching a request and all relevant results can be **retrieved** paginated via an offset parameter. For requests that match a large number of records, and hence constitute bulk retrieval of tRNA sequences, a second API endpoint provides export to tsv, csv, json, or fasta which is available at <https://tdb.bioinf.uni-leipzig.de/export>. Note that the unique tRNAdb identifier **every entry is connected to** is stable, hence over changing releases, existing identifiers will not change and remain relevant for retrieval purposes. For usage details see the documentation page of the tRNAdb website.

Website

The tRNAdb web interface (see Figure 1) can be found at <https://tdb.bioinf.uni-leipzig.de/>. It provides user-friendly access to the tRNA data stored in our database and can be functionally separated into three subpages: “Search”, “Details”, and “Taxonomy Tree”. The main functionality of the website is the “Search” page which allows users to filter tRNA by organism, origin (e.g., nuclear DNA, chemically modified RNA, mitochondrial DNA) amino acid, anticodon, membership of taxonomic clades, external references (publications, genome IDs, external datasources) as well as sequence motifs and structural constraints. Species can be filtered precisely via NCBI taxonomy ID as well as their scientific species name and species synonyms as annotated in the NCBI taxonomy database. For every

[illegible]

To expand the set of mitochondrial tRNAs included in the tRNAdb, a total of 16 630 mitochondrial genome sequences, as well as their

annotations were obtained from NCBI Nucleotide (13), accessed **October 2024**. We used MITOS2 (7) without detection of proteins and rRNA and the refseq89m models on each mitochondrial genome to search for tRNAs. The predicted sequences were then compared to the NCBI annotation. All tRNAs that matched exactly to the NCBI annotation were added to tRNAdb. In case of discrepancies of location, we checked whether the structure of the acceptor stem predicted by Infernal (11) passed human curation. In the affirmative case, the sequence predicted by MITOS2 was deemed correct and included into the database. In case of discrepancies in the assignment of the strand (reading direction), we retained the MITOS2 prediction if no plausible anticodon could be identified in the anticodon stem and loop of the NCBI annotation. After filtering (see “Alignment and quality control section” below), this resulted in 305 905 novel predicted mtRNAs that were added to the tRNAdb.

Nuclear tRNAs

Genomic tRNAs were sourced from the GtRNAdb, using the data release from 2024 (3). Information retrieved included sequences, genomic locations and genome IDs. After filtering, 216 856 novel genomic tRNAs were added to the tRNAdb. GtRNAdb contains only a limited number of archaeal tRNA entries (below 9 000), covering a narrow taxonomic range within archaea. Since GtRNAdb also contains no virus-derived tRNAs, we decided to further expand tRNAdb in these two areas.

Archaeal tRNAs

Archaeal genomes were sourced from the Bacterial and Viral Bioinformatics Resource Center (BV-BRC) database (12), accessed **October 2024**. Only good quality genomes were retained for further analysis, and genome quality was assessed by CheckM2 ($\geq 85\%$ complete, $< 5\%$ contaminated) (5). These archaeal genomes cover a broad taxonomic range and include all major archaeal superphyla (Asgard, DPANN, TACK, and Euryarchaeota). We did not rely on the BV-BRC annotation of tRNA genes, as we found it inconsistent with predictions using the most up-to-date version of tRNAscan-SE 2.0.12 (4). Notably, tRNA-Ile2 is often mislabelled as tRNA-Met in BV-BRC, the most frequent mismatch we observed in isotype assignment (34%). This likely reflects the agmatidine wobble-base modification unique to archaea (4), highlighting the necessity to use domain specific models. Therefore, all genomes were annotated using tRNAscan-SE in archaeal mode. Selenocysteine tRNAs were identified using the SeC-specific covariance model, since it has an atypical secondary structure (4). As independent validation, all tRNAscan predictions were re-scored with Infernal (11) against tRNAinf-arch and tRNAinf-arch-SeC models. After filtering (see “Alignment and quality control section” below), 21 127 tRNA genes from 228 unique archaeal genomes were added, of which 214 were not represented in the 2009 tRNAdb, substantially expanding archaeal coverage relative to the legacy dataset.

Phage tRNAs

Genome sequences of all viruses annotated with the host term “Bacteria” were obtained from NCBI Virus (10), accessed 17 August 2026. Putative tRNA genes were identified using tRNAscan-SE v2.0.12 (4) in bacterial mode. Predicted tRNAs were retained only if they were 70–90nt in length, had an overall tRNAscan-SE inf

score > 60 , and had both HMM and secondary-structure scores > 10 . Predictions with undefined isotypes or classified as pseudogenes were excluded. After filtering (see “Alignment and quality control section” below), 7 099 viral tRNA genes from 1 558 unique genomes were added to tRNAdb.

Chemically modified tRNAs

Modified tRNA sequences, as well as their genomic variants, were obtained from the MODOMIC (15) database and the Protein Data Bank (PDB) (2). Sequences retrieved from the PDB having only a single modification located at the first or last base, were removed. In order display alignments properly, legacy one-letter codes for modifications were retained. The dictionary of one letter codes for modifications was extended to include any that were not present in the legacy tRNAdb 2009. For these novel modifications, we also identified the name of the compound using the PDB entry. The InChIKeys and InChis were extracted from PubChem (9) to check whether the compound was already present in the MODOMIC database. A list of all modifications with the one letter codes used in tRNAdb and MODOMIC can be found in the supplement and on the database webpage. Modified mitochondrial tRNA sequences from *H. sapiens* (18) and *B. taurus* (17) where added from the supplement of the respective publications. The included canonical base numberings were directly used for alignment and structural annotation.

Alignment and quality control

For all novel sequences, we used Infernal (11) to align sequences to the respective tRNA covariance models (eukaryotic, bacterial or archaeal models from tRNAscan for nuclear tRNAs, the respective MITOS2 models for mitochondrial tRNAs). When an anticodon was annotated, we compared the annotated anticodon with the one identified by Infernal (at the center of the anticodon loop). To filter tRNAs with suspicious structural features, we also checked the length of the D-loops, AC-loops, T-loops, as well as the number of unpaired bases between D-stem and AC-stem. For the AC-loop, we used a strict length of 7nt. For the other structural features, we **leveraged the maximum length** of the respective structural feature in tRNAdb 2009, and allowed a single nucleotide difference in length (resulting in 18 and 28 nt for D and T-loops respectively and 5 nt between D and AC-stem) and discarded **a total of 404 tRNAs** that did not make one of the four thresholds. As a last filtering step, all entries that were duplicates with respect to sequence and taxonomic ID were unified when necessary, so that every sequence/species pair in the tRNAdb is unique.

Statistics

The database lists 330 261 entries of mitochondrial tRNA sequences comprising 219 388 unique sequences. The vast majority of these originate in metazoa (see Fig. 4a), which reflects the distribution of mitochondrial genomes at the NCBI. The genomic part of the database contains a total of 228 955 entries comprising 93 257 unique sequences. The majority, precisely 149 749 entries, are bacterial tRNAs. However, the difference of genomic tRNA representation in tRNAdb is much smaller for unique sequences, see Fig 4b.

Mitochondrial tRNAs show a relatively even distribution with respect to their abundance, ranging from 14 692 entries for Glu to 16 734 entries for Met. The relatively even distribution can also be




Figure 4 Number of tRNA sequences per clade. a: mitochondrial tRNAs at kingdom level. b: genomic tRNAs at domain level. Light gray shows number of all sequences, dark gray number of unique sequences.




Figure 6 Distribution of species included in the tRNAdb, separated by clade. a) mitochondrial b) genomic tRNAs. Light gray shows number species in tRNAdb 2026 only, dark gray number of species in tRNAdb 2009.




Figure 5 Distribution of tRNAs per amino acid and anticodon. a) mitochondrial tRNA sequences b) genomic tRNA sequences. Light gray shows number of all sequences, dark gray number of unique sequences.

seen in the unique sequences: Asp has the most with 12 446, while Ser2 has the least amount of unique sequences with 10 184. The exception to these even distributions are Leu1 (10 857 entries, 6 209 unique sequences) and to a smaller extent Ser1 (12 979 and 9 457, resp.), which code for the two amino acids that have two tRNAs in many mitochondria. Mitochondrial tRNA Lys is the only tRNA utilizing a notable amount of a secondary anticodon (Fig 5a).

In contrast, genomic tRNAs show a much wider variety in their abundance, with the most abundant being Leu with 23 995 entries and 10 233 unique sequences. Besides the non-standard Suppressor and SeC tRNAs, His has the fewest entries with 5 290 (2 305 unique sequences), while Phe and Trp have the fewest unique sequences with 2 093 (in 5 855 and 5 579 entries, respectively). Notably, many genomic tRNAs (Asn, Asp, Cys, His, Phe, Trp, Tyr) show a clear preference for one codon (see Fig. 5b). Not surprisingly, the number of entries as well as the number of unique sequences per amino acid strongly correlates with the number of main ($n > 100$) anticodons an amino acids uses ($\rho > 0.93$, $p < 10^{-8}$ for both).

In comparison to the first version of tRNAdb, we greatly expanded sequences belonging to clades that were either not included before,

such as mitochondrial tRNAs from viridiplantae or fungi, or had only very sparse entries, such as archaea and viroidal genomic tRNAs (see Fig 6). In total, tRNAdb2026 contains genomic entries from 6 086 species, and mitochondrial entries from 15 796 different species.

Conclusion

The updated tRNAdb has been substantially expanded by an 11-fold increase in available tRNA sequences and its infrastructure was considerably modernized. Alongside a redesigned web interface, we provide a public API and a considerable increase in the number of included entries. Among these new entries now are modified mitochondrial tRNAs and, for the first time, mitochondrial tRNAs from plants and fungi. tRNAdb thus continues to serve as an important resource for the tRNA community.

However, because most entries in the tRNAdb stem from semi-automatic annotation together with manual curation, fringe cases that deviate from established tRNA properties are likely to be rejected, even though these are often of high biological interest. We therefore rely on the community to report such sequences, so they can be incorporated into future releases. This feedback may in turn motivate updated search algorithms capable of capturing a broader range of non-standard tRNAs.

Conflicts of interest

The authors declare that they have no competing interests.

Funding

This work was funded in part by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy - EXC-3105/1 - 533765739, This work is supported in part by funds from the National Funding Body (NFB: # XXXXXXXX and # YYYYYYYY).

Data availability

The data underlying the tRNAdb2026 are available in public repositories like GtRNAdb, NCBI nucleotide and BV-BRC. Links

to the sources as well as all other relevant information are provided at the detail pages of the database.

Author contributions statement

BJS developed front and back end of the database. SHB and PFS designed the new scope of the database. SHB, EC, MF, and TS searched and identified new tRNAs, FJ, MM, and JP provided data and helped porting the database. SHB, TS, and BJS performed data cleansing and data preprocessing. SHB developed the scripts for automated image generation. PFS, BJS, EC and SHB wrote the manuscript. All authors reviewed the manuscript.

Acknowledgments

References

1. T Abe, T Ikemura, Y Ohara, H Uehara, M Kinouchi, S Kanaya, Y Yamada, A Muto, and H Inokuchi. tRNADB-CE: tRNA gene database curated manually by experts. *Nucleic Acids Res.*, 37:D163–D168, 2009.
2. Helen M. Berman, John Westbrook, Zukang Feng, Gary Gilliland, T. N. Bhat, Helge Weissig, Ilya N. Shindyalov, and Philip E. Bourne. The protein data bank. *Nucleic Acids Research*, 28(1):235–242, 01 2000.
3. Patricia P. Chan and Todd M. Lowe. Gtrnadb 2.0: an expanded database of transfer rna genes identified in complete and draft genomes. *Nucleic Acids Research*, 44(D1):D184–D189, 01 2016.
4. Patricia P Chan, Brian Y Lin, Allysia J Mak, and Todd M Lowe. trnascan-se 2.0: improved detection and functional classification of transfer rna genes. *Nucleic Acids Research*, 49(16):9077–9096, 09 2021.
5. Alex Chklovski, Donovan H. Parks, Ben J. Woodcroft, and Gene W. Tyson. Checkm2: a rapid, scalable and accurate tool for assessing microbial genome quality using machine learning. *Nature Methods*, 20(8):1203–1212, 2023.
6. Peter J. A. Cock, Tiago Antao, Jeffrey T. Chang, Brad A. Chapman, Cymon J. Cox, Andrew Dalke, Iddo Friedberg, Thomas Hamelryck, Frank Kauff, Bartek Wilczynski, and Michiel J. L. de Hoon. Biopython: freely available python tools for computational molecular biology and bioinformatics. *Bioinformatics*, 25(11):1422–1423, 06 2009.
7. Alexander Donath, Frank Jühling, Marwa Al-Arab, Stephan H Bernhart, Franziska Reinhardt, Peter F Stadler, Martin Middendorf, and Matthias Bernt. Improved annotation of protein-coding genes boundaries in metazoan mitochondrial genomes. *Nucleic acids research*, 47(20):10543–10552, 2019.
8. Frank Jühling, Mario Mörl, Roland K Hartmann, Mathias Sprinzl, Peter F Stadler, and Jörn Pütz. tRNAdb 2009: compilation of tRNA sequences and tRNA genes. *Nucleic Acids Res.*, 37:D159–D162, 2009.
9. Sunghwan Kim, Jie Chen, Tiejun Cheng, Asta Gindulyte, Jia He, Siqian He, Qingliang Li, Benjamin A Shoemaker, Paul A Thiessen, Bo Yu, Leonid Zaslavsky, Jian Zhang, and Evan E Bolton. Pubchem 2025 update. *Nucleic Acids Research*, 53(D1):D1516–D1525, 01 2025.
10. National Center for Biotechnology Information. NCBI Virus. <https://www.ncbi.nlm.nih.gov/labs/virus/vssi/#/>, 2004. Accessed: 17 August 2026.
11. Eric P. Nawrocki and Sean R. Eddy. Infernal 1.1: 100-fold faster RNA homology searches. *Bioinformatics*, 29(22):2933–2935, 2013.
12. Robert Olson, Rida Assaf, Thomas Brettin, Neal Conrad, Clark Cucinell, James J Davis, Don Dempsey, Allan Dickerman, Emily M Dietrich, Ronald W Kenyon, Mehmet Kuscuglu, Elliot Lefkowitz, Jian Lu, Dustin Machi, Catherine Macken, Chunhong Mao, Anna Niewiadomska, Marcus Nguyen, Gary J Olsen, and Rick Stevens. Introducing the bacterial and viral bioinformatics resource center (bv-brc): a resource combining patric, ird and vipr. *Nucleic Acids Research*, 51:D678–D689, 11 2022.
13. Eric Sayers, Jeffrey Beck, Evan Bolton, J. Brister, Jessica Chan, Ryan Connor, Michael Feldgarden, Anna M Fine, Kathryn Funk, Jinna Hoffman, Sivakumar Kannan, Christopher Kelly, William Klimke, Sunghwan Kim, Stacy Lathrop, Aron Marchler-Bauer, Terence D Murphy, Chris O’Sullivan, Erin Schmieder, and Kim D Pruitt. Database resources of the national center for biotechnology information in 2025. *Nucleic Acids Research*, 53:D20–D29, 11 2024.
14. Conrad Schoch, Stacy Ciufo, Carol Hotton, Sivakumar Kannan, Rogneda Khovanskaya, Detlef Leipe, Richard McVeigh, Kathleen O’Neill, Barbara Robbertse, Shobha Sharma, Vladimir Soussov, John Sullivan, Lu Sun, Sean Turner, and Ilene Karsch-Mizrachi. Ncbi taxonomy: A comprehensive update on curation, resources and tools. *Database*, 2020, 08 2020.
15. Dominik Sordyl, Etienne Boileau, Agata Bernat, Satyabrata Maiti, Sunandan Mukherjee, S Naeim Moafinejad, Masoud Amiri Farsani, Anastasiya Shavina, Andrea Cappannini, Giada Agostini, Silvestro G Conticello, Filip Stefaniak, Christoph Dieterich, Elżbieta Purta, and Janusz M Bujnicki. Modomics: a database of rna modifications and related information. 2025 update and 20th anniversary. *Nucleic Acids Research*, 54(D1):D219–D225, 01 2026.
16. Mathias Sprinzl and Konstantin S. Vassilenko. Compilation of trna sequences and sequences of trna genes. *Nucleic Acids Research*, 33(suppl_1):D139–D140, 01 2005.
17. Takeo Suzuki and Tsutomu Suzuki. A complete landscape of post-transcriptional modifications in mammalian mitochondrial trnas. *Nucleic Acids Research*, 42(11):7346–7357, 06 2014.
18. Takeo Suzuki, Yuka Yashiro, Ittoku Kikuchi, Yuma Ishigami, Hironori Saito, Ikuya Matsuzawa, Shunpei Okada, Mari Mito, Shintaro Iwasaki, Ding Ma, Xuewei Zhao, Kana Asano, Huan Lin, Yohei Kirino, Yuriko Sakaguchi, and Tsutomu Suzuki. Complete chemical structures of human mitochondrial trnas. *Nature Communications*, 11:4269, 2020.