



**NTNU – Trondheim**  
Norwegian University of  
Science and Technology

# DLA based agglomeration of carbon rich aerosols

Simen Berntsen  
September 22, 2015

Specialization project applied physics  
Department of Physics  
Norwegian University of Science and Technology (NTNU).

Supervisor: Alex Hansen.

# Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>Theory</b>	<b>2</b>
2.1	On-lattice DLA . . . . .	2
2.2	Off-lattice DLA . . . . .	2
<b>3</b>	<b>Method</b>	<b>3</b>
<b>A</b>	<b>References</b>	<b>4</b>

# 1 Introduction

This is the section where the introduction to the project will be written. For now, please just read the notes:

This project has been on applying a Diffusion Limited Aggregation (DLA) approach to the clumping of aerosols in the lower atmosphere in relation to emission from fossil fuelled based combustion (etc.). Diffusion Limited Aggregation is an algorithmic approach to growth of structures where diffusion (in other words, lack of mass) is the limiting factor. It was first proposed by T. A. Witten Jr. and L. M. Sanders [2].

## 2 Theory

The fundamental algorithm of DLA is as follows; a seed particle (the particle we follow) is placed at the centre. This particle will be the basis of the growing cluster. Another particle is then introduced at some distance from this seeding particle. The starting position of this particle can in principle be infinitely far away from the particle, but for practical matters, it is often placed rather close to the seeding particle (??? is this WLOG?). This particle will perform a random walk in the space around the seeding particle, with the intention that they at some point in time, the walking particle will "hit" (i.e. be sufficiently close to) the seeding particle, and the two will clump together.

### 2.1 On-lattice DLA

In the case of on-lattice DLA, the random walk performed is usually discrete along the axes of the lattice. That means that the particle may only walk, and latch to the cluster, along these directions. Usually, the particle moves a length equivalent to the lattice constant per jump, but this may be varied.

### 2.2 Off-lattice DLA

For the off-lattice DLA, the walking particles are to restricted to walk along axes, but can move in any direction.

### **3 Method**

The algorithm used is loosely based on [1].

## A References

- [1] KUIJPERS, K. R., DE MARTÍN, L., AND VAN OMMEN, J. R. Optimizing off-lattice diffusion-limited aggregation. *Computer Physics Communications* 185, 3 (2014), 841 – 846.
- [2] WITTEN, T. A., AND SANDER, L. M. Diffusion-limited aggregation, a kinetic critical phenomenon. *Phys. Rev. Lett.* 47 (Nov 1981), 1400–1403.