Analysis of Basic Experimental Designs Using R and JASP

NE Milla, Jr.

2026-02-25

Loading libraries

```
library(readxl)
library(ExpDes)
library(agricolae)
library(tidyverse)
library(ggstatsplot)
library(car)
library(patchwork)
library(emmeans)
library(multcomp)
```

Overview of CRD & RCBD

Completely Randomized Design

- Simplest experimental design

- Frequently used to compare treatments when environmental conditions are fairly uniform (*homogeneous experimental units*)
- Each treatment is applied at *random* to several experimental units
- If the experimental units are not homogeneous, experimental error would be large and difficult to detect treatment effect
- Applicable for experiments in controlled conditions

Randomized Complete Block Design

- RCBD is one of the most widely used experimental designs
- Suited for field experiments where there exists one extraneous factor which can be used as *stratifying* variable to form homogeneous experimental units
- *Blocking* is used to minimize variability in experimental units so as to increase precision of the experiment
- Units within a block a homogeneous; units between blocks are heterogeneous
- Complete set of treatments are assigned independently at random in every block

Analysis of Variance (ANOVA)

- A method of partitioning the total variance in the response into different components which can be attributed to different sources
    - Systematic variation- effect of manipulated factors
    - Random variation- experimental error
- Method of comparing the effect of treatments on the response variable

- Method of comparing the means of three or more treatments
- **Model Assumptions**:
  - Residuals (experimental error) are *independently* distributed as $N(0, \sigma_\epsilon^2)$
  - Response is linearly related to the treatments
- One-way ANOVA for CRD experiments
- Two-way ANOVA for RCBD experiments

ANOVA for CRD: An example

```
#Importing the data
yield <- read_excel("Crop yield.xlsx")
```

ANOVA with post hoc test

```
with(yield, crd(treat = Variety,
                resp = Yield,
                quali = TRUE,
                mcomp = "tukey"))
```

```
## ------------------------------------------------------------------------
## Analysis of Variance Table
## ------------------------------------------------------------------------
##            DF    SS     MS     Fc     Pr>Fc
## Treatament  3 188.2 62.733 5.6901 0.0075605
## Residuals  16 176.4 11.025
## Total      19 364.6
## ------------------------------------------------------------------------
## CV = 13.78 %
##
## ------------------------------------------------------------------------
## Shapiro-Wilk normality test
## p-value:  0.7255893
## According to Shapiro-Wilk normality test at 5% of significance, residuals can
be considered normal.
## ------------------------------------------------------------------------
##
## ------------------------------------------------------------------------
## Homogeneity of variances test
## p-value:  0.7323728
## According to the test of bartlett at 5% of significance, residuals can be con
sidered homocedastic.
## ------------------------------------------------------------------------
##
```

```
## Tukey's test
## ------------------------------------------------------------------------
## Groups Treatments Means
## a     B   28.6
## ab    D   25
##  b    C   22.4
##  b    A   20.4
## ------------------------------------------------------------------------
```

Presenting results visually

```
#Run ANOVA the traditional way
mod1 <- aov(Yield ~ Variety,
            data = yield)
summary(mod1)
```

```
##             Df Sum Sq Mean Sq F value  Pr(>F)
## Variety      3  188.2   62.73    5.69 0.00756 **
## Residuals   16  176.4   11.02
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
#Compute summary statistics by Stocks
trt.means1 <- emmeans(mod1, specs="Variety")
trt.means1
```

```
##  Variety emmean   SE df lower.CL upper.CL
##  A         20.4 1.48 16     17.3     23.5
##  B         28.6 1.48 16     25.5     31.7
##  C         22.4 1.48 16     19.3     25.5
##  D         25.0 1.48 16     21.9     28.1
##
## Confidence level used: 0.95
```

```
#Add the letters
trt.means.cld1 <- cld(trt.means1,Letters=letters)
trt.means.cld1
```

```
##  Variety emmean   SE df lower.CL upper.CL .group
##  A         20.4 1.48 16     17.3     23.5  a
##  C         22.4 1.48 16     19.3     25.5  a
##  D         25.0 1.48 16     21.9     28.1  ab
##  B         28.6 1.48 16     25.5     31.7   b
##
## Confidence level used: 0.95
## P value adjustment: tukey method for comparing a family of 4 estimates
## significance level used: alpha = 0.05
## NOTE: If two or more means share the same grouping symbol,
##       then we cannot show them to be different.
##       But we also did not show them to be the same.
```

```
#Create graphs with SE and letters
ggplot(trt.means.cld1, aes(Variety, emmean,label = .group)) +
  geom_col(fill = "yellow", col="black") +
  geom_errorbar(aes(ymin=emmean - SE, ymax = emmean + SE),
                width = 0.2, size = 0.7) +
  geom_text(vjust=-2.1) +
  labs(x= "variety", y= "Yield") +
  lims(y = c(0,35)) +
  theme_classic()
```

```
## Warning: Using `size` aesthetic for lines was deprecated in ggplot2 3.4.0.
## ℹ Please use `linewidth` instead.
## This warning is displayed once per session.
## Call `lifecycle::last_lifecycle_warnings()` to see where this warning was
## generated.
```



Analysis of Experiments in RCBD

Importing the data (Make sure to set the appropriate working directory)

```
agrondata <- read_excel("agrondata.xlsx")
head(agrondata) #displays the 1st 6 rows of the data frame
```

```
## # A tibble: 6 × 10
##   treatment block   DTE  HT15  HT30  HT45  HT60   DTF   DTM HERBAGE
##       <dbl> <dbl> <dbl> <dbl> <dbl> <dbl> <dbl> <dbl> <dbl>   <dbl>
## 1         1     1     4  56.2 102.   152.  210.    44    60  158000
## 2         1     2     4  53.9  92.9  140   194.    44    60  192000
## 3         1     3     4  53.4  90.4  134.  189.    44    60  180000
## 4         1     4     4  52.2  89    129.  183.    44    60  155000
## 5         1     5     4  54.6 102.   150.  207.    44    60  185000
## 6         2     1     7  47.3  73.5  131.  173.    76    90  255000
```

ANOVA for RCBD experiment with post hoc

```
with(agrondata, rbd(treat = treatment,
                    block = block,
                    resp = HT15,
                    quali = TRUE,
                    mcomp = "tukey"))
```

```
## ------------------------------------------------------------------------
## Analysis of Variance Table
## ------------------------------------------------------------------------
##            DF      SS      MS      Fc    Pr>Fc
## Treatament  2 2416.67 1208.33 232.588 0.000000
## Block       4   74.27   18.57   3.574 0.059085
## Residuals   8   41.56    5.20
## Total      14 2532.50
## ------------------------------------------------------------------------
## CV = 5.7 %
##
## ------------------------------------------------------------------------
## Shapiro-Wilk normality test
## p-value:  0.1688107
## According to Shapiro-Wilk normality test at 5% of significance, residuals can
be considered normal.
## ------------------------------------------------------------------------
##
## ------------------------------------------------------------------------
## Homogeneity of variances test
## p-value:  0.07919401
## According to the test of oneillmathews at 5% of significance, the variances c
an be considered homocedastic.
## ------------------------------------------------------------------------
```

```
##
## Tukey's test
## ------------------------------------------------------------------------
## Groups Treatments Means
## a     1   54.06
##  b    2   42.6
##   c   3   23.3
## ------------------------------------------------------------------------
```

Presenting the results visually

```
#Convert integer codes to factors
agrondata <- agrondata %>%
  mutate(Trt = factor(treatment),
         Blk = factor(block))

#Run ANOVA the traditional way
mod2 <- aov(HT15 ~ Trt + Blk,
            data = agrondata)
summary(mod2)
```

```
##             Df Sum Sq Mean Sq F value   Pr(>F)
## Trt          2 2416.7  1208.3 232.588 8.17e-08 ***
## Blk          4   74.3    18.6   3.574   0.0591 .
## Residuals    8   41.6     5.2
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
#Compute summary statistics by Stocks
trt.means2 <- emmeans(mod2, specs="Trt")
trt.means2
```

```
##  Trt emmean   SE df lower.CL upper.CL
##  1     54.1 1.02  8     51.7     56.4
##  2     42.6 1.02  8     40.2     45.0
##  3     23.3 1.02  8     20.9     25.7
##
## Results are averaged over the levels of: Blk
## Confidence level used: 0.95
```

```
#Add the letters
trt.means.cld2 <- cld(trt.means2,Letters=letters)
trt.means.cld2
```

```
##  Trt emmean   SE df lower.CL upper.CL .group
##  3     23.3 1.02  8     20.9     25.7  a
##  2     42.6 1.02  8     40.2     45.0   b
##  1     54.1 1.02  8     51.7     56.4    c
##
## Results are averaged over the levels of: Blk
## Confidence level used: 0.95
## P value adjustment: tukey method for comparing a family of 3 estimates
## significance level used: alpha = 0.05
## NOTE: If two or more means share the same grouping symbol,
##       then we cannot show them to be different.
##       But we also did not show them to be the same.
```

```
#Create graphs with SE and letters
ggplot(trt.means.cld2, aes(Trt, emmean,label = .group)) +
  geom_col(fill = "lightblue", col="black") +
  geom_errorbar(aes(ymin=emmean - SE, ymax = emmean + SE),
                width = 0.2, size = 0.7) +
  geom_text(vjust=-1.5) +
  labs(x= "Treatment", y= "Mean Plant Height (15 days)") +
  lims(y = c(0,60)) +
  theme_classic()
```



Residual analysis

- To check validity of model assumptions

```
#Extract the residuals
residuals <- resid(mod2)

#Generate the normal probability plot of the residuals
qqnorm(residuals)

#Draws a reference line
qqline(residuals, col = "red")
```




```
#A formal test of normality
shapiro.test(residuals)
```

```
##
##  Shapiro-Wilk normality test
##
## data:  residuals
## W = 0.91625, p-value = 0.1688
```

```
#Runs Levene's test of equal variance (homogeneity)
leveneTest(residuals ~ Trt, data = agrondata)
```

```
## Levene's Test for Homogeneity of Variance (center = median)
##       Df F value Pr(>F)
## group  2  0.4724 0.6346
##       12
```