

Convergence of Koebe's Iteration

David Gu

Computer Science Department
Stony Brook University

gu@cs.stonybrook.edu

August 29, 2023

Convergence of Koebe Iteration Method

Koebe Iteration Algorithm

Input: Poly annulus M , $\partial M = \gamma_0 - \gamma_1 - \cdots - \gamma_n$;

Output: Conformal map $\varphi : M \rightarrow \mathbb{D}$, where \mathbb{D} is a circle domain.

- ① Compute a slit map, map the surface to the circular slit domain $f : M \rightarrow \mathbb{C}$, γ_0 and γ_k are mapped to the exterior and interior circular boundary of \mathbb{C} ;
- ② Fill the inner circle using Delaunay refinement mesh generation;
- ③ Repeat step 1 and 2, fill all the holes step by step;

Koebe Iteration Method




Figure: Slit map.

Koebe Iteration Method




Figure: Hole filling and slit map.

Koebe Iteration Method




Figure: Hole filling and slit map.

Koebe Iteration Method



Figure: All holes are filled.

Koebe Iteration Algorithm

- ④ Punch a hole at the k -th inner boundary;
- ⑤ Compute a conformal map, to map the surface onto a canonical planar annulus;
- ⑥ Fill the inner circular hole;
- ⑦ Repeat step 4 through 6, each time punch a different hole, until the process converges.

Koebe Iteration Method



Koebe Iteration Method



Koebe Iteration Method



Koebe Iteration Method



Koebe Iteration Method



Koebe Iteration Method



Koebe Iteration Method



Koebe Iteration Method



Figure: Final result.

Area, Diameter Estimate

Lemma

Suppose A is a topological annulus on \mathbb{C} , the conformal module of A is $\mu^{-1} > 1$, the exterior and interior boundaries of A are Jordan curves Γ_0 and Γ_1 , $\partial A = \Gamma_0 - \Gamma_1$, then we have the area and diameter estimates:

$$\alpha(\Gamma_1) \leq \mu^2 \alpha(\Gamma_0), \quad (1)$$

and

$$[\operatorname{diam} \Gamma_1]^2 \leq \frac{\pi}{2 \log \mu^{-1}} \alpha(\Gamma_0), \quad (2)$$

where $\alpha(\Gamma_k)$ is the area bounded by Γ_k , $k = 0, 1$.

Area, Diameter Estimate




Figure: Topological annulus with conformal module μ^{-1} .

Area, Diameter Estimate

Proof.

Let holomorphic function g maps $\{1 \leq |w| \leq \mu^{-1}\}$ to A ,

$$g(w) = w + a_0 + \frac{a_1}{w} + \frac{a_2}{w^2} + \dots$$

By Gnowell area estimate, we have

$$\alpha(\Gamma_1) = \pi \left(1 - \sum_{n=1}^{\infty} n|a_n|^2 \right)$$

$$\alpha(\Gamma_0) = \pi \left(\mu^{-2} - \sum_{n=1}^{\infty} n|a_n|^2 \mu^{2n} \right)$$

hence, this proves the area inequality (1)

$$\alpha(\Gamma_0) - \mu^{-2} \alpha(\Gamma_1) = \pi \sum_{n=1}^{\infty} n|a_n|^2 (\mu^{-2} - \mu^{2n}) \geq 0$$

Area, Diameter Estimate

Continued

The diameter $\text{diam}\Gamma_1$ is determined by $g(\{1 < |w| < \rho\})$, where $\rho \in (1, \mu^{-1})$. The diameter is bounded by half of the boundary length $g(|w| = \rho)$, we have

$$2\text{diam}\Gamma_1 \leq \int_{|w|=\rho} |g'(w)| dw = \int_0^{2\pi} |g'(\rho e^{i\theta})| \rho d\theta = \int_0^{2\pi} |g'(\rho e^{i\theta})| \sqrt{\rho} \sqrt{\rho} d\theta,$$

By Schwartz inequality, we have

$$[2\text{diam}\Gamma_1]^2 \leq \int_0^{2\pi} |g'(\rho e^{i\theta})|^2 \rho d\theta \int_0^{2\pi} \rho d\theta = 2\pi\rho \int_0^{2\pi} |g'(\rho e^{i\theta})|^2 \rho d\theta$$

Area, Diameter Estimate

Continued

Equivalent

$$\frac{2}{\pi\rho}[\operatorname{diam}\Gamma_1]^2 \leq \int_0^{2\pi} |g'(\rho e^{i\theta})|^2 \rho d\theta$$

Integrate with respect to ρ ,

$$\int_1^{\mu^{-1}} \frac{2}{\pi\rho} [\operatorname{diam}\Gamma_1]^2 d\rho \leq \int_1^{\mu^{-1}} \int_0^{2\pi} |g'(\rho e^{i\theta})|^2 \rho d\theta d\rho = \alpha(\Gamma_0) - \alpha(\Gamma_1).$$

Calculate left

$$\frac{2 \log \mu^{-1}}{\pi} [\operatorname{diam}\Gamma_1]^2 \leq \alpha(\Gamma_0) - \alpha(\Gamma_1) \leq \alpha(\Gamma_0).$$

This proves inequality (2).

Multiple Reflected Domain

Definition (Multi-reflected circle domain)

Given an m -level embedding relation tree of a circle domain C , the union of all nodes in the tree is called a multiple-reflected circle domain,

$$\Omega_m = \bigcup_{k \leq m} \bigcup_{(i)=i_1 i_2 \cdots i_k} C^{(i)} = \hat{\mathbb{C}} \setminus \bigcup_{(i)=i_1 i_2 \cdots i_m} \bigcup_{k \neq i_1} \alpha(\Gamma_k^{(i)})$$

where $\alpha(\Gamma)$ is the area bounded by Γ .

Suppose we have a holomorphic univalent map $g_m : \Omega_m \rightarrow \hat{\mathbb{C}}$, define

$$C_m := g_m(C^0)$$

$$C_m^{(i)} := g_m(C^{(i)})$$

$$\Gamma_{m,k} := g_m(\Gamma_k)$$

$$\Gamma_{m,k}^{(i)} := g_m(\Gamma_k^{(i)})$$

Symmetric Relation

According to the reflection generation tree, we have the symmetry

$$C^{i_1 i_2 \dots i_{m-1}} \mid C^{i_1 i_2 \dots i_{m-1} i_m} \quad (\Gamma_{i_m})$$

this symmetric relation is preserved by the holomorphic map g_m :

$$C_m^{i_1 i_2 \dots i_{m-1}} \mid C_m^{i_1 i_2 \dots i_{m-1} i_m} \quad (\Gamma_{m,i_m})$$

therefore g_m maps the embedding relation tree of $\{C^{(i)}\}$ to the embedding relation tree of $\{C_m^{(i)}\}$.

Hole Area Estimation

Lemma

Suppose the boundaries of C_m are $\Gamma_{m,1}, \Gamma_{m,2}, \dots, \Gamma_{m,n}$. In the m -level embedding relation tree of C_m , the total area of the holes bounded by the interior boundaries of leaf nodes is less than μ^{4m} times the total area of holes bounded by $\Gamma_{m,k}$'s,


$$\sum_{(i)=i_1 i_2 \dots i_m} \sum_{k \neq i_1} \alpha(\Gamma_{m,k}^{(i)}) \leq \mu^{4m} \sum_{i=1}^n \alpha(\Gamma_{m,i}). \quad (3)$$

Proof.

Using area estimate (1) and induction on m .



Koebe's Iteration



Koebe's Iteration

Key Observation

Given a multi-annulus \mathcal{R} , there is a biholomorphic map $g : \mathcal{C} \rightarrow \mathcal{R}$ maps a circle domain \mathcal{C} to \mathcal{R} . During the process of Koebe's iteration, the domain of the mapping \mathcal{C} can be extended to the image of the multiple reflection, (multiple reflected circle domain), which eventually covers the whole augmented complex plane $\hat{\mathbb{C}}$.

Koebe's Iteration




Figure: Reflection tree of the poly-annulus.

Koebe's Iteration

Lemma

During Koebe's iteration, at the mn -th step, the algorithm generates a univalent holomorphic function g_{mn} , its domain is extended to the m -level reflected circle domain,

$$g_{mn} : \Omega_m \rightarrow \hat{\mathbb{C}}.$$

Proof.

Initial domain is C_0 , $\infty \in C_0$, the complementary sets are

$D_{0,1}, D_{0,2}, \dots, D_{0,n}$, $\partial D_{0,i} = \Gamma_{0,i}$, $i = 1, 2, \dots, n$.

There is a biholomorphic function, $f : C_0 \rightarrow \mathcal{C}$, the complementary of \mathcal{C} is the set D_1, D_2, \dots, D_n , where D_i 's are disks, $\partial D_i = \Gamma_i$ is a canonical circle. In the neighborhood of ∞ , $f(z) = z + O(z^{-1})$. □

Koebe's Iteration

continued.

By Riemann mapping theorem, there is a Riemann mapping

$$h_1 : \hat{\mathbb{C}} \setminus D_{0,1} \rightarrow \hat{\mathbb{C}} \setminus \mathbb{D},$$

maps $\Gamma_{0,1}$ to the unit circle $\Gamma_{1,1}$, C_0 to C_1 , satisfying the normalization condition,

$$h_1(\infty) = \infty, \quad h'_1(\infty) = 1,$$

and

$$D_{1,k} = h_1(D_{0,k}), \quad k = 2, \dots, n.$$

Repeat this procedure, at $k \leq n$ step, construct a Riemann mapping,

$$h_k : \hat{\mathbb{C}} \setminus D_{k-1,k} \rightarrow \hat{\mathbb{C}} \setminus \mathbb{D},$$

which maps $\Gamma_{k-1,k}$ to the unit circle, C_{k-1} to C_k , $h_k(\infty) = \infty$ and $h'(\infty) = 1$.

Koebe's Iteration

continued.

We recursively define the symbols as follows:

$$C_k = h_k(C_{k-1})$$

$$\Gamma_{k,i} = h_k(\Gamma_{k-1,i}), i \neq k$$

$$D_{k,i} = h_k(D_{k-1,i}), i \neq k$$

$D_{k,k}$ is the unit disk \mathbb{D} , $\Gamma_{k,k}$ the unit circle. We construct a biholomorphic map $f_k : C_0 \rightarrow C_k$:

$$f_k = h_k \circ h_{k-1} \circ \cdots \circ h_1$$

and the biholomorphic map from the circle domain \mathcal{C} to C_k , $g_k : \mathcal{C} \rightarrow C_k$,

$$g_k := f_k \circ f^{-1},$$

g_k satisfies normalization condition $g_k(\infty) = \infty$, $g'_k(\infty) = 1$.

Koebe's Iteration

continued.

We generalize the domain of g_k to multiple reflected circle domain.

Because $\Gamma_{1,1}$ is a canonical circle, C_1 can be reflected about $\Gamma_{1,1}$ to C_1^1 ,

$$C_1|C_1^1 \quad (\Gamma_{1,1})$$

$h_2 : \hat{\mathbb{C}} \setminus D_{1,2} \rightarrow \hat{\mathbb{C}} \setminus \mathbb{D}$, hence h_2 is well defined on $D_{1,1}$. we denote

$$C_2^1 := h_2(C_1^1), \quad C_2^1|C_2 \quad (\Gamma_{2,1}).$$


when $k = 2, 3, \dots, n$, the Riemann mapping h_k is well defined on $C_k \cup D_{k,1}$, domain

$$C_k^1 := h_k \circ h_{k-1} \circ \dots \circ h_1(C_1^1), \quad k = 2, \dots, n,$$

satisfying

$$C_k^1|C_k \quad (\Gamma_{k,1}).$$

Koebe's Iteration



Koebe's Iteration

continued.

But the map h_{n+1} on $D_{n,1}$ is not defined. We can use Schwartz reflection to define C_{n+1}^1 . Consider the composition:

$$\beta_n := h_n \circ h_{n-1} \circ \cdots \circ h_2 : C_1 \rightarrow C_n,$$

β_n is well defined on $D_{1,1}$.

$$h_{n+1} \circ \beta_n : C_1 \rightarrow C_{n+1},$$

maps the circle $\Gamma_{1,1}$ to the circle $\Gamma_{n+1,1}$, but is not defined on $D_{1,1}$. By Schwartz reflection principle, the map $h_{n+1} \circ \beta_n$ can be extended to

$$H_{n+1} : C_1 \cup C_1^1 \rightarrow C_{n+1} \cup C_{n+1}^1,$$

where

$$C_{n+1}^1 | C_n \quad (\Gamma_{n+1,1}).$$

Koebe's Iteration

Continued.

$$\begin{array}{ccc} C_1 \cup C_1^1 & \xrightarrow{\beta_n} & C_n \cup C_n^1 \\ H_{n+1} \downarrow & & \downarrow H_{n+1} \circ \beta_n^{-1} := h_{n+1} \\ C_{n+1} \cup C_{n+1}^1 & \xrightarrow{Id} & C_{n+1} \cup C_{n+1}^1 \end{array}$$

we obtain the composition map

$$H_{n+1} \circ \beta_n^{-1} : C_n \cup C_n^1 \rightarrow C_{n+1} \cup C_{n+1}^1.$$

for convenience, we still use h_{n+1} to represent $H_{n+1} \circ \beta_n^{-1}$. Hence, we extend the domain of h_{n+1} to C_n^1 , $h_{n+1} : C_n \cup C_n^1 \rightarrow C_{n+1} \cup C_{n+1}^1$. Repeat this procedure, we conclude: for all $k \geq 1$, C_k^1 and C_k are symmetric,

$$C_k^1 | C_k \quad (\Gamma_{k,1}).$$

Koebe's Iteration

Continued.

Similarly, when $k = 2$, $\Gamma_{2,2}$ is a circle, C_2^2 and C_2 are symmetric about $\Gamma_{2,2}$. When $k > 2$, we define

$$C_k^2 := h_k \circ h_{k-1} \circ \cdots \circ h_3(C_2^2),$$

similarly, for every h_{kn+2} map, we use Schwartz reflection principle to extend analytically. For all $k \geq 2$, C_k^2 and C_k are symmetric:

$$C_k^2|C_k \quad (\Gamma_{k,2}).$$

Similarly, for any $i = 3, \dots, n$, we use Schwartz reflection principle to extend the domain and define C_k^i symmetric to C_k , for all $k \geq i$,

$$C_k^i|C_k \quad (\Gamma_{k,i}).$$

Koebe's Iteration

Continued.

After the first round of iterations, all C_k^i , $i = 1, 2, \dots, n$ are defined. Since $\Gamma_{n+1,1}$ is the unit circle, we define C_{n+1}^{i1} to be the mirror image of C_{n+1}^i with respect to $\Gamma_{n+1,1}$, $C_{n+1}^{11} = C_{n+1}$, but all other C_{n+1}^{i1} are newly generated domains $i \neq 1$. Apply the extended Riemann mapping, we get a series of mirror images:

$$C_k^{i1} | C_k^i \quad (\Gamma_{k,1}), \quad \forall k \geq n + 1, i = 2, 3, \dots, n.$$

Similarly, we can define mirror image domains:

$$C_k^{ij} | C_k^i \quad (\Gamma_{k,j}), \quad \forall k \geq n + j.$$

Koebe's Iteration

Continued.

After mn iterations, we obtain m -level mirror images $C_k^{i_1 i_2 \dots i_m}$, satisfying the symmetric relation:

$$C_k^{i_1 i_2 \dots i_m i_{m+1}} | C_k^{i_1 i_2 \dots i_m} = (\Gamma_{k, i_{m+1}}), \quad k \geq mn + i_{m+1},$$

Now the j -th boundary of $C_k^{i_1 i_2 \dots i_m i_{m+1}}$ is denoted as $\Gamma_{k,j}^{i_1 i_2 \dots i_m i_{m+1}}$,

$$\partial C_k^{i_1 i_2 \dots i_m i_{m+1}} = \Gamma_{k,i_1}^{i_1 i_2 \dots i_m i_{m+1}} - \bigcup_{j \neq i_1}^n \Gamma_{k,j}^{i_1 i_2 \dots i_m i_{m+1}}.$$

Koebe's Iteration

Continued.

Consider $g_k^{-1} = f \circ f_k^{-1}$, for all k we have

$$C = g_k^{-1}(C_k)$$

similarly,

$$C^{i_1 i_2 \dots i_m} = g_k^{-1}(C_k^{i_1 i_2 \dots i_m})$$

and its boundaries

$$\Gamma_j^{i_1 i_2 \dots i_m} = g_k^{-1}(\Gamma_{k,j}^{i_1 i_2 \dots i_m}).$$

Error Estimate

The circle domain $C = C^0$ is reflected about $\Gamma_{i_1}, \Gamma_{i_2}, \dots, \Gamma_{i_k}$ sequentially, to a k -level mirror reflection image $C^{i_1 i_2 \dots i_k}$, its interior boundary is

$$\Gamma_j^{i_1 i_2 \dots i_k} = \Gamma_j^{(i)}, \quad j \neq i_1,$$

such that i_l and i_{l+1} are not equal. After analytic extension, g_k is defined on the augmented complex plane with $n(n-1)^{k-1}$ disks removed. The boundaries of these disks are

$$\bigcup_{i_1 i_2 \dots i_k, i_l \neq i_{l+1}} \bigcup_{j \neq i_1} \Gamma_j^{i_1 i_2 \dots i_k}$$

Error Estimate

We choose a big circle Γ_ρ , enclosing all the initial boundaries Γ_j . For any point $w \in C^0$, by Cauchy's formula

$$g_k(w) - w = \frac{1}{2\pi i} \oint_{\Gamma_\rho} \frac{g_k(s) - w}{s - w} ds - \sum_{(i),j} \frac{1}{2\pi i} \oint_{\Gamma_j^{(i)}} \frac{g_k(s) - w}{s - w} ds$$

at ∞ neighborhood, $g_k(w) - w = O(w^{-1})$, when $\rho \rightarrow \infty$

$$\frac{1}{2\pi i} \oint_{\Gamma_\rho} \frac{g_k(s) - w}{s - w} ds = \frac{1}{2\pi i} \oint_{\Gamma_\rho} \frac{g_k(s) - s}{s - w} + \frac{s - w}{s - w} ds \rightarrow 0.$$

Error Estimate

Since w is outside all $\Gamma_j^{(i)}$, integration

$$\frac{1}{2\pi i} \oint_{\Gamma_j^{(i)}} \frac{w}{s-w} ds = 0,$$


for any complex number $c_j^{(i)}$, integration

$$\frac{1}{2\pi i} \oint_{\Gamma_j^{(i)}} \frac{c_j^{(i)}}{s-w} ds = 0$$

we obtain

$$g_k(w) - w = - \sum_{(i),j} \frac{1}{2\pi i} \oint_{\Gamma_j^{(i)}} \frac{g_k(s) - c_j^{(i)}}{s-w} ds$$

Multiple Reflection



Error Estimate

In the initial circle domain C^0 , let distance constant

$$\delta := \min_{i \neq j} \text{dist}(\Gamma_i, \Gamma_j^i),$$

we have $\delta > 0$. Since $\Gamma_j^{(i)} \subset \Gamma_{i_{m-1}}^{i_m}$, $|s - w| > \delta$. Define

$$\delta_{k,j}^{(i)} := \text{diam} \left(\Gamma_{k,j}^{(i)} \right),$$

the curve $\Gamma_{k,j}^{(i)} = g_k(\Gamma_j^{(i)})$ is enclosed by the circle centered at $c_j^{(i)}$ and with the diameter $\delta_{k,j}^{(i)}$, then for all $s \in \Gamma_j^{(i)}$,

$$|g_k(s) - c_j^{(i)}| \leq \delta_{k,j}^{(i)},$$

the length of the integration is $\pi \delta_j^{(i)}$, where $\delta_j^{(i)} = \text{diam}(\Gamma_j^{(i)})$.

Error Estimate

$$\begin{aligned}|g_k(w) - w| &\leq \sum_{(i),j} \frac{1}{2\pi} \oint_{\Gamma_j^{(i)}} \frac{|g_k(s) - c_j^{(i)}|}{|s - w|} |ds| \leq \sum_{(i),j} \frac{1}{2\pi} \frac{\delta_{k,j}^{(i)}}{\delta} \pi \delta_j^{(i)} \\&= \sum_{(i),j} \frac{1}{2\delta} \delta_{k,j}^{(i)} \delta_j^{(i)} \leq \sum_{(i),j} \frac{1}{4\delta} \left([\delta_j^{(i)}]^2 + [\delta_{k,j}^{(i)}]^2 \right)\end{aligned}$$

For the first term,

$$\sum_{(i),j} [\delta_j^{(i)}]^2 = \frac{4}{\pi} \sum_{(i),j} \alpha(\Gamma_j^{(i)}) \leq \mu^{4m} \sum_j \alpha(\Gamma_j) = \frac{4}{\pi} \mu^{4m} \gamma_1,$$

where $\sum_j \alpha(\Gamma_j) = \gamma_1$.

Error Estimate

For the second term, consider the topological annulus bounded by $\tilde{\Gamma}_{k,j}^{(i)}$ and $\Gamma_{k,j}^{(i)}$, by the diameter estimation (2), we obtain

$$[\delta_{j,k}^{(i)}]^2 \leq \frac{\pi}{2 \log \mu^{-1}} \alpha(\tilde{\Gamma}_{k,j}^{(i)}),$$

By inequality (3), we obtain

$$\sum_{(i).j} [\delta_{j,k}^{(i)}]^2 \leq \frac{\pi}{2 \log \mu^{-1}} \sum_{(i).j} \alpha(\tilde{\Gamma}_{k,j}^{(i)}) \leq \frac{\pi \mu^{4m}}{2 \log \mu^{-1}} \sum_j \alpha(\tilde{\Gamma}_{k,j}) = \frac{\pi \mu^{4m}}{2 \log \mu^{-1}} \gamma_2,$$

where $\gamma_2 = \sum_j \alpha(\tilde{\Gamma}_{k,j})$.

Koebe's Quarter Theorem

Theorem (Koebe Quarter Theorem)

The image of an injective analytic function $\varphi : \mathbb{D} \rightarrow \mathbb{C}$ from the unit disk \mathbb{D} onto a subset of the complex plane contains the disk whose center is $\varphi(0)$ and whose radius is $|\varphi'(0)|/4$.




Figure: Koebe's quarter theorem.

Error Estimate

We estimate γ_1 and γ_2 . The circle Γ_ρ enclose all the circles $\tilde{\Gamma}_i$, then $\gamma_1 < \pi\rho^2$. Using $g_k(w)$, we estimate γ_2 . g_k is univalent on $|w| > \rho$, in the neighborhood of ∞ , $g_k(w) = w + O(w^{-1})$. Perform coordinate change $\zeta = 1/w$, $\eta = 1/z$, construct univalent holomorphic function $\varphi : \zeta \rightarrow \eta$,

$$\varphi(\zeta) = \frac{1}{g_k(1/\zeta)},$$

φ is defined on the disk $|\zeta| < \rho^{-1}$, $\varphi(0) = 0$ and $\varphi'(0) = 1$. By Koebe 1/4 theorem,

$$\left\{ |\eta| < \frac{1}{4\rho} \right\} \subset \varphi \left(\left\{ |\zeta| < \frac{1}{\rho} \right\} \right),$$

equivalently

$$\{|z| > 4\rho\} \subset g_k(\{|w| > \rho\}),$$

hence all $\tilde{\Gamma}_{k,j}$ are included in the interior of $|z| < 4\rho$, hence the total area of all holes

$$\gamma_2 = \sum_j \alpha(\tilde{\Gamma}_{k,j}) < 16\pi\rho^2.$$

Error Estimate

We proved the convergence rate of Koebe's iteration.

Theorem (Convergence Rate of Koebe's Iteration)

In the Koebe's iteration, when $k > mn$,

$$|g_k(w) - w| \leq \frac{1}{4\delta} \left(\frac{4}{\pi} \pi \rho^2 + \frac{\pi}{2 \log \mu^{-1}} 16 \pi \rho^2 \right) \mu^{4m}.$$

This shows μ controls the convergence rate.