



| NHP ID  | DOB       | Sex | Origin             | Weight at injection | Volume of virus injection | ROA          | Nab titer at injection | Nab titer at euthanasia | Prophylactic drug administration protocol   | In-life observations                                 |
|---------|-----------|-----|--------------------|---------------------|---------------------------|--------------|------------------------|-------------------------|---|--|
| Mxxx-xx | 2025-0-00 | X   | Chinese/Indonesian | Xx.xx kg            | Xxx uL                    | Intravitreal | 1:20                   | 1:1,000                 | 1 mg/kg prednisone 1 week prior to injection through end of study; 10 uL of triamcinolone intravitreal at time of injection | Mild inflammation at 1 week clearing by 3 weeks p.i. |






Fundus imaging  
Micron X



OCT Imaging  
Heidelberg OCT



ERG recordings

|                      | -14 dpi   | 0 dpi   | 14 dpi  | 30 dpi  | 60 dpi  |
|----------------------|---|---|---|---|---|
| ERG trace            |  |  |  |  |  |
| Recording conditions |   |   |   |   |   |
| a-wave amplitude     | 16uV at 19ms  |   |   |   |   |
| b-wave amplitude     | 57uV at 34ms  |   |   |   |   |





IOP measurements (mm of Hg):  
Tonopen

| OD/OS     | 09/17/24 | 10/15/24 | 11/12/24   |
|-----------|----------|----------|------------|
| 5.75/5.85 | 6.0/6    | 4.2/4.5  | NA/3.1-3.5 |

| NHP ID  | DOB       | Sex | Origin             | Weight at injection | Volume of virus injection | ROA          | Nab titer at injection | Nab titer at euthanasia | Prophylactic drug administration protocol   | In-life observations                                 |
|---------|-----------|-----|--------------------|---------------------|---------------------------|--------------|------------------------|-------------------------|---|--|
| Mooo-00 | 2025-1-11 | X   | Chinese/Indonesian | Xx.xx kg            | Xxx uL                    | Intravitreal | 1:20                   | 1:1,000                 | 1 mg/kg prednisone 1 week prior to injection through end of study; 10 uL of triamcinolone intravitreal at time of injection | Mild inflammation at 1 week clearing by 3 weeks p.i. |







### Fundus imaging

#### Micron X






| -14 dpi  | 0 dpi   | 14 dpi   | 30 dpi  |
|--|---|--|---|
|  |  |  |  |

### OCT Imaging

#### Heidelberg OCT

|         |  |  |
|---------|--|--|
| -14 dpi |     |  |
| 0 dpi   |   |  |
| 30 dpi  |   |  |

ERG recordings

|                      | -14 dpi   | 0 dpi   | 14 dpi  | 30 dpi  | 60 dpi  |
|----------------------|---|---|---|---|---|
| ERG trace            |  |  |  |  |  |
| Recording conditions |   |   |   |   |   |
| a-wave amplitude     | 16uV at 19ms  |   |   |   |   |
| b-wave amplitude     | 57uV at 34ms  |   |   |   |   |

IOP measurements (mm of Hg):  
Tonopen

| OD/OS     | 09/17/24 | 10/15/24 | 11/12/24   |
|-----------|----------|----------|------------|
| 5.75/5.85 | 6.0/6    | 4.2/4.5  | NA/3.1-3.5 |