PRE-PUBLICATION — PUBLISH-ACCESS



**B E V E R L Y  I N D E X  L L C**

*Confidential White Paper*

# NOST-Based Mortality Forecasting

## A Conceptual Framework for Predictive Life-Course Surveillance Through Cognitive-Linguistic Signal Analysis

### Codename: THANATOS

NOUS Platform — Cross-Domain Meta-Analysis Module

**Author: Randolph R Beverly Jr**
President & Chief Architect, Beverly Index LLC
February 2026

**PROPRIETARY & CONFIDENTIAL**
*This document contains trade secrets and proprietary information of Beverly Index LLC.*

PRE-PUBLICATION — PUBLISH-ACCESS



Beverly Index LLC | Access Code: PUBLISH-ACCESS | Unauthorized distribution prohibited

PRE-PUBLICATION — PUBLISH-ACCESS



# 1. EXECUTIVE SUMMARY

This white paper introduces THANATOS, a proposed cross-domain meta-analysis module for the NOUS Platform, designed to leverage Neural Oscillation Signature Theory (NOST) for the forecasting of mortality risk through natural causes. THANATOS represents a paradigmatic extension of Beverly Index's existing diagnostic capabilities into the domain of predictive life-course surveillance.

The foundational premise is direct: if NOST can detect the presence and progression of conditions across the NOUS Platform's four established clinical domains—mental health, cerebrovascular, cardiovascular, and oncology—then the aggregation of longitudinal NOSS degradation signatures across those domains can yield statistically meaningful predictions about mortality trajectories. These four domains collectively account for approximately 75% of all natural-cause mortality in developed nations.

THANATOS does not propose to predict the moment of death. Rather, it establishes a framework for identifying accelerating multi-system cognitive-linguistic decline that correlates with elevated mortality risk within defined time horizons (6-month, 12-month, and 36-month windows). The module introduces three core analytical constructs:

- **Δ-NOSS (Delta-NOSS):** A temporal derivative measuring the rate of change in neural oscillation signature degradation across sequential analyses.
- **Mortality Convergence Index (MCI):** A composite score quantifying simultaneous multi-domain NOSS deterioration as a proxy for systemic biological decline.
- **Terminal Linguistic Marker Profile (TLMP):** A constellation of late-stage language pattern shifts—syntactic simplification, semantic narrowing, prosodic flattening, temporal-reference regression—that characterize the final phases of cognitive decline preceding death from natural causes.

The addressable market for mortality forecasting technology spans life insurance underwriting and actuarial science ($850B+ global life insurance premium market), palliative and hospice care optimization ($27B U.S. market), pharmaceutical clinical trial design, elder care and aging-in-place monitoring, and prenatal mortality prediction through integration with the GENESIS initiative (Domain 5). Conservative projections estimate THANATOS could generate $15–25M in incremental annual revenue by Year 3 of deployment, with significantly higher potential in insurance and actuarial licensing.

This document presents the theoretical foundation, technical architecture, clinical validation pathway, ethical governance framework, regulatory strategy, market analysis, and implementation timeline for the THANATOS module.



Beverly Index LLC | Access Code: PUBLISH-ACCESS | Unauthorized distribution prohibited

PRE-PUBLICATION — PUBLISH-ACCESS



# 2. THEORETICAL FOUNDATION

## 2.1 From Diagnosis to Prognosis: The NOST Continuum

NOST's established capability rests on the detection of pathological signatures embedded in language and speech patterns. Each clinical domain within the NOUS Platform maps to specific neural oscillation disruption patterns that manifest as measurable deviations in cognitive-linguistic output. The critical insight enabling THANATOS is that these deviations are not binary (present/absent) but exist on a continuum of severity that correlates with disease progression and, ultimately, with proximity to mortality.

A single NOST analysis provides a cross-sectional diagnostic snapshot—a point-in-time assessment of neural oscillation signature integrity. Repeated analyses over time, however, reveal a trajectory: the rate and pattern of signature degradation. This temporal dimension transforms NOST from a diagnostic instrument into a prognostic one. The transition requires three conceptual advances.

## 2.2 Δ-NOSS: The Temporal Derivative of Neural Decline

Delta-NOSS (Δ-NOSS) represents the first derivative of NOSS classification scores over time. Where a standard NOSS score captures the current state of neural oscillation signature integrity within a given domain, Δ-NOSS captures the velocity of change between sequential measurements.

Formally, for a given clinical domain d and time interval between analyses $t_1$ and $t_2$:

$$\Delta\text{-NOSS}(d) = [\text{NOSS}(d, t_2) - \text{NOSS}(d, t_1)] / (t_2 - t_1)$$

A negative Δ-NOSS indicates deterioration; the magnitude indicates the rate of decline. Critical thresholds can be established empirically through longitudinal validation studies, with preliminary theoretical modeling suggesting the following risk stratification:

| Δ-NOSS Range | Classification | Implied Trajectory | Monitoring Interval |
|---|---|---|---|
| > -0.5 / month | Stable | Normal aging | Annual |
| -0.5 to -1.5 / month | Declining | Accelerated deterioration | Quarterly |
| -1.5 to -3.0 / month | Critical | Rapid systemic decline | Monthly |
| < -3.0 / month | Terminal | Imminent mortality risk | Continuous |

Second-order derivatives ($\Delta^2$-NOSS) capture acceleration of decline—a critical distinction between linear deterioration (typical of chronic disease) and exponential collapse (suggesting acute decompensation or cascading system failure).



Beverly Index LLC | Access Code: PUBLISH-ACCESS | Unauthorized distribution prohibited

PRE-PUBLICATION — PUBLISH-ACCESS



## 2.3 Mortality Convergence Index (MCI)

Natural death rarely results from isolated single-system failure. More commonly, it involves cascading multi-system decline in which the failure of one biological system accelerates the deterioration of others. The Mortality Convergence Index (MCI) is designed to capture this phenomenon through the lens of cross-domain NOSS degradation.

The MCI aggregates Δ-NOSS values across all active clinical domains, weighted by domain-specific mortality contribution coefficients derived from epidemiological data:

$$MCI = \Sigma\ [w(d) \times |\Delta\text{-}NOSS(d)| \times S(d)]$$

Where w(d) is the epidemiological weighting for domain d, and S(d) is a severity scalar reflecting the absolute NOSS score (lower absolute scores amplify the MCI contribution, reflecting the increased mortality risk associated with advanced disease states).

The convergence component is captured by an interaction term that increases when multiple domains simultaneously exhibit accelerating decline. Two domains declining simultaneously contribute more than the sum of their individual contributions—a mathematical reflection of the clinical reality of multi-system failure.

| MCI Score Range | Risk Classification | Estimated Mortality Horizon |
|---|---|---|
| 0.0 – 2.0 | Low Risk | > 36 months |
| 2.0 – 5.0 | Elevated Risk | 12 – 36 months |
| 5.0 – 8.0 | High Risk | 6 – 12 months |
| > 8.0 | Critical Risk | < 6 months |

## 2.4 Terminal Linguistic Marker Profile (TLMP)

Existing research in psycholinguistics, gerontology, and palliative care has identified measurable language pattern shifts that occur in the final months of life. These shifts are not domain-specific but reflect a generalized cognitive withdrawal that precedes biological death. NOST's architecture is uniquely positioned to detect these markers because they manifest in precisely the dimensions NOST already measures.

The TLMP framework identifies seven primary terminal linguistic markers:

- **Syntactic Simplification:** Progressive reduction in sentence complexity, subordinate clause usage, and recursive grammatical structures. Mean sentence length decreases; syntactic tree depth compresses.
- **Semantic Narrowing:** Contraction of active vocabulary, reduced use of abstract and metaphorical language, increasing reliance on concrete and immediate referents.



Beverly Index LLC | Access Code: PUBLISH-ACCESS | Unauthorized distribution prohibited

PRE-PUBLICATION — PUBLISH-ACCESS



- **Temporal-Reference Regression:** Shift from future-oriented to past-oriented temporal framing. Reduction in planning language, goal-setting constructions, and prospective temporal markers.
- **Prosodic Flattening:** Decreased variation in pitch, rhythm, and intonation. Narrowing of dynamic range in speech production. Reduced emotional prosody.
- **Absolutist Intensification:** Increased use of absolute terms (“always,” “never,” “everything,” “nothing”) reflecting cognitive rigidity and reduced nuance processing.
- **Hedging Collapse:** Paradoxical reduction in uncertainty markers (“maybe,” “perhaps,” “I think”), not indicating increased certainty but rather diminished metacognitive capacity.
- **Social Disengagement Markers:** Reduced use of second-person pronouns, collaborative language, and interactional discourse markers. Increasing self-referential isolation in linguistic output.

The TLMP is computed as a composite score across these seven dimensions, calibrated against age-adjusted baselines to distinguish terminal decline from normal aging-related language change.

PRE-PUBLICATION — PUBLISH-ACCESS



Beverly Index LLC | Access Code: PUBLISH-ACCESS | Unauthorized distribution prohibited

PRE-PUBLICATION — PUBLISH-ACCESS



# 3. TECHNICAL ARCHITECTURE

## 3.1 THANATOS Module Integration

THANATOS operates as a cross-domain meta-analysis layer sitting atop the four existing NOUS Platform clinical domains. It does not require new data collection modalities; rather, it synthesizes the outputs of existing NOST analyses to generate mortality risk assessments. The module integrates with the REMIEL Engine for data processing and the Empyrean 3D visualization system for risk trajectory rendering.

The architecture consists of four processing stages:

- **Stage 1 – Longitudinal NOSS Aggregation:** Compilation of all historical NOSS scores for a given subject across all domains into a unified temporal dataset. Minimum viable longitudinal record: three analyses spaced at least 30 days apart.
- **Stage 2 – Δ-NOSS Computation:** Calculation of first- and second-order temporal derivatives for each domain. Application of smoothing algorithms to account for measurement noise and day-to-day linguistic variation.
- **Stage 3 – MCI Synthesis:** Cross-domain aggregation of Δ-NOSS values with epidemiological weighting and convergence interaction terms. Generation of composite mortality risk score.
- **Stage 4 – TLMP Overlay:** Independent assessment of terminal linguistic markers across the most recent analysis, compared against the subject's own historical baseline and population-level age-adjusted norms. TLMP score modifies the MCI-derived risk assessment upward or downward.

## 3.2 Data Requirements and Minimum Viable Analysis

THANATOS requires longitudinal data to function. A single-point analysis cannot generate a mortality forecast. The minimum viable analysis requirements are:

| Analysis Tier | Data Requirements | Forecast Capability |
|---|---|---|
| **Preliminary Risk Screen** | 3+ analyses over 90+ days, 2+ domains | Directional risk indication only |
| **Standard Mortality Forecast** | 6+ analyses over 180+ days, 3+ domains | 36-month risk horizon with confidence intervals |
| **Enhanced Mortality Forecast** | 12+ analyses over 365+ days, 4 domains | 6/12/36-month stratified risk with trajectory modeling |
| **Continuous Surveillance** | Monthly analyses, all domains, 2+ years history | Real-time risk monitoring with alert thresholds |

## 3.3 Output Specification



Beverly Index LLC | Access Code: PUBLISH-ACCESS | Unauthorized distribution prohibited

PRE-PUBLICATION — PUBLISH-ACCESS



The THANATOS module generates a structured output report containing the following elements: a THANATOS Risk Score (0–100 scale, where higher scores indicate higher mortality risk), domain-specific Δ-NOSS trajectories with trend visualization, MCI composite score with convergence mapping, TLMP assessment with marker-level detail, confidence interval and statistical significance metrics, recommended monitoring cadence, and clinical referral triggers.

All outputs are designed for interpretation by Certified NOST Analysis Practitioners (CNAPs) and are explicitly not intended for direct patient communication without clinical mediation.

PRE-PUBLICATION — PUBLISH-ACCESS



Beverly Index LLC | Access Code: PUBLISH-ACCESS | Unauthorized distribution prohibited

PRE-PUBLICATION — PUBLISH-ACCESS



# 4. CLINICAL VALIDATION PATHWAY

## 4.1 Phase 1: Retrospective Validation (Months 1–12)

The initial validation phase involves retrospective analysis of existing longitudinal datasets where both language/speech samples and mortality outcomes are available. Target datasets include large-scale aging studies (e.g., the Health and Retirement Study, the English Longitudinal Study of Ageing), clinical trial datasets with mortality endpoints, and hospice/palliative care records with documented language samples.

The objective is to establish whether NOST-analyzed language patterns from these historical samples exhibit the Δ-NOSS decay curves and TLMP profiles predicted by the theoretical model. This phase requires no new data collection and can leverage existing REMIEL Engine processing capabilities.

## 4.2 Phase 2: Prospective Cohort Study (Months 6–36)

A prospective longitudinal cohort study tracking 2,000–5,000 participants aged 55+ across multiple clinical sites. Participants undergo quarterly NOST analyses across all four domains while receiving standard medical care. Primary endpoint: all-cause mortality at 12, 24, and 36 months. Secondary endpoints: disease-specific mortality, hospitalization, and functional decline.

This study design allows for the calibration of Δ-NOSS thresholds, MCI weighting coefficients, and TLMP composite scoring against actual mortality outcomes. Statistical targets include area under the receiver operating characteristic curve (AUC-ROC) of 0.80+ for 12-month mortality prediction and 0.75+ for 36-month prediction.

## 4.3 Phase 3: Comparative Validation (Months 24–48)

Head-to-head comparison of THANATOS mortality forecasts against established mortality prediction instruments (Charlson Comorbidity Index, APACHE II/IV, Palliative Performance Scale) and traditional actuarial life tables. The objective is to demonstrate that NOST-based cognitive-linguistic mortality forecasting provides incremental predictive value beyond existing methods—particularly in the 6–36 month horizon where current instruments perform most poorly.

## 4.4 Phase 4: Real-World Evidence Generation (Months 36–60)

Integration of THANATOS into clinical workflows at partner sites to generate real-world evidence of predictive accuracy, clinical utility, and impact on care decisions. This phase supports regulatory submissions, payer negotiations, and market adoption.



Beverly Index LLC | Access Code: PUBLISH-ACCESS | Unauthorized distribution prohibited

PRE-PUBLICATION — PUBLISH-ACCESS



# 5. ETHICAL GOVERNANCE FRAMEWORK

## 5.1 The Moral Weight of Mortality Prediction

Predicting death carries extraordinary ethical gravity that exceeds any other function of the NOUS Platform. False positives could cause severe psychological harm, existential distress, and inappropriate medical decision-making. False negatives could create dangerous complacency. The THANATOS module therefore requires an ethical governance framework of commensurate rigor.

## 5.2 Core Ethical Principles

- **Non-Maleficence Above All:** THANATOS outputs must never be communicated directly to patients or subjects without clinical mediation by a qualified healthcare provider. The system is a clinical decision-support tool, not a patient-facing prognostic instrument.
- **Probabilistic Framing:** All outputs must be communicated as probability distributions, never as deterministic predictions. The language of THANATOS reporting must emphasize risk elevation, not prediction of death.
- **Intervention Orientation:** Every THANATOS risk assessment must be accompanied by actionable clinical recommendations. The purpose is to identify preventable decline and enable earlier intervention, not to forecast inevitable outcomes.
- **Informed Consent:** Subjects must provide explicit informed consent for mortality risk assessment, separate from consent for standard diagnostic NOST analysis. Consent protocols must include psychological preparation and access to counseling support.
- **Right Not to Know:** Subjects retain an absolute right to decline mortality risk information even after analysis has been performed. This right extends to family members and non-clinical third parties.
- **Anti-Discrimination Safeguards:** THANATOS outputs must never be used to deny individuals access to healthcare, insurance, employment, or other services. Contractual and technical controls must prevent discriminatory application.

## 5.3 Ethics Advisory Board

Beverly Index will establish a dedicated THANATOS Ethics Advisory Board comprising bioethicists specializing in end-of-life issues, palliative care physicians, patient advocates, health law scholars, AI ethics researchers, and representatives from underserved and minority health communities. This board will have authority to halt or modify THANATOS deployment if ethical concerns arise during validation or commercial rollout.

## 5.4 Insurance Market Ethical Considerations

The use of THANATOS in insurance underwriting presents particular ethical complexity. While life insurers represent a significant potential revenue stream, unrestricted access to mortality



Beverly Index LLC | Access Code: PUBLISH-ACCESS | Unauthorized distribution prohibited

PRE-PUBLICATION — PUBLISH-ACCESS



forecasting data could enable discriminatory underwriting practices that harm vulnerable populations. Beverly Index will adopt a tiered access model: aggregate population-level risk analytics for actuarial modeling (permitted), individual-level risk scoring for new policy underwriting (restricted, requiring ethical review), and individual-level scoring for coverage denial or premium adjustment of existing policyholders (prohibited).

PRE-PUBLICATION — PUBLISH-ACCESS



Beverly Index LLC | Access Code: PUBLISH-ACCESS | Unauthorized distribution prohibited

PRE-PUBLICATION — PUBLISH-ACCESS



# 6. REGULATORY STRATEGY

## 6.1 FDA Classification

THANATOS will likely be classified as a Class III medical device under FDA regulations, requiring Premarket Approval (PMA) rather than the 510(k) pathway available to lower-risk diagnostics. The mortality forecasting function represents a novel intended use with significant clinical risk, and there is no predicate device. Beverly Index should engage in Pre-Submission (Q-Sub) dialogue with the FDA early in the development process to confirm classification, agree on clinical evidence requirements, and establish the regulatory pathway.

## 6.2 European Regulatory Pathway (EU MDR)

Under the EU Medical Device Regulation (EU MDR 2017/745), THANATOS would likely be classified as Class IIb or Class III, requiring a Notified Body assessment and potentially clinical investigation under the Clinical Investigation Regulation. Beverly Index's existing Milan presence provides a strategic advantage for EU regulatory engagement.

## 6.3 Breakthrough Device Designation

THANATOS may qualify for FDA Breakthrough Device Designation, which provides intensive interaction and guidance on efficient device development, priority review, and a streamlined pathway to market. The novel mechanism (cognitive-linguistic mortality forecasting) and potential for significant advantage over existing alternatives support a strong application.

## 6.4 Software as a Medical Device (SaMD)

THANATOS will be regulated as Software as a Medical Device under the International Medical Device Regulators Forum (IMDRF) framework. The SaMD classification will depend on the significance of the information to the healthcare decision (critical—drives clinical management of a life-threatening condition) and the healthcare situation or condition (critical). This positions THANATOS at the highest SaMD risk category (Category IV), requiring the most rigorous clinical evidence.



Beverly Index LLC | Access Code: PUBLISH-ACCESS | Unauthorized distribution prohibited

PRE-PUBLICATION — PUBLISH-ACCESS



# 7. MARKET ANALYSIS AND REVENUE PROJECTIONS

## 7.1 Addressable Markets

| Market Segment | Global TAM | THANATOS Use Case | Revenue Model |
|---|---|---|---|
| **Life Insurance & Actuarial** | $850B+ premiums | Risk assessment, underwriting optimization | Per-assessment licensing |
| **Palliative & Hospice Care** | $27B (U.S.) | Early hospice referral, care planning | SaaS platform fee |
| **Pharmaceutical Clinical Trials** | $80B+ CRO market | Mortality endpoint proxy, patient stratification | Per-trial licensing |
| **Elder Care & Aging-in-Place** | $1.6T+ global | Continuous monitoring, decline alerts | Per-patient subscription |
| **Prenatal (GENESIS Integration)** | $45B global | Stillbirth & maternal mortality forecasting | Integrated with Domain 5 |

## 7.2 Revenue Projections

Conservative revenue projections for THANATOS, assuming regulatory clearance in Year 2 and initial market entry in Year 3:

| Revenue Stream | Year 3 | Year 4 | Year 5 |
|---|---|---|---|
| **Clinical Platform (SaaS)** | $4–6M | $8–12M | $15–20M |
| **Insurance Licensing** | $5–8M | $12–18M | $25–40M |
| **Pharma Partnerships** | $3–5M | $6–10M | $10–15M |
| **Elder Care Monitoring** | $3–6M | $8–15M | $18–25M |
| **TOTAL** | **$15–25M** | **$34–55M** | **$68–100M** |

These projections are incremental to existing NOUS Platform revenue and assume THANATOS is offered as a premium tier above standard NOST diagnostic services.



Beverly Index LLC | Access Code: PUBLISH-ACCESS | Unauthorized distribution prohibited

PRE-PUBLICATION — PUBLISH-ACCESS



# 8. INTELLECTUAL PROPERTY STRATEGY

## 8.1 Patent Applications

THANATOS introduces several patentable innovations that should be filed as continuations or new applications building on Beverly Index's existing 21 pending patent applications:

- **Delta-NOSS Temporal Analysis Method:** System and method for computing temporal derivatives of neural oscillation signature scores for prognostic health assessment.
- **Mortality Convergence Index:** System and method for multi-domain cognitive-linguistic signal aggregation to generate composite mortality risk scores.
- **Terminal Linguistic Marker Detection:** System and method for identifying terminal cognitive-linguistic patterns predictive of mortality through automated language analysis.
- **Longitudinal Cognitive Surveillance for Life-Course Risk Assessment:** System and method for continuous monitoring of cognitive-linguistic biomarkers to forecast health trajectory and mortality risk.

## 8.2 Trade Secret Protection

The specific weighting coefficients, threshold values, smoothing algorithms, and interaction terms within the Δ-NOSS, MCI, and TLMP calculations constitute trade secrets that should be protected through confidentiality controls rather than patent disclosure. These operational parameters represent the competitive moat of THANATOS and should be documented in Beverly Index's internal IP registry with appropriate access controls.

## 8.3 EU IP Considerations

Filing strategy should include European Patent Office (EPO) applications through the Patent Cooperation Treaty (PCT) pathway, leveraging Beverly Index's Milan presence for EU jurisdictional advantages. Particular attention should be given to the European approach to software patentability, which requires demonstration of "technical character"—the medical device classification of THANATOS supports this requirement.

PRE-PUBLICATION — PUBLISH-ACCESS



Beverly Index LLC | Access Code: PUBLISH-ACCESS | Unauthorized distribution prohibited

PRE-PUBLICATION — PUBLISH-ACCESS



# 9. IMPLEMENTATION TIMELINE

| Phase | Activities | Milestones |
|---|---|---|
| **Months 1–6** | Theoretical framework formalization, patent drafting, ethics board recruitment, retrospective dataset identification | White paper publication, 4 patent applications filed, ethics board seated |
| **Months 6–12** | Retrospective validation study, Δ-NOSS algorithm development, FDA Pre-Submission meeting, initial TLMP calibration | Preliminary validation results, regulatory pathway confirmed, REMIEL integration spec complete |
| **Months 12–24** | Prospective cohort study launch, MCI weighting calibration, insurance market engagement, GENESIS mortality module design | Cohort enrollment complete, first insurance LOI, GENESIS integration spec |
| **Months 24–36** | Comparative validation, Breakthrough Device application, beta deployment at clinical sites, actuarial partnership development | AUC-ROC targets met, Breakthrough designation decision, beta site feedback |
| **Months 36–48** | PMA submission preparation, real-world evidence generation, commercial pricing finalization, EU regulatory filing | PMA submitted, EU filing complete, commercial launch preparation |
| **Months 48–60** | FDA review and approval, commercial launch, insurance market entry, elder care platform deployment | Regulatory clearance, first commercial revenue, Year 3 revenue targets |

PRE-PUBLICATION — PUBLISH-ACCESS



Beverly Index LLC | Access Code: PUBLISH-ACCESS | Unauthorized distribution prohibited

PRE-PUBLICATION — PUBLISH-ACCESS



# 10. STRATEGIC POSITIONING

## 10.1 Narrative Framework

THANATOS must be positioned not as a “death prediction” tool but as a life-course surveillance system that identifies preventable decline early enough to change outcomes. The core narrative is: “We don’t predict death. We identify the trajectory toward death early enough to intervene.” This framing is essential for clinical adoption, regulatory acceptance, public trust, and ethical integrity.

## 10.2 Competitive Landscape

No existing product offers non-invasive, language-based mortality forecasting. Current mortality prediction instruments (Charlson Comorbidity Index, APACHE, Palliative Performance Scale) require clinical data inputs—lab values, diagnoses, functional assessments—that are expensive to obtain and available only in clinical settings. THANATOS’s ability to generate mortality risk assessments from cognitive-linguistic analysis alone represents a category-defining innovation with no direct competitors.

## 10.3 Integration with NOUS Platform Strategy

THANATOS represents the logical culmination of the NOUS Platform’s multi-domain architecture. By synthesizing signals across mental health, cerebrovascular, cardiovascular, and oncology domains, THANATOS transforms the NOUS Platform from a collection of domain-specific diagnostic tools into an integrated life-course health intelligence system. With the addition of GENESIS (Domain 5), THANATOS can extend mortality surveillance across the full human lifespan, from prenatal viability through end-of-life.

This positions Beverly Index not merely as a diagnostic technology company but as the architect of the world’s first comprehensive cognitive-linguistic health surveillance infrastructure—a platform that can assess, monitor, and forecast human health from the womb to the end of life through the medium of language itself.

**E N D  O F  D O C U M E N T**

Beverly Index LLC — Cheyenne, Wyoming | Milan, Italy
rrbjr@remielengine.ai | +1 307 204-6354



Beverly Index LLC | Access Code: PUBLISH-ACCESS | Unauthorized distribution prohibited