

Chapter 2: Polynomial Interpolation

Contents

1 Polynomial Interpolation	1
1.1 Van Der Monde Matrix Example	1
1.2 Lagrange Form	2

1 Polynomial Interpolation

In this chapter we study how to interpolate a data set with a polynomial.

Problem Description: Given $(n+1)$ points, say (x_i, y_i) , where $i = 0, 1, 2, \dots, n$, with distinct x_i , not necessarily sorted, we want to find a polynomial of degree n :

$$P_n(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0$$

such that it interpolates these points, i.e.,

$$P_n(x) = y_i, i = 0, 1, 2, \dots, n$$

The goal is to determine the coefficients $a_n, a_{n-1}, \dots, a_1, a_0$. Note the total number of data points is 1 larger than the degree of the polynomial.

Why should we do this?

- Find the values between the points for discrete data set
- To approximate a (probably complicated) function by a polynomial
- Then, it is easier to do computations such as derivative, integrations, etc. . .

1.1 Van Der Monde Matrix Example

Interpolate the given data set with a polynomial of degree 2:

$$\begin{array}{c|c|c|c} x_i & 0 & 1 & 2/3 \\ y_i & 1 & 0 & 0.5 \end{array}$$

Answer Let

$$P_2(x) = a_2 x^2 + a_1 x + a_0$$

We need to find the coefficients a_2, a_1, a_0 . By the interpolating properties, we have 3 equations:

$$\begin{aligned} x = 0, y = 1 : P_2(0) &= a_0 = 1 \\ x = 1, y = 0 : P_2(1) &= a_2 + a_1 + a_0 = 0 \end{aligned}$$

$$x = \frac{2}{3}, y = 0.5 : P_2\left(\frac{2}{3}\right) = \left(\frac{4}{9}\right)a_2 + \left(\frac{2}{3}\right)a_1 + a_0 = 0.5$$

Here we have 3 linear equations and 3 unknowns (a_2, a_1, a_0) .

The equations:

$$\begin{aligned} a_0 &= 1 \\ a_2 + a_1 &= 0 \\ \frac{4}{9}a_2 + \frac{2}{3}a_1 + a_0 &= 0.5 \end{aligned}$$

In matrix-vector form:

$$\begin{pmatrix} 0 & 0 & 1 \\ 1 & 1 & 1 \\ \frac{4}{9} & \frac{2}{3} & 1 \end{pmatrix} \begin{pmatrix} a_2 \\ a_1 \\ a_0 \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \\ 0.5 \end{pmatrix}$$

Easy to solve in Matlab, or do it by hand:

$$\begin{aligned} a_2 &= -\frac{3}{4} \\ a_1 &= \frac{1}{4} \\ a_0 &= 1 \end{aligned}$$

Then

$$P_2(x) = -\frac{3}{4}x^2 - \frac{1}{4}x + 1$$

1.1.1 The general case

- X: a $(n+1) \times (n+1)$ matrix, given (Van Der Monde matrix)
- a: unknown vector, with length $(n + 1)$
- y: given vector, with length $(n + 1)$

Theorem If x_i 's are distinct, then X is invertible, therefore a has a unique solution.

In Matlab, the command `vander(x)`, where x is a vector that contains the interpolation points $x = [x_1, x_2, \dots, x_n]$, will generate this matrix.

Bad News: X has a very large condition number for large n, therefore not effective to solve if n is large.

1.2 Lagrange Form

Given points: x_0, x_1, \dots, x_n

Define the **cardinal functions** $I_0, I_1, \dots, I_n : \in P^n$, satisfying the properties

Here δ_{ij} is called the Kronecker's delta.

Locally supported in discrete sense. The cardinal functions $I_i(x)$ can be written as:

Lagrange form of the interpolation polynomial can be simply expressed as

The general case. For the general case with $(n + 1)$ points, we have

$$P_n(x_i) = y_i, \quad i = 0, 1, 2, \dots, n$$

We will have $(n + 1)$ equations and $(n + 1)$ unknowns:

$$\begin{aligned} P_n(x_0) = y_0 & : x_0^n a_n + x_0^{n-1} a_{n-1} + \dots + x_0 a_1 + a_0 = y_0 \\ P_n(x_1) = y_1 & : x_1^n a_n + x_1^{n-1} a_{n-1} + \dots + x_1 a_1 + a_0 = y_1 \\ & \vdots \\ P_n(x_n) = y_n & : x_n^n a_n + x_n^{n-1} a_{n-1} + \dots + x_n a_1 + a_0 = y_n \end{aligned}$$

Putting this in matrix-vector form

$$\begin{pmatrix} x_0^n & x_0^{n-1} & \dots & x_0 & 1 \\ x_1^n & x_1^{n-1} & \dots & x_1 & 1 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ x_n^n & x_n^{n-1} & \dots & x_n & 1 \end{pmatrix} \begin{pmatrix} a_n \\ a_{n-1} \\ \vdots \\ a_0 \end{pmatrix} = \begin{pmatrix} y_0 \\ y_1 \\ \vdots \\ y_n \end{pmatrix}$$

i.e.

$$\mathbf{X} \vec{a} = \vec{y}$$

Figure 1:

$$l_i(x_j) = \delta_{ij} = \begin{cases} 1 & , \quad i = j \\ 0 & , \quad i \neq j \end{cases} \quad i = 0, 1, \dots, n$$

Figure 2:

$$\begin{aligned}
l_i(x) &= \prod_{j=0, j \neq i}^n \left(\frac{x - x_j}{x_i - x_j} \right) \\
&= \frac{x - x_0}{x_i - x_0} \cdot \frac{x - x_1}{x_i - x_1} \cdots \frac{x - x_{i-1}}{x_i - x_{i-1}} \cdot \frac{x - x_{i+1}}{x_i - x_{i+1}} \cdots \frac{x - x_n}{x_i - x_n}
\end{aligned}$$

Verify:

$$l_i(x_i) = 1$$

and for $i \neq k$

$$l_i(x_k) = 0$$

$$l_i(x_k) = \delta_{ik}.$$

Figure 3:

$$P_n(x) = \sum I_j(x) * y_j$$

It is easy to check the interpolating property:

$$P_n(x_j) = \sum I_i(x_j) * y_i = y_j$$

for every j . (The cardinal function is 1 at x_i , so y_i is multiplied by 1).

Example 2 Write the Lagrange polynomial for the data

$$\begin{array}{c|c|c|c}
x_i & 0 & 1 & 2/3 \\
y_i & 1 & 0 & 0.5
\end{array}$$

Answer We first compute the cardinal functions

$$\begin{aligned}
I_0(x) &= \frac{(x - x_1)(x - x_2)}{(x_0 - x_1)(x_0 - x_2)} = \frac{(x - 2/3)(x - 1)}{(0 - 2/3)(0 - 1)} = \frac{3}{2}(x - \frac{2}{3})(x - 1) \\
I_1(x) &= \frac{(x - x_0)(x - x_2)}{(x_1 - x_0)(x_1 - x_2)} = \frac{(x - 0)(x - 1)}{(2/3 - 0)(2/3 - 1)} = -\frac{9}{2}x(x - 1) \\
I_2(x) &= \frac{(x - x_0)(x - x_1)}{(x_2 - x_0)(x_2 - x_1)} = \frac{(x - 0)(x - 2/3)}{(1 - 0)(1 - 2/3)} = 2x(x - \frac{2}{3})
\end{aligned}$$

so

$$P_2(x) = I_0(x)y_0 + I_1(x)y_1 + I_2(x)y_2 = \frac{3}{2}(x - \frac{2}{3})(x - 1) - \frac{9}{2}x(x - 1)(0.5) + 0 = -\frac{3}{4}x^2 - \frac{1}{4}x + 1$$

Pros and cons of Lagrange polynomial:

- (+) Elegant Formula
- (-) Slow to compute, since each cardinal function is different
- (-) Not flexible: if one changes a point x_j , or add on an additional point x_{n+1} , one must recompute all cardinal functions.