

# Matrix Formulas That Is Usefull In Least Square Analysis

(You could always swap indexes properly to get similar results)

## 1 Difference of Quadratic Forms, Inverse Matrices and Solution of Linear Systems

For invertable  $(n_1 + n_2 + n_3) \times (n_1 + n_2 + n_3)$  matrix  $R$  (no need symmetric)

$$R = \begin{bmatrix} R^{(xx)} & R^{(xy)} & R^{(xz)} \\ R^{(yx)} & R^{(yy)} & R^{(yz)} \\ R^{(zx)} & R^{(zy)} & R^{(zz)} \end{bmatrix}^{-1} = \begin{bmatrix} Q^{(xx)} & Q^{(xy)} & Q^{(xz)} \\ Q^{(yx)} & Q^{(yy)} & Q^{(yz)} \\ Q^{(zx)} & Q^{(zy)} & Q^{(zz)} \end{bmatrix} \quad (1)$$

For any  $\begin{bmatrix} \vec{r}^{(x|x)} & \vec{r}^{(x|y)} & \vec{r}^{(x|z)} \end{bmatrix}$  ( $n_4 \times (n_1 + n_2 + n_3)$  matrix)

Define

(replace transpose  $\cdot^T$  to conjugate transpose  $\cdot^H$  for matrix with complex entries)

$$\begin{aligned} SS_{R[x,z]} &= \begin{bmatrix} \vec{r}^{(x|x)} & \vec{r}^{(x|z)} \end{bmatrix} \begin{bmatrix} R^{(xx)} & R^{(xz)} \\ R^{(zx)} & R^{(zz)} \end{bmatrix}^{-1} \begin{bmatrix} \vec{r}^{(x|x)} & \vec{r}^{(x|z)} \end{bmatrix}^T \\ SS_{R[x,y,z]} &= \begin{bmatrix} \vec{r}^{(x|x)} & \vec{r}^{(x|y)} & \vec{r}^{(x|z)} \end{bmatrix} \begin{bmatrix} R^{(xx)} & R^{(xy)} & R^{(xz)} \\ R^{(yx)} & R^{(yy)} & R^{(yz)} \\ R^{(zx)} & R^{(zy)} & R^{(zz)} \end{bmatrix}^{-1} \begin{bmatrix} \vec{r}^{(x|x)} & \vec{r}^{(x|y)} & \vec{r}^{(x|z)} \end{bmatrix}^T \\ \begin{bmatrix} \vec{a}^{(11)} & \vec{a}^{(12)} & \vec{a}^{(13)} \end{bmatrix} \begin{bmatrix} R^{(xx)} & R^{(xy)} & R^{(xz)} \\ R^{(yx)} & R^{(yy)} & R^{(yz)} \\ R^{(zx)} & R^{(zy)} & R^{(zz)} \end{bmatrix} &= \begin{bmatrix} \vec{r}^{(x|x)} & \vec{r}^{(x|y)} & \vec{r}^{(x|z)} \end{bmatrix} \\ \begin{bmatrix} \vec{a}_T^{(11)} & \vec{a}_T^{(12)} & \vec{a}_T^{(13)} \end{bmatrix} \begin{bmatrix} R^{(xx)} & R^{(xy)} & R^{(xz)} \\ R^{(yx)} & R^{(yy)} & R^{(yz)} \\ R^{(zx)} & R^{(zy)} & R^{(zz)} \end{bmatrix}^T &= \begin{bmatrix} \vec{r}^{(x|x)} & \vec{r}^{(x|y)} & \vec{r}^{(x|z)} \end{bmatrix} \end{aligned}$$

Then the difference of the quadratic form is

(Assume the inverse matrics in the expressions are all valid, pseudo-inverse could also work)

$$SS_{R[x,y,z]} - SS_{R[x,z]} = \vec{a}^{(12)} \left( Q^{(yy)} \right)^{-1} \left( \vec{a}_T^{(12)} \right)^T \quad (2)$$

The  $\left( Q^{(yy)} \right)^{-1}$  might be expressed as

(Swap index  $x$  and  $z$  will give another expression)

$$\begin{aligned} \left( Q^{(yy)} \right)^{-1} &= R^{(yy)} - R^{(yx)} \left( R^{(xx)} \right)^{-1} R^{(xy)} \\ &\quad - \left( R^{(yz)} - R^{(yx)} \left( R^{(xx)} \right)^{-1} R^{(xz)} \right) \left( R^{(zz)} - R^{(zx)} \left( R^{(xx)} \right)^{-1} R^{(xz)} \right)^{-1} \left( R^{(zy)} - R^{(zx)} \left( R^{(xx)} \right)^{-1} R^{(xy)} \right) \end{aligned} \quad (3)$$

The inverse matrices have relation

$$\begin{bmatrix} P^{(xx)} & P^{(xy)} \\ P^{(yx)} & P^{(yy)} \end{bmatrix} \triangleq \begin{bmatrix} R^{(xx)} & R^{(xy)} \\ R^{(yx)} & R^{(yy)} \end{bmatrix}^{-1} = \begin{bmatrix} Q^{(xx)} & Q^{(xy)} \\ Q^{(yx)} & Q^{(yy)} \end{bmatrix} - \begin{bmatrix} Q^{(xz)} \\ Q^{(yz)} \end{bmatrix} \left( Q^{(zz)} \right)^{-1} \begin{bmatrix} Q^{(zx)} & Q^{(zy)} \end{bmatrix} \quad (4)$$

$$\left( Q^{(zz)} \right)^{-1} = R^{(zz)} - \begin{bmatrix} R^{(zx)} & R^{(zy)} \end{bmatrix} \begin{bmatrix} R^{(xx)} & R^{(xy)} \\ R^{(yx)} & R^{(yy)} \end{bmatrix}^{-1} \begin{bmatrix} R^{(xz)} \\ R^{(yz)} \end{bmatrix}$$

If we solve

$$\begin{bmatrix} \vec{b}^{(11)} & \vec{b}^{(12)} \end{bmatrix} \begin{bmatrix} R^{(xx)} & R^{(xy)} \\ R^{(yx)} & R^{(yy)} \end{bmatrix} = \begin{bmatrix} \vec{r}^{(x|x)} & \vec{r}^{(x|y)} \end{bmatrix}$$

Then there is relation

(There are 6 of these relations)

$$b^{(12)} = a^{(12)} - a^{(13)} \left( Q^{(zz)} \right)^{-1} Q^{(zy)} \quad (5)$$

$$a^{(12)} \left( I - \left( Q^{(yy)} \right)^{-1} Q^{(yz)} \left( Q^{(zz)} \right)^{-1} Q^{(zy)} \right) = b^{(12)} + b_{[xz]}^{(13)} \left( Q^{(zz)} \right)^{-1} Q^{(zy)} \quad (6)$$

Note

$$\begin{aligned} \left( Q^{(zz)} \right)^{-1} Q^{(zy)} &= - \left( R^{(zy)} - R^{(zx)} \left( R^{(xx)} \right)^{-1} R^{(xy)} \right) \left( R^{(yy)} - R^{(yx)} \left( R^{(xx)} \right)^{-1} R^{(xy)} \right)^{-1} \\ &= - \begin{bmatrix} R^{(zx)} & R^{(zy)} \end{bmatrix} \begin{bmatrix} P^{(xy)} \\ P^{(yy)} \end{bmatrix} \\ I - \left( Q^{(yy)} \right)^{-1} Q^{(yz)} \left( Q^{(zz)} \right)^{-1} Q^{(zy)} &= \left( Q^{(yy)} \right)^{-1} P^{(yy)} \end{aligned}$$

Also

$$\vec{b}^{(11)} = \vec{a}^{(11)} - a^{(13)} \left( Q^{(zz)} \right)^{-1} Q^{(zx)}$$

In matrix form

$$\begin{aligned} \begin{bmatrix} \vec{b}^{(11)} & \vec{b}^{(12)} \end{bmatrix} &= \begin{bmatrix} \vec{a}^{(11)} & \vec{a}^{(12)} \end{bmatrix} - a^{(13)} \left( Q^{(zz)} \right)^{-1} \begin{bmatrix} Q^{(zx)} & Q^{(zy)} \end{bmatrix} \\ \begin{bmatrix} \vec{b}^{(11)} & \vec{b}^{(12)} \end{bmatrix} &= \begin{bmatrix} \vec{a}^{(11)} & \vec{a}^{(12)} \end{bmatrix} + a^{(13)} \begin{bmatrix} R^{(zx)} & R^{(zy)} \end{bmatrix} \begin{bmatrix} R^{(xx)} & R^{(xy)} \\ R^{(yx)} & R^{(yy)} \end{bmatrix}^{-1} \\ b^{(21)} &= a^{(21)} - a^{(23)} \left( Q^{(zz)} \right)^{-1} Q^{(zx)} \\ \vec{b}^{(22)} &= \vec{a}^{(22)} - a^{(23)} \left( Q^{(zz)} \right)^{-1} Q^{(zy)} \\ \begin{bmatrix} \vec{b}^{(21)} & \vec{b}^{(22)} \end{bmatrix} &= \begin{bmatrix} \vec{a}^{(21)} & \vec{a}^{(22)} \end{bmatrix} - a^{(23)} \left( Q^{(zz)} \right)^{-1} \begin{bmatrix} Q^{(zx)} & Q^{(zy)} \end{bmatrix} \\ \begin{bmatrix} \vec{b}^{(11)} & \vec{b}^{(12)} \\ \vec{b}^{(21)} & \vec{b}^{(22)} \end{bmatrix} &= \begin{bmatrix} \vec{a}^{(11)} & \vec{a}^{(12)} \\ \vec{a}^{(21)} & \vec{a}^{(22)} \end{bmatrix} - \begin{bmatrix} a^{(13)} \\ a^{(23)} \end{bmatrix} \left( Q^{(zz)} \right)^{-1} \begin{bmatrix} Q^{(zx)} & Q^{(zy)} \end{bmatrix} \\ \begin{bmatrix} \vec{b}^{(11)} & \vec{b}^{(12)} \\ \vec{b}^{(21)} & \vec{b}^{(22)} \end{bmatrix} &= \begin{bmatrix} \vec{a}^{(11)} & \vec{a}^{(12)} \\ \vec{a}^{(21)} & \vec{a}^{(22)} \end{bmatrix} + \begin{bmatrix} a^{(13)} \\ a^{(23)} \end{bmatrix} \begin{bmatrix} R^{(zx)} & R^{(zy)} \end{bmatrix} \begin{bmatrix} R^{(xx)} & R^{(xy)} \\ R^{(yx)} & R^{(yy)} \end{bmatrix}^{-1} \\ \begin{bmatrix} a^{(13)} \\ a^{(23)} \end{bmatrix} \begin{bmatrix} R^{(zx)} & R^{(zy)} \end{bmatrix} &= \begin{bmatrix} \vec{r}^{(x|x)} & \vec{r}^{(x|y)} \\ \vec{r}^{(y|x)} & \vec{r}^{(y|y)} \end{bmatrix} - \begin{bmatrix} \vec{a}^{(11)} & \vec{a}^{(12)} \\ \vec{a}^{(21)} & \vec{a}^{(22)} \end{bmatrix} \begin{bmatrix} R^{(xx)} & R^{(xy)} \\ R^{(yx)} & R^{(yy)} \end{bmatrix} \\ \begin{bmatrix} a^{(13)} \\ a^{(23)} \end{bmatrix} \begin{bmatrix} R^{(zx)} & R^{(zy)} \end{bmatrix} \begin{bmatrix} R^{(xx)} & R^{(xy)} \\ R^{(yx)} & R^{(yy)} \end{bmatrix}^{-1} &= \begin{bmatrix} \vec{r}^{(x|x)} & \vec{r}^{(x|y)} \\ \vec{r}^{(y|x)} & \vec{r}^{(y|y)} \end{bmatrix} \begin{bmatrix} R^{(xx)} & R^{(xy)} \\ R^{(yx)} & R^{(yy)} \end{bmatrix}^{-1} - \begin{bmatrix} \vec{a}^{(11)} & \vec{a}^{(12)} \\ \vec{a}^{(21)} & \vec{a}^{(22)} \end{bmatrix} \end{aligned}$$

Q: partial coef and coef??-should correct.

## 1.1 Partial correlation coefficient

$$\begin{bmatrix} \vec{a}^{(11)} & \vec{a}^{(12)} & \vec{a}^{(13)} \end{bmatrix} \begin{bmatrix} R^{(xx)} & R^{(xy)} & O \\ R^{(yx)} & R^{(yy)} & O \\ O & O & R^{(zz)} \end{bmatrix} = \begin{bmatrix} \vec{r}^{(x|x)} & \vec{r}^{(x|y)} & 0 \end{bmatrix}$$

$$\left(Q^{(yy)}\right)^{-1} = R^{(yy)} - R^{(yx)} \left(R^{(xx)}\right)^{-1} R^{(xy)}$$

$$x'_t = x_t - \alpha Y$$

## 2 Appendix: Formulas Used to Derive Above Results

### 2.1 Binomial inverse theorem

$$(A + UBV)^{-1} = A^{-1} - A^{-1}UB(B + BVA^{-1}UB)^{-1}BVA^{-1}$$

Specially

$$(A - BD^{-1}C)^{-1} = A^{-1} + A^{-1}B(D - CA^{-1}B)^{-1}CA^{-1}$$

### 2.2 Inverse of $(n_1 + n_2) \times (n_1 + n_2)$ matrix

(Assume the inverse matrices in the expressions are all valid)

$$\left( R^{(zy)} - R^{(zx)} \left( R^{(xx)} \right)^{-1} R^{(xy)} \right) \left( R^{(yy)} - R^{(yx)} \left( R^{(xx)} \right)^{-1} R^{(xy)} \right)^{-1}$$

Use row elimination, get

$$\begin{aligned} \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix}^{-1} &= \begin{bmatrix} (a_{11} - a_{12}a_{22}^{-1}a_{21})^{-1} & - (a_{11} - a_{12}a_{22}^{-1}a_{21})^{-1} a_{12}a_{22}^{-1} \\ - (a_{22} - a_{21}a_{11}^{-1}a_{12})^{-1} a_{21}a_{11}^{-1} & (a_{22} - a_{21}a_{11}^{-1}a_{12})^{-1} \end{bmatrix} \\ &= \begin{bmatrix} a_{11}^{-1} + a_{11}^{-1}a_{12}(a_{22} - a_{21}a_{11}^{-1}a_{12})^{-1}a_{21}a_{11}^{-1} & -a_{11}^{-1}a_{12}(a_{22} - a_{21}a_{11}^{-1}a_{12})^{-1} \\ -a_{22}^{-1}a_{21}(a_{11} - a_{12}a_{22}^{-1}a_{21})^{-1} & a_{22}^{-1} + a_{22}^{-1}a_{21}(a_{11} - a_{12}a_{22}^{-1}a_{21})^{-1}a_{12}a_{22}^{-1} \end{bmatrix} \end{aligned}$$

#### 2.2.1 Known inverse matrix, get inverse of sub matrix

Known

$$\begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix}^{-1} = \begin{bmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{bmatrix}$$

Then  $a_{22}^{-1}$  can get from

$$a_{22}^{-1} = b_{22} - b_{21}(b_{11})^{-1}b_{12}$$

### 2.3 Inverse of $(n_1 + n_2 + n_3) \times (n_1 + n_2 + n_3)$ maxtrix

$$\begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}^{-1} = UD^{-1}L$$

$$U = \begin{bmatrix} I & -a_{11}^{-1}a_{12} & a_{11}^{-1}a_{12}(a_{22} - a_{21}a_{11}^{-1}a_{12})^{-1}(a_{23} - a_{21}a_{11}^{-1}a_{13}) - a_{11}^{-1}a_{13} \\ O & I & - (a_{22} - a_{21}a_{11}^{-1}a_{12})^{-1}(a_{23} - a_{21}a_{11}^{-1}a_{13}) \\ O & O & I \end{bmatrix}$$

$$L = \begin{bmatrix} & I & & O & O \\ & -a_{21}a_{11}^{-1} & & I & O \\ (a_{32} - a_{31}a_{11}^{-1}a_{12})(a_{22} - a_{21}a_{11}^{-1}a_{12})^{-1}a_{21}a_{11}^{-1} - a_{31}a_{11}^{-1} & & - (a_{32} - a_{31}a_{11}^{-1}a_{12})(a_{22} - a_{21}a_{11}^{-1}a_{12})^{-1} & I \end{bmatrix}$$

$$D = \begin{bmatrix} a_{11} & O & O \\ O & a_{22} - a_{21}a_{11}^{-1}a_{12} & O \\ O & O & a_{33} - a_{31}a_{11}^{-1}a_{13} - (a_{32} - a_{31}a_{11}^{-1}a_{12})(a_{22} - a_{21}a_{11}^{-1}a_{12})^{-1}(a_{23} - a_{21}a_{11}^{-1}a_{13}) \end{bmatrix}$$

Note:

$$\begin{bmatrix} 1 & a_{12} & a_{13} \\ 0 & 1 & a_{23} \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} b_1 & 0 & 0 \\ 0 & b_2 & 0 \\ 0 & 0 & b_3 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ c_{21} & 1 & 0 \\ c_{31} & c_{32} & 1 \end{bmatrix} = \begin{bmatrix} b_1 + a_{12}b_2c_{21} + a_{13}b_3c_{31} & a_{12}b_2 + a_{13}b_3c_{32} & a_{13}b_3 \\ b_2c_{21} + a_{23}b_3c_{31} & b_2 + a_{23}b_3c_{32} & a_{23}b_3 \\ b_3c_{31} & b_3c_{32} & b_3 \end{bmatrix}$$