



Figure C. Several phenotypes under one rule: what is observed, and how far it is checked

ONE LANE: evidence from Lane 6 (PR #11, merged @ f72db9e); no second-lane reproduction yet.
Claim statuses checked against research/claims.md @ 76a8dc9.



b Resolution and timestep checks: the coexistence band moves, and at R 26 the circler's own rule falls outside it



c At the coexistence rule ($\mu 0.155, \sigma 0.020$): persistence is not the same as a verified attractor

	persistence at the coexistence rule (one lane)			same lane, stronger tests		independent	
	T10 R13	T10 R26	T40 R13	clean rerun	returns after 1001 attenuation	second lane reproduces	catalog identity
glider (Orbium cells)	glider m 0.487	glider m 0.487	→ static m 0.387	no trace at this rule	≤ 10% (2 phases)	not yet	Orbium
circler S102	circler m 0.523	died	circler m 0.497	bitwise S102-5bfac8f95f	at 5%: kept in 1 of 2 phases	not yet	OG2g
static ring S103	static m 0.379	static m 0.379	static m 0.379	bitwise S103-1d8c158cdd	bitwise, ≤ 20% (1 phase, dossier)	not yet	C01a

● glider ■ static ■ filled the world ◆ circler ● died ▨ glider + circler coexist (location shifts with T, R)

C038 · reproduced C039 · num. fragile C040 · observed C042 · reproduced C043 · observed

····· $\sigma = 0.020$ (coexistence rule) | sampled σ without coexistence