
Key Proofs

Bart Snapp and Brad Findell

January 4, 2019

Contents

Introduction	3
The Ximera Environment	3
I Math 1	5
Vertical Angles	5
Parallel Lines	7
Isosceles Triangle Theorem	9
Quadrilaterals	13
Inscribed Angles	18
II Math 2	20
Similar Right Triangles	20
Midsegment Theorem	22

Introduction

Online proof project description.

Student performance is generally quite poor on Ohio's end-of-course exams for Algebra 1, Geometry, Math 1, and Math 2, especially on items involving proof. In response to concerns that the items are too difficult, the following pages provide examples of more accessible computer-scorable proof items, mostly in Geometry. The proofs are written to focus on the most important steps and reasons in the argument.

For the convenience of teachers using an integrated curriculum, items are separated into two groups: those appropriate for Math 1, and those appropriate for Math 2, according to Ohio's assessments.

Fix note: This is work in progress.

Draft items are first written as complete proofs in order to consider which expressions, words, or phrases students might be prompted to enter.

Questions to reviewers are written in red. Please send comments to Brad Findell, findell.2@osu.edu, Department of Mathematics, The Ohio State University.

The Ximera Environment

Students complete the proofs by filling in blanks, pulling down menus, and selecting correct answers. In Ximera, some answers are checked automatically when they are chosen. Others answers require pressing Enter, clicking the blue question mark, or clicking the blue "Check Work" button. Give it a try!

Example 1. *Some problems are multiple-choice:*

Multiple Choice:

- (a) *Don't pick me.*
- (b) *Not me either.*
- (c) *Pick me!*

Author(s): Brad Findell

(d) *Also an incorrect choice*

Example 2. *Some problems are select-all that are correct:*

Select All Correct Answers:

(a) *Don't pick me.*

(b) *Pick me!*

(c) *Pick me too!*

(d) *I'm a correct choice too.*

Example 3. *Some problems use (purple haze/ purple rain/ pull-down menus).*

Example 4. *Some problems are fill in the blank: $3 \times 2 = \boxed{?}$*

Part I


Math 1

Vertical Angles

Proofs updated.

Fix note: Below are three different proofs. Please consider them separately. And in each proof, which of the details should be included, and which should be omitted?

Problem 1 Point P is the intersection of lines m and n . Prove that $\angle 1 \cong \angle 3$.



Fix note: When students write equations about linear pairs, they often write two equations for non-overlapping linear pairs—which doesn't help. The figure above is intended to help avoid that dead end, but it would be worthwhile to discuss that dead end anyway.

- (a) $\angle 1 \cong \angle 3$ because they are both (complementary/ supplementary/ opposite / congruent) to $\angle 2$.

Detail: First write down equations about linear pairs of angles:

$$m\angle 1 + m\angle 2 = 180^\circ$$

$$m\angle 3 + m\angle 2 = 180^\circ$$


By comparing the two equations, some students will see clearly that $m\angle 1 = m\angle 3$. A more formal approach would be to do some algebra.

- (b) A rotation of $(90^\circ / 180^\circ / 360^\circ)$ about P maps m onto itself, maps n onto itself, and swaps $\angle 1$ and $\angle 3$. Because rotations preserve angle measures, it must be that $\angle 1 \cong \angle 3$.

Detail: Line m is the union of two opposite rays with endpoint P . Check that the 180° rotation about P swaps these opposite rays. The same idea holds for line n so that together the sides of $\angle 1$ become the sides of $\angle 3$ and vice versa.

- (c) Reflecting about the (bisector/ supplement/ opposite) of $\angle 2$ swaps $\angle 1$ and $\angle 3$. Because reflections preserve angle measures, it follows that $\angle 1 \cong \angle 3$.

Detail: The reflection swaps the two rays that are the sides of $\angle 2$. Because reflections take lines to lines, that reflection must swap not just the rays but lines m and n .



Parallel Lines

Proofs updated.


Parallel postulate: Given a line and a point not on the line, there is exactly one line through the given point parallel to the given line.

Theorems to prove:

- (a) A 180° rotation about a point on a line takes the line to itself.
- (b) A 180° rotation about a point not on a line takes the line to a parallel line.
- (c) If two parallel lines are cut by a transversal alternate interior and corresponding angles are congruent.
- (d) If two lines are cut by a transversal so that alternate interior (or corresponding) angles are congruent, then the lines are parallel.

Fix note: Below are two different proofs. Please consider them separately. And in each proof, which of the details should be included, and which should be omitted?

Problem 2 Given that parallel lines m and n are cut by transversal k , prove that alternate interior angles are congruent.



First proof:

Author(s): Brad Findell

- (a) Let A and B be the intersections of transversal k with lines m and n , respectively.
- (b) Let P be the midpoint of \overline{BC}
- (c) Rotate 180° about P , which takes k to itself.
- (d) The rotation maps B to C and C to B because distances are preserved.
- (e) The rotation maps m to a parallel line through C , which must be n by the uniqueness of parallels.
- (f) The rotation maps n to m by the same reasoning.
- (g) The rotation swaps $\angle 2$ and $\angle 3$. These alternate interior angles must be congruent because the rotation preserves angle measures.

Second proof:


- (a) Let A and B be the intersections of transversal k with lines m and n , respectively.
 - (b) Translate line m along line k to the right by distance BC .
 - (c) The translation maps B to C , and it maps m to n because the translation maintains parallels.
 - (d) The translation maps $\angle 1$ to $\angle 3$. These corresponding angles must be congruent because the translation preserves angle measures.
-

Isosceles Triangle Theorem

Proofs updated.

Fix note: Below are several different proofs, along with one that is not a proof. Please consider them separately.
Any (or all) of the proofs might be extended to conclude that, in the case of an isosceles triangle, the perpendicular bisector, angle bisector, median, and altitude all lie on the same line.

Problem 3 Prove that the base angles of an isosceles triangle are congruent.




- (a) Beginning with the given figure on the left, Morgan draws \overline{CD} and marks the figure intending that this new segment is a(n) (median/ angle bisector / perpendicular bisector/ altitude).
- (b) Based on the marked figure, Morgan claims that the $\triangle ACD \cong \triangle \boxed{?}$ by (SAS/ SSS/ SSA/ ASA/ HL).
- (c) Finally, Morgan concludes that $\angle A \cong \angle \boxed{?}$, as they are corresponding parts of congruent triangles.

Problem 4 Prove that the base angles of an isosceles triangle are congruent.


Author(s): Brad Findell

Isosceles Triangle Theorem



- Beginning with the given figure on the left, Deja draws \overline{CD} and marks the figure intending that this new segment is a(n) (median / angle bisector / perpendicular bisector / altitude).
- Based on the marked figure, Deja claims that the $\triangle ACD \cong \triangle \boxed{?}$ by (SAS / SSS / SSA / ASA / HL).
- Finally, Deja concludes that $\angle A \cong \angle \boxed{?}$, as they are corresponding parts of congruent triangles.


Problem 5 Prove that the base angles of an isosceles triangle are congruent.



- Beginning with the given figure on the left, Elle draws \overline{CD} and marks the figure intending that this new segment is a(n) (median / angle bisector / perpendicular bisector / altitude).
- Based on the marked figure, Deja claims that the $\triangle ACD \cong \triangle \boxed{?}$ by (SAS / SSS / SSA / ASA / HL).
- Finally, Deja concludes that $\angle A \cong \angle \boxed{?}$, as they are corresponding parts of congruent triangles.

Isosceles Triangle Theorem

Problem 6 Simon and Taylor are trying to prove that the base angles of an isosceles triangle are congruent.



Beginning with the given figure on the left, Simon draws \overline{CD} and marks the second figure intending that this new segment is a perpendicular bisector of \overline{AB} .

Taylor claims that a perpendicular bisector of a side of a triangle usually misses the opposite vertex. So without using other properties of isosceles triangles or perpendicular bisectors, the figure should allow for that possibility.


Fix note: Taylor's claim needs some work, as is the case for the choices below.

Choose the best response to their argument:

Multiple Choice:

- (a) Simon is correct, and $\triangle ACD \cong \triangle BCD$ by SAS.
- (b) Simon is correct, and $\triangle ACD \cong \triangle BCD$ by SSS
- (c) Taylor is correct, and the perpendicular bisector cannot be used to complete this proof.
- (d) Neither of them are correct.

Problem 7 Prove that the base angles of an isosceles triangle are congruent.



Isosceles Triangle Theorem


- (a) Examining the given figure on the left, Lissy notices symmetry in the triangle and claims that the triangle is congruent to itself by a (translation / reflection / rotation).
 - (b) Based on the marked figure, Lissy claims that the $\triangle ACB \cong \triangle \boxed{?}$ by (SAS / SSS / SSA / ASA / HL).
 - (c) Finally, Lissy concludes that $\angle A \cong \angle \boxed{?}$, as they are corresponding parts of congruent triangles.
-

Quadrilaterals

Proof.

Problem 8 Adapted from Ohio's 2017 Geometry released item 13.

Two pairs of parallel lines intersect to form a parallelogram as shown.




Complete the following proof that opposite angles of a parallelogram are congruent:

- (a) $\angle 1 \cong \angle 2$ as (opposite angles / alternate interior angles / corresponding angles) for parallel lines (m and n / k and l).
- (b) $\angle 3 \cong \angle 2$ as (opposite angles / alternate interior angles / corresponding angles) for parallel lines (m and n / k and l).
- (c) Then $\angle 1 \cong \angle 3$ because they are both congruent to $\angle 2$.

Problem 9 Adapted from Ohio's 2018 Geometry released item 21.

Given the parallelogram $WXYZ$, prove that $\overline{WX} \cong \overline{YZ}$.

Author(s): Brad Findell




Fix note: It really would help to have an online environment that allows students to mark diagrams.

Complete the proof below:

- (a) $\angle ZWY \cong \angle XYW$ as (alternate interior angles / corresponding angles / opposite angles) for parallel segments (\overline{WZ} and \overline{XY} / \overline{WX} and \overline{YZ}).
- (b) $\angle ZYW \cong \angle XWY$ for the same reason, this time for parallel segments (\overline{WZ} and \overline{XY} / \overline{WX} and \overline{YZ}).
- (c) $\overline{WY} \cong \overline{YW}$ because a segment is congruent to itself.
- (d) $\triangle WYZ \cong \triangle YWX$ by (SAS / ASA / SSS).
- (e) Then $\overline{YZ} \cong \overline{WX}$ as corresponding parts of congruent triangles.

Fix note: Maybe number the angles.

Problem 10 Use symmetry to prove properties of parallelograms.



Consider a 180° rotation about M , the midpoint of diagonal \overline{AC} . Show that this rotation maps the parallelogram onto itself.

Fix note: The following details are subtle, especially distinguishing between mapping the sides (i.e., segments) and the lines containing the sides. Can any of this be omitted or abbreviated? Which parts might students supply?

- (a) The rotation maps A to C and C to A because a 180° rotation about a point on a line takes the line to itself and preserves lengths.
- (b) The rotation maps \overleftrightarrow{AB} to a parallel line through C (the image of A), which by the uniqueness of parallels must be \overleftrightarrow{CD} . Similarly, the rotation maps \overleftrightarrow{CD} to \overleftrightarrow{AB} , \overleftrightarrow{AD} to \overleftrightarrow{CB} , and \overleftrightarrow{CB} to \overleftrightarrow{AD} .
- (c) Furthermore, the intersection of \overleftrightarrow{AB} and \overleftrightarrow{CD} , which is B , must map to the intersection of their images, \overleftrightarrow{CD} and \overleftrightarrow{AD} , which is D . And likewise, D must map to B .
- (d) Because vertices are mapped to vertices, sides are mapped to opposite sides, angles are mapped to opposite angles, and thus the parallelogram is mapped onto itself.

Because the 180° rotation about M maps the parallelogram onto itself, we can conclude that


Now we see that the symmetry provides the properties for free:

- opposite sides are congruent (sides are mapped to opposite sides),
- opposite angles are congruent (angles are mapped to opposite angles), and
- the diagonals bisect each other.

Detail: The 180° rotation about M swaps \overrightarrow{MB} and \overrightarrow{MD} , so they must be opposite rays, and thus B , M , and D are collinear.

Because the rotation preserves lengths, $MB = MD$, so that M is also the midpoint of \overline{BD} , which means that the diagonals bisect each other.


Problem 11 Quadrilateral $ABCD$ is a kite as marked. Prove that \overleftrightarrow{BD} is the perpendicular bisector of \overline{AC} .



Key theorem: The points on a perpendicular bisector are exactly those that are equidistant from the endpoints of a segment.

Because B and D are each equidistant from A and C, they each must lie on the perpendicular bisector of \overline{AC} , which implies that \overleftrightarrow{BD} is its perpendicular bisector.

Problem 12 Another proof:



Fix note: Do we need a step about \overleftrightarrow{BX} and \overleftrightarrow{BD} being the same line?


Fix note: Here, for the record, are the steps for a paragraph proof.

- (a) $\overline{BD} \cong \overline{BD}$, so that $\triangle BAD \cong \triangle BCD$ by SSS.
- (b) $\angle ABD \cong \angle CBD$ by CPCTC.
- (c) $\overline{BX} \cong \overline{BX}$, so that $\triangle ABX \cong \triangle CBX$ by SAS.
- (d) $\angle BXA \cong \angle BXC$ by CPCTC, and they are a linear pair, so $\overline{BX} \perp \overline{AC}$.
- (e) $\overline{AX} \cong \overline{CX}$ by CPCTC, so X is the midpoint of \overline{AC} .
- (f) Thus, \overleftrightarrow{BD} is the perpendicular bisector of \overline{AC} .


Inscribed Angles

Proofs updated.

Problem 13 In the figure below, \overline{AC} is a diameter of a circle with center P . Prove that $\angle ABC$ is a right angle.



- (a) Beginning with the diagram on the left, Natalia draws \overline{PB} and marks the diagram to show segments that she knows to be congruent because each one is a $\boxed{?}$ of the circle.



- (b) Natalia sees that $\triangle APB$ and $\triangle BPC$ are $\boxed{?}$ triangles, so she marks the figure to show angles that must congruent.

Fix note: Do we need a statement or citation of the theorem?

- (c) In order to do some algebra with these congruent angles, Natalia labels their measures x and y , as shown in the picture on the right.
- (d) She writes an equation for the sum of the angles of $\triangle ABC$:


$$\boxed{?} = 180^\circ$$

Fix note: Need a prompt about dividing the equation by 2.

(e) Since $m\angle ABC = \boxed{?}$, she concludes that $m\angle ABC = 90^\circ$.

Problem 14 A special case of the relationship between an inscribed angle and the corresponding central angle.

In the figure below, \overline{AC} is a diameter of a circle with center P . Prove that $z = 2x$.



Because z is the measure of an angle exterior to $\triangle APB$, it is equal to the sum of the measures of the remote interior angles. In other words $z = 2x$.

Alternatively, without using the exterior angle theorem, one might proceed as follows:

- (a) $\angle ABP + x + x = 180^\circ$ because of the angle sum in $\triangle ABP$.
- (b) $\angle ABP + z = 180^\circ$ because they form a linear pair.
- (c) Then $z = 2x$ by comparing the two equations.

Fix note: This handles the special case in which one side of the inscribed angle is a diameter. For the general result, consider two cases: (1) When the center of the circle is in the interior of the inscribed angle; and (2) When the center of the circle is not in the interior of the inscribed angle.


Part II

Math 2

Similar Right Triangles

Proofs updated.

Problem 15 *Adapted from Ohio's 2017 Geometry released item 17.*




Complete the following proof that $\triangle DAC$ is similar to $\triangle DBA$:

- (a) $\triangle ABC \sim \triangle \square$ by (AA similarity / CPCTC / right triangle similarity) because they share $\angle B$ and they each have a right angle.
- (b) $\triangle ABC \sim \triangle DAC$ for the same reason because they share ($\angle A / \angle B / \angle C$) and they each have a right angle.
- (c) $\triangle DAC \sim \triangle \square$ because they are both similar to $\triangle ABC$.

Problem 16 *A different proof, also adapted from Ohio's 2017 Geometry released item 17.*

Author(s): Brad Findell

Similar Right Triangles



Complete the following proof that $\triangle DAC$ is similar to $\triangle DBA$:

- (a) $\angle B$ and $\angle BAD$ are complementary because they are acute angles in a right triangle.
- (b) $\angle DAC$ and $\angle BAD$ are complementary because they are adjacent angles that form $\angle BAC$, which is right.
- (c) $\angle B \cong \angle DAC$ because they are both complementary to $\angle BAD$.
- (d) $\triangle DAC \sim \triangle \boxed{?}$ by (AA similarity / CPCTC / right triangle similarity) because $\angle B \cong \angle DAC$ and they each have a right angle.

Midsegment Theorem

Proofs updated.

Definition 1. In a triangle, a midsegment is a line joining the midpoints of two sides.

Theorem 1. Midsegment Theorem: A midsegment in a triangle is parallel to and half the length of the corresponding side.


Fix note: The typical proof uses similarity. This one uses parallelograms.

In preparation for the midsegment theorem, the class proved some useful theorems about parallelograms.

Theorem 2. Antonia's Theorem: If the diagonals of a quadrilateral bisect each other, then the quadrilateral is a parallelogram.

Theorem 3. Lu's Theorem: If one pair of sides of a quadrilateral are congruent and parallel, then the quadrilateral is a parallelogram.

Problem 17 To prove the midsegment theorem for $\triangle ABC$ with midpoints D and E of sides AC and BC , respectively, Jesse extends \overline{DE} to a point X such that $EX = DE$, as shown in the marked figure.




Jesse organizes her reasoning in the following flow chart:

Fix note: The flowchart omits reasons to reduce clutter. The most significant steps are green whereas the details are blue.

Author(s): Brad Findell

Midsegment Theorem



Fix note: For the following questions, does it help to name the theorems? Would theorem numbers be better? Or maybe we should write each theorem in words.

In the proof above, why may Jesse conclude that quadrilateral $CDBX$ a parallelogram?

Multiple Choice:

- (a) By the Pythagorean Theorem.
- (b) By Lu's Theorem.
- (c) By Antonia's Theorem.
- (d) By the Parallelogram Theorem.

In the proof above, why may Jesse conclude that quadrilateral $DABX$ a parallelogram?

Multiple Choice:

- (a) By the Pythagorean Theorem.
- (b) By Lu's Theorem.
- (c) By Antonia's Theorem.
- (d) By the Parallelogram Theorem.

Fix note: Here, for the record, are sentences for a paragraph proof.

- (a) $CE = BE$ and $DX = EX$, as given.
- (b) Quadrilateral $CDBX$ is a parallelogram by Antonia's Theorem.
- (c) $XB = CD$ because opposite sides of a parallelogram are congruent.
- (d) $XB = AD$ because they are both equal to CD .
- (e) $\overline{XB} \parallel \overline{CD}$ because $CDBX$ is a parallelogram.
- (f) $\overline{XB} \parallel \overline{AD}$ because A , C , and D are collinear.
- (g) Quadrilateral $DABX$ is a parallelogram by Lu's Theorem.
- (h) $\overline{DX} \parallel \overline{AB}$ because $DABX$ is a parallelogram.
- (i) $\overline{DE} \parallel \overline{AB}$ because D , E , and X are collinear.
- (j) $AB = DX = DE + EX = 2DE$
- (k) $DE = \frac{1}{2}AB$
- (l) Summary: DE is parallel to AB and half its length.