


Genomic Identification and Laboratory Characterization of a Putative Transcriptional Repressor of Virulence in *Pseudomonas aeruginosa* UCBPP-PA14

Patrick K. Taylor^{#1}, Wing Yin Venus Lau^{#1}, Baofeng Jia¹, Fiona S. L. Brinkman^{*1}, and Amy H. Y. Lee^{*1}

Authors contributed equally to this work, *Co-contributing senior authors

1. Department of Molecular Biology and Biochemistry, Simon Fraser University, Canada.

Antivirulence as an alternative approach to antibiotics¹



Pathogen-associated genes (PAGs) as antivirulence drug targets

Genes found exclusively in bacterial pathogens, but not in related non-pathogens²

- Likely associated with virulence-related functions
- Likely not essential for bacterial growth and survival
- Likely more pathogen-specific drug targets


Pseudomonas aeruginosa – a WHO priority pathogen^{3,4}

- Gram-negative, ecologically diverse, opportunistic plant and animal pathogen
- Common cause of morbidity and mortality in immunocompromised patients
- Broad range of intrinsic and extrinsic AMR Mechanisms

Hypothesis:

Novel antivirulence drug targets can be discovered via comparative genomic analyses to identify genes solely present in pathogens and potentially under positive selection.

Genus-specific approach of detecting PAGs: *Pseudomonas*



Three methods of PAG identification to increase accuracy


55 *Pseudomonas aeruginosa* specific PAGs were identified and prioritized, 3 with evidence of positive selection in *Pseudomonas aeruginosa* PA14



A) A total of 55 PAGs were identified by all 3 methods in *P. aeruginosa*.

B) Of the 55 PAGs, 17 were found in more than 3 genomes in the reduced dataset, 3 of which showed positive selection in *P. aeruginosa* PA14.

PA14_RS12700 knockout does not affect antimicrobial sensitivity and biofilm formation, but is associated with hyper-virulence phenotype



A) The PA14_RS12700 transcriptional repressor knockout strain showed no difference in susceptibility to β -lactam antibiotics or heavy metals.

B) PA14_RS12700 displays a hyper-virulent phenotype in a *C. elegans* infection model (Gehan-Breslow-Wilcoxon test; $p < 0.0001$)

SUMMARY & FUTURE DIRECTIONS

- Using genomics, we identified and prioritized 17 PAGs in the *P. aeruginosa* PA14 genome that are conserved in *Pseudomonas* pathogens but not in *Pseudomonas* non-pathogens.
- Two genes of interest, PA14_RS12700 and PA14_RS12695, have sequence similarities with Bla/Cop/Mec sensor-regulator systems for β -lactam and heavy metal resistance in Gram positives.
- In *P. aeruginosa* PA14, loss of the transcriptional repressor, PA14_RS12700, results in increased virulence, but no difference in resistance observed.
- **Computationally predicted PAGs warrant further study including possible development as targets for novel antivirulence therapeutics.** However, there is a need to understand their mechanisms of action.
- Future plan: Characterize an overexpression strain of the BlaR-like component, PA14_RS12695.

Funding & Acknowledgements

SFU

CIHR IRSC
Canadian Institutes of Health Research

NSERC CRSNG
Natural Sciences and Engineering Research Council of Canada

GenomeCanada

GenomeAtlantic

Genome British Columbia

References

1. Lau WYV, et al. eBioMedicine. 2023;88:104429. doi: 10.1016/j.ebiom.2022.104429
2. Ho Sui SJ, et al. (2009) <https://doi.org/10.1371/journal.pone.0008094>
3. Von Wintersdorff CJH, et al. Front Microbiol. 2016;7:173. fmicb.201
4. <https://www.who.int/news/item/27-02-2017-who-publishes-list-of-bacteria-for-which-new-antibiotics-are-urgently-needed>



Scan For Copy