

the 1990s, the incidence of *S. flexneri* has increased in the United Kingdom [10]. In the United States, *S. flexneri* has been reported as the most common serotype of *S. flexneri* in the 1990s [11].

There is a paucity of data on the epidemiology of *S. flexneri* in the United Kingdom. In the 1980s, *S. flexneri* was the most commonly isolated serotype of *S. flexneri* in the United Kingdom [12]. In the 1990s, *S. flexneri* was the most commonly isolated serotype of *S. flexneri* in the United Kingdom [13].

The purpose of this study was to determine the prevalence of *S. flexneri* in the United Kingdom. The study was conducted in the United Kingdom, where *S. flexneri* is the most commonly isolated serotype of *S. flexneri* in the United Kingdom [12].

The study was conducted in the United Kingdom, where *S. flexneri* is the most commonly isolated serotype of *S. flexneri* in the United Kingdom [12]. The study was conducted in the United Kingdom, where *S. flexneri* is the most commonly isolated serotype of *S. flexneri* in the United Kingdom [12].

The study was conducted in the United Kingdom, where *S. flexneri* is the most commonly isolated serotype of *S. flexneri* in the United Kingdom [12]. The study was conducted in the United Kingdom, where *S. flexneri* is the most commonly isolated serotype of *S. flexneri* in the United Kingdom [12].

The study was conducted in the United Kingdom, where *S. flexneri* is the most commonly isolated serotype of *S. flexneri* in the United Kingdom [12]. The study was conducted in the United Kingdom, where *S. flexneri* is the most commonly isolated serotype of *S. flexneri* in the United Kingdom [12].

The study was conducted in the United Kingdom, where *S. flexneri* is the most commonly isolated serotype of *S. flexneri* in the United Kingdom [12]. The study was conducted in the United Kingdom, where *S. flexneri* is the most commonly isolated serotype of *S. flexneri* in the United Kingdom [12].

The study was conducted in the United Kingdom, where *S. flexneri* is the most commonly isolated serotype of *S. flexneri* in the United Kingdom [12]. The study was conducted in the United Kingdom, where *S. flexneri* is the most commonly isolated serotype of *S. flexneri* in the United Kingdom [12].

The study was conducted in the United Kingdom, where *S. flexneri* is the most commonly isolated serotype of *S. flexneri* in the United Kingdom [12]. The study was conducted in the United Kingdom, where *S. flexneri* is the most commonly isolated serotype of *S. flexneri* in the United Kingdom [12].