

Swirl String Theory (SST) Canon v0.3.1

*Omar Iskandarani**

August 26, 2025

Abstract

This Canon is the single source of truth for Swirl String Theory (SST): definitions, constants, boxed master equations, and notational conventions. It consolidates core structure up to the previous baseline (§§1–12.6) and promotes five results to canonical status: (i) Swirl Coulomb constant Λ and hydrogen soft-core, (ii) circulation–metric corollary (frame-dragging analogue), (iii) corrected swirl time-rate (Swirl Clock) law, (iv) Kelvin-compatible swirl Hamiltonian density, and (v) swirl pressure law (Euler corollary).

Versioning Semantic versions: vMAJOR.MINOR.PATCH. This file: **v0.3.1**. Every paper/derivation must state the Canon version it depends on.

Core Postulates (SST)

1. **Swirl medium:** Physics is formulated on \mathbb{R}^3 with absolute reference time. Dynamics occur in an incompressible, inviscid swirl condensate, which plays the role of a universal substrate.
2. **Strings as swirls:** Particles and excitations correspond to closed, possibly linked or knotted swirl strings with quantized circulation.
3. **String-induced gravitation:** Macroscopic attraction emerges from coherent swirl fields and swirl-pressure gradients. The effective gravitational coupling G_{swirl} is fixed by canonical constants.
4. **Swirl clocks:** Local proper-time rate depends on tangential swirl velocity. Higher swirl density slows local clocks relative to the asymptotic frame.
5. **Quantization from topology and circulation:** Discrete quantum numbers track directly to linking, writhe, twist, and circulation quantization of swirl strings.
6. **Taxonomy:** Unknotted excitations behave as bosonic string modes; chiral hyperbolic knots map to quarks; torus knots map to leptons (taxonomy documented separately).

Hydrodynamic analogy only; no mechanical “æther” is assumed in the mainstream presentation.

* Independent Researcher, Groningen, The Netherlands

Email: info@omariskandarani.com

ORCID: [0009-0006-1686-3961](https://orcid.org/0009-0006-1686-3961)

DOI: [10.5281/zenodo.16934536](https://doi.org/10.5281/zenodo.16934536)

License: CC-BY-NC 4.0 International

1 Swirl Quantization Principle

1.1 Local Circulation Quantization

The circulation of the swirl velocity field around any closed loop is quantized:

$$\Gamma = \oint \vec{v}_{\text{swirl}} \cdot d\vec{\ell} = n\kappa, \quad n \in \mathbb{Z}, \quad (1)$$

with circulation quantum

$$\kappa = \frac{h}{m_{\text{eff}}}. \quad (2)$$

This parallels the Onsager–Feynman quantization condition in superfluids, but here is elevated to a fundamental postulate of the swirl condensate.

1.2 Topological Quantization

Closed swirl filaments may form knots and links. Each topological class corresponds to a discrete excitation state:

$$\mathcal{H}_{\text{swirl}} = \{\text{trefoil, figure-eight, Hopf link, } \dots\}. \quad (3)$$

Quantum numbers such as mass, charge, and chirality are encoded in the knot invariants (linking, twist, writhe).

1.3 Unified Principle

We define Swirl Quantization as the joint discreteness of circulation and topology:

$$\text{Swirl Quantization} \equiv \left(\Gamma = n\kappa \right) \cup \left(\text{Knot spectrum } \mathcal{H}_{\text{swirl}} \right).$$

This principle underlies both the discrete particle spectrum and the emergence of fundamental interactions in Swirl String Theory.

Quantum Mechanics	Swirl String Theory
Canonical Quantization: $[x, p] = i\hbar$	Swirl Quantization Principle: $\Gamma = n\kappa, \quad n \in \mathbb{Z}$ $\mathcal{H}_{\text{swirl}} = \{\text{trefoil, figure-eight, Hopf link, } \dots\}$
Discreteness arises from operator commutators	Discreteness arises from circulation integrals and topology
Particles = eigenstates of Hamiltonian operator	Particles = knotted swirl states with quantized circulation and invariants

2 Chronos–Kelvin Invariant (Canonical)

Setting. Consider a thin, material swirl loop (nearly solid–body core) of instantaneous material radius $R(t_-)$ convected by an incompressible, inviscid medium. Let $\omega := \|\boldsymbol{\omega}\|$ denote the vorticity magnitude on the loop and r_c the canonical string radius. The local Swirl Clock is

$$S_t \equiv \frac{dt_{\text{local}}}{dt_{\infty}} = \sqrt{1 - \frac{v_t^2}{c^2}} = \sqrt{1 - \frac{\omega^2 r_c^2}{c^2}}, \quad v_t := \omega r_c. \quad (4)$$

Material derivatives are taken with respect to absolute \mathcal{A} ether-Time: $\frac{D}{Dt_-} := \frac{\partial}{\partial t_-} + \mathbf{v} \cdot \nabla$.

Theorem 2.1 (Chronos–Kelvin Invariant) *For any such loop without reconnection or source terms, Kelvin’s theorem implies the material invariant*

$$\boxed{\frac{D}{Dt_-}(R^2\omega) = 0} \iff \boxed{\frac{D}{Dt_-}\left(\frac{c}{r_c} R^2 \sqrt{1 - S_t^2}\right) = 0} \quad (5)$$

Proof (one line). Kelvin’s circulation theorem for an inviscid, barotropic medium gives $\frac{D}{Dt_-}\Gamma = 0$ with $\Gamma = \oint \mathbf{v} \cdot d\boldsymbol{\ell}$ [1, 2, 3]. For a nearly solid-body core, $\Gamma = 2\pi R v_t = 2\pi R^2\omega$; hence $\frac{D}{Dt_-}(R^2\omega) = 0$. Using (4), $R^2\omega = \frac{c}{r_c} R^2 \sqrt{1 - S_t^2}$, which yields (5).

Dimensional consistency. $[R^2\omega] = \text{m}^2\text{s}^{-1}$; and $\left[\frac{c}{r_c} R^2 \sqrt{1 - S_t^2}\right] = \text{s}^{-1} \cdot \text{m}^2 = \text{m}^2\text{s}^{-1}$.

Clock–radius transport law (corollary). From $R^2\omega = \text{const}$ and (4),

$$\frac{dS_t}{dt_-} = \frac{2(1 - S_t^2)}{S_t} \frac{1}{R} \frac{dR}{dt_-}. \quad (6)$$

Hence expansion ($dR/dt_- > 0$) pushes $S_t \rightarrow 1$ (clocks speed up), while contraction slows clocks ($S_t \downarrow$), preserving (5).

PV analogue (optional). With a uniform background rotation Ω_{bg} and column thickness H , the Ertel/PV structure gives the SST counterpart

$$\frac{D}{Dt_-} \left(\frac{\omega + \Omega_{\text{bg}}}{H} \right) = 0, \quad (7)$$

the standard potential-vorticity conservation rewritten in SST terms [4, 3].

Conditions (Canon). Incompressible, inviscid medium; barotropic swirl pressure; material loop without reconnection or external injection; absolute Æther-Time parametrization. These are the same hypotheses under which Kelvin/Helmholtz invariants hold.

Limits. Weak-swirl ($\omega r_c \ll c$): $S_t \simeq 1 - \frac{1}{2}(\omega r_c/c)^2$ and (5) reduces to the classical $R^2\omega = \text{const}$. Core on-axis limit: $v_t \rightarrow \mathbf{v}_\odot$ gives $S_t \rightarrow \sqrt{1 - (\mathbf{v}_\odot/c)^2}$, keeping (5) valid.

3 Foundational Identities

Let \mathbf{v} be the swirl velocity ($\nabla \cdot \mathbf{v} = 0$), $\boldsymbol{\omega} = \nabla \times \mathbf{v}$. For inviscid, barotropic flow [1, 2, 3, 5]:

$$\textbf{Kelvin circulation: } \frac{d\Gamma}{dt} = 0, \quad \Gamma = \oint_{\mathcal{C}(t)} \mathbf{v} \cdot d\boldsymbol{\ell}. \quad (\text{F1})$$

$$\textbf{Vorticity transport: } \frac{\partial \boldsymbol{\omega}}{\partial t} = \nabla \times (\mathbf{v} \times \boldsymbol{\omega}). \quad (\text{F2})$$

$$\textbf{Helicity: } h = \mathbf{v} \cdot \boldsymbol{\omega}, \quad H = \int h, dV \text{ (invariant up to reconnections). [6]} \quad (\text{F3})$$

These underpin knotted swirl string stability and reconnection energetics in SST.

4 Canonical Constants and Symbols

Primary SST constants (SI unless noted)

- Swirl speed scale (core): $\|\mathbf{v}_\odot\| = 1.093\,85 \times 10^6 \text{ m s}^{-1}$ (evaluate at $r = r_c$).
- String (core) radius: $r_c = 1.408\,97 \times 10^{-15} \text{ m}$.
- Effective fluid density: $\rho_f = 7.000\,00 \times 10^{-7} \text{ kg m}^{-3}$.
- Mass-equivalent density: $\rho_m = 3.893\,44 \times 10^{18} \text{ kg m}^{-3}$.
- EM-like maximal force: $F_{\text{EM}}^{\text{max}} = 2.905\,35 \times 10^1 \text{ N}$.
- Gravitational maximal force (reference scale): $F_{\text{G}}^{\text{max}} = 3.025\,63 \times 10^{43} \text{ N}$.
- Golden ratio: $\varphi = (1 + \sqrt{5})/2 \approx 1.618\,03$.

Universal constants

- $c = 299\,792\,000 \text{ m s}^{-1}$, $t_p = 5.391\,25 \times 10^{-44} \text{ s}$.
- Fine-structure constant (identified): $\alpha \approx 7.297\,35 \times 10^{-3}$.

Effective densities (mainstream field-theory style).

$\rho_f \equiv$ effective fluid density,

$$\rho_E \equiv \frac{1}{2} \rho_f \|\mathbf{v}_\odot\|^2 \quad (\text{swirl energy density}), \quad \rho_m \equiv \rho_E / c^2 \quad (\text{mass-equivalent density}).$$

Note: The local Python `constants_dict` used in simulations must mirror these values exactly; papers should quote the Canon version.

5 Canon Governance (Binding)

Definitions

Formal System. Let $\mathcal{S} = (\mathcal{P}, \mathcal{D}, \mathcal{R})$ denote the SST formal system: postulates \mathcal{P} , definitions \mathcal{D} , and admissible inference rules \mathcal{R} (variational derivation, Noether, dimensional analysis, asymptotic matching, etc.).

Canonical statement. A statement X is canonical iff X is a theorem or identity provable in \mathcal{S} :

$$\mathcal{P}, \mathcal{D} \vdash_{\mathcal{R}} X,$$

and X is consistent with all previously accepted canonical items in the current major version.

Empirical statement. A statement Y is empirical iff it asserts a measured value, fit, or protocol:

$$Y \equiv \text{“observable } \mathcal{O} \text{ has value } \hat{o} \pm \delta o \text{ under procedure II.”}$$

Empirical items calibrate symbols (e.g., v_\odot , r_c , ρ_f) but are not premises in proofs.

Status Classes

- **Axiom / Postulate (Canonical).** Primitive assumptions of SST (e.g., incompressible, inviscid medium; absolute time; Euclidean space).
- **Definition (Canonical).** Introduces symbols by construction (e.g., swirl Coulomb constant Λ by surface-pressure integral).
- **Theorem / Corollary (Canonical).** Proven consequences (e.g., Euler–SST radial balance; Swirl Clocks time-scaling).
- **Constitutive Model (Canonical if derived; otherwise Semi-empirical).** Ties fields/observables; canonical when deduced from \mathcal{P}, \mathcal{D} .
- **Calibration (Empirical).** Recommended numerical values with uncertainties for canonical symbols.
- **Research Track (Non-canonical).** Conjectures or alternatives pending proof or axiomatization.

Canonicity Tests (all required)

1. **Derivability** from \mathcal{P}, \mathcal{D} via \mathcal{R} .
2. **Dimensional Consistency** (SI throughout; correct limits).
3. **Symmetry Compliance** (Galilean + absolute time; foliation; incompressibility).
4. **Recovery Limits** (Newtonian gravity, Coulomb/Bohr, linear waves).
5. **Non-Contradiction** with accepted canonical theorems.
6. **Parameter Discipline** (no ad-hoc fits).

Examples (from current Canon)

- *Canonical (Definition):* $\Lambda \equiv \int_{S_r^2} p_{\text{swirl}} r^2 d\Omega$.
- *Canonical (Theorem):* $\frac{1}{\rho_f} \frac{dp_{\text{swirl}}}{dr} = \frac{v_\theta(r)^2}{r}$ for steady, azimuthal drift (Euler balance).
- *Empirical (Calibration):* $v_\phi = 1.09384563 \times 10^6 \text{ m s}^{-1}$ with procedure $f\Delta x$.
- *Consistency Check (Not a premise):* Hydrogen soft-core reproduces a_0, E_1 ; validates choices but remains a check, not an axiom.

6 What is Canonical in SST—and Why

[Postulate] **Incompressible, inviscid medium with absolute time and Euclidean space.** $\nabla \cdot \mathbf{v}_\phi = 0$, $\nu = 0$. This fixes the kinematic arena and legal inference rules.

[Definition] **Vorticity, circulation, helicity.** $\omega_\phi = \nabla \times \mathbf{v}_\phi$, $\Gamma = \oint \mathbf{v}_\phi \cdot d\ell$, $h = \mathbf{v}_\phi \cdot \omega_\phi$, $H = \int h dV$. Classical constructs canonized as primary SST kinematic invariants.

[Theorem] **Kelvin/vorticity transport/helicity invariants.** For inviscid, barotropic flow:

$$\frac{d\Gamma}{dt} = 0, \quad \frac{\partial \boldsymbol{\omega}_\odot}{\partial t} = \nabla \times (\mathbf{v}_\odot \times \boldsymbol{\omega}_\odot), \quad H \text{ invariant up to reconnections.}$$

[Definition] **Swirl Coulomb constant Λ .**

$$\boxed{\Lambda \equiv \int_{S_r^2} p_{\text{swirl}}(r) r^2 d\Omega} \Rightarrow [\Lambda] = \text{J m} = \text{N m}^2.$$

In SST Canon this evaluates symbolically to $\Lambda = 4\pi\rho_m v_\odot^2 r_c^4$.

[Theorem] **Hydrogen soft-core potential and Coulomb recovery.**

$$V_{\text{SST}}(r) = -\frac{\Lambda}{\sqrt{r^2 + r_c^2}} \xrightarrow{r \gg r_c} -\frac{\Lambda}{r},$$

yielding Bohr scalings $a_0 = \hbar^2/(\mu\Lambda)$, $E_n = -\mu\Lambda^2/(2\hbar^2 n^2)$.

[Theorem] **Euler–SST radial balance (swirl pressure law).** For steady, purely azimuthal drift $v_\theta(r)$,

$$0 = -\frac{1}{\rho_f} \frac{dp_{\text{swirl}}}{dr} + \frac{v_\theta(r)^2}{r} \Rightarrow \boxed{\frac{1}{\rho_f} \frac{dp_{\text{swirl}}}{dr} = \frac{v_\theta(r)^2}{r}}.$$

For flat curves $v_\theta \rightarrow v_0$: $p_{\text{swirl}}(r) = p_0 + \rho_f v_0^2 \ln(r/r_0)$.

[Definition \rightarrow Corollary] **Effective swirl line element (analogue-metric form).** In (t, r, θ, z) with azimuthal drift $v_\theta(r)$,

$$ds^2 = -(c^2 - v_\theta^2) dt^2 + 2v_\theta r d\theta dt + dr^2 + r^2 d\theta^2 + dz^2,$$

co-rotating to $ds^2 = -c^2(1 - v_\theta^2/c^2)dt^2 + \dots$, giving the Swirl Clock factor $\frac{dt_{\text{local}}}{dt_\infty} = \sqrt{1 - \frac{v_\theta^2}{c^2}}$.

[Definition] **SST Hamiltonian density (Kelvin-compatible).**

$$\mathcal{H}_{\text{SST}} = \frac{1}{2}\rho_f \|\mathbf{v}_\odot\|^2 + \frac{1}{2}\rho_f r_c^2 \|\boldsymbol{\omega}_\odot\|^2 + \lambda(\nabla \cdot \mathbf{v}_\odot).$$

Empirical Calibrations (not premises, but binding numerically)

- [Empirical] $v_\odot = 1.09384563 \times 10^6 \text{ m s}^{-1}$.
- [Empirical] $r_c = 1.40897017 \times 10^{-15} \text{ m}$.
- [Empirical] $\rho_m = 3.8934358266918687 \times 10^{18} \text{ kg m}^{-3}$.

Non-Canonical (Research Track)

Blackbody via swirl temperature, EM/SST minimal coupling, etc., remain conjectural until proven under \mathcal{S} .

Consistency & Dimension Checks (illustrative)

$$[\Lambda] = [\rho_m][v_\odot^2][r_c^4] = \frac{\text{kg}}{\text{m}^3} \cdot \frac{\text{m}^2}{\text{s}^2} \cdot \text{m}^4 = \frac{\text{kg m}^3}{\text{s}^2} = \text{J m}.$$

Soft-core Coulomb recovery: $V_{\text{SST}}(r) \rightarrow -\Lambda/r$ as $r/r_c \rightarrow \infty$.

7 Canonical Coarse-Graining of ρ_f from a Swirl-String Bath

Scope. The medium is modeled as an incompressible, inviscid fluid populated by thin swirl strings. We derive the bulk effective fluid density ρ_f via coarse-graining of line-supported mass and vorticity, relying on Euler kinematics and Kelvin-Helmholtz invariants.

7.1 Axioms and Definitions

A representative string carries:

$$(D1) \quad \mu_* \equiv \rho_m \pi r_c^2 \quad [\text{kg/m}], \quad (8)$$

$$(D2) \quad \Gamma_* \equiv \oint \mathbf{v}_\zeta \cdot d\boldsymbol{\ell} \simeq \kappa_\Gamma r_c v_\zeta, \quad \kappa_\Gamma = 2\pi \text{ (near-solid-body core)}. \quad (9)$$

Let $\nu \equiv N_{\text{str}}/A$ [m^{-2}] be the areal string density. Then:

$$(C1) \quad \rho_f = \mu_* \nu, \quad (10)$$

$$(C2) \quad \langle \boldsymbol{\omega}_\zeta \rangle = \Gamma_* \nu \hat{\mathbf{t}}_{\text{avg}} \Rightarrow |\langle \boldsymbol{\omega}_s \rangle| = \Gamma_* \nu. \quad (11)$$

7.2 First-Principles Derivation

Combining (C1)–(C2):

$$\boxed{\rho_f = \mu_* \frac{\langle \boldsymbol{\omega}_s \rangle}{\Gamma_*} = \frac{\rho_m \pi r_c^2}{\kappa_\Gamma r_c v_\zeta} \langle \boldsymbol{\omega}_s \rangle = \frac{\rho_m r_c}{2 v_\zeta} \langle \boldsymbol{\omega}_s \rangle} \quad (\kappa_\Gamma = 2\pi). \quad (12)$$

For uniform solid-body rotation Ω , $\langle \boldsymbol{\omega}_s \rangle = 2\Omega$,

$$\boxed{\rho_f = \frac{\rho_m r_c}{v_\zeta} \Omega} \quad [\text{kg/m}^3]. \quad (13)$$

Energy and tension scales.

$$\boxed{u_{\text{swirl}} = \frac{1}{2} \rho_f v_\zeta^2}, \quad \boxed{T_* = \frac{1}{2} \mu_* v_\zeta^2}.$$

7.3 Numerical Calibration (SST Canonical Constants)

With $\rho_m = 3.8934358266918687 \times 10^{18} \text{ kg/m}^3$, $r_c = 1.40897017 \times 10^{-15} \text{ m}$, $v_\zeta = 1.09384563 \times 10^6 \text{ m/s}$, one finds

$$\Gamma_* = 2\pi r_c v_\zeta = 9.68361920 \times 10^{-9} \text{ m}^2/\text{s}, \quad T_* = 1.45267535 \times 10^1 \text{ N}.$$

From (13),

$$\rho_f = (5.01509060 \times 10^{-3}) \Omega,$$

so the Canon baseline $\rho_f = 7.0 \times 10^{-7} \text{ kg/m}^3$ occurs at

$$\boxed{\Omega_* = 1.39578735 \times 10^{-4} \text{ s}^{-1} \text{ (period } \approx 12.5 \text{ h)}}.$$

8 Master Equations (Boxed, Definitive)

8.1 Master Energy and Mass Formula (SST)

$$\boxed{E_{\text{SST}}(V) = \frac{4}{\alpha \varphi} \left(\frac{1}{2} \rho_f v_{\odot}^2 \right) V} \quad [\text{J}], \quad \boxed{M_{\text{SST}}(V) = \frac{E_{\text{SST}}(V)}{c^2}} \quad [\text{kg}].$$

Numerics per unit volume: $\frac{1}{2} \rho_f v_{\odot}^2 \approx 4.1877439 \times 10^5 \text{ J m}^{-3}$, $\frac{4}{\alpha \varphi} \approx 3.3877162 \times 10^2$, $\Rightarrow E/V \approx 1.418688 \times 10^8 \text{ J m}^{-3}$, $M/V \approx 1.57850 \times 10^{-9} \text{ kg m}^{-3}$.

8.2 Swirl–Gravity Coupling

$$\boxed{G_{\text{string}} = \frac{v_{\odot} c^5 t_p^2}{2 F_{\text{EM}}^{\text{max}} r_c^2}}$$

Numerically $\approx 6.674302 \times 10^{-11} \text{ m}^3 \text{ kg}^{-1} \text{ s}^{-2}$ with the Canon constants.

8.3 Topology–Driven Mass Law (invariant form)

Let $T(p, q)$ be a torus knot/link, $n = \text{gcd}(p, q)$ components, braid index $b(T) = \min(|p|, |q|)$, Seifert genus $g(T)$ (with standard link adjustment). Using ropelength $\mathcal{L}_{\text{tot}}(T)$ and string core radius r_c :

$$\boxed{M(T(p, q)) = \left(\frac{4}{\alpha} \right) b(T)^{-3/2} \varphi^{-g(T)} n^{-1/\varphi} \left(\frac{1}{2} \rho_f v_{\odot}^2 \right) \frac{\pi r_c^3 \mathcal{L}_{\text{tot}}(T)}{c^2}}.$$

Dimensionality follows from the factor $\frac{1}{2} \rho_f v_{\odot}^2$ (J m^{-3}) times a volume.

8.4 Swirl Clocks (Local Time-Rate)

$$\boxed{\frac{dt_{\text{local}}}{dt_{\infty}} = \sqrt{1 - \frac{\|\boldsymbol{\omega}_{\odot}\|^2 r_c^2}{c^2}} = \sqrt{1 - \frac{\|\mathbf{v}_{\odot}\|^2}{c^2}} \quad (r = r_c)}.$$

Historical (deprecated) variant without a length scale is retained only for traceability.

8.5 Swirl Angular Frequency Profile

$$\boxed{\Omega_{\text{swirl}}(r) = \frac{v_{\odot}}{r_c} e^{-r/r_c}}, \quad \Omega_{\text{swirl}}(0) = \frac{v_{\odot}}{r_c}.$$

8.6 Vorticity Potential (Canonical Form)

$$\Phi(\vec{r}, \boldsymbol{\omega}_{\odot}) = \frac{v_{\odot}^2}{2 F_{\text{EM}}^{\text{max}}} \boldsymbol{\omega}_{\odot} \cdot \vec{r}.$$

Dimensional remark: Use with the SST Lagrangian ensuring $\rho_f \Phi$ has energy density units.

9 Unified SST Lagrangian (Definitive Form)

Let \mathbf{v}_{\odot} be the velocity, ρ_f constant (incompressible), $\boldsymbol{\omega}_{\odot} = \nabla \times \mathbf{v}_{\odot}$, and λ enforce incompressibility.

$$\boxed{\mathcal{L}_{\text{SST}} = \frac{1}{2} \rho_f \|\mathbf{v}_{\odot}\|^2 - \rho_f \Phi(\vec{r}, \boldsymbol{\omega}_{\odot}) + \lambda (\nabla \cdot \mathbf{v}_{\odot}) + \eta \int (\mathbf{v}_{\odot} \cdot \boldsymbol{\omega}_{\odot}) dV + \mathcal{L}_{\text{couple}}[\Gamma, \mathcal{K}]}.$$

Here $\mathcal{L}_{\text{couple}}$ encodes coupling to quantized circulation Γ and knot invariants \mathcal{K} (linking, writhe, twist).

10 Notation, Ontology, and Glossary

- **Absolute time (A-time):** global time parameter of the medium.
- **Chronos Time (C-time):** asymptotic observer time (dt_∞).
- **Swirl Clocks:** local clocks set by $\|\omega_\odot\|$ or $\|\mathbf{v}_\odot\|$ per Sec. 9.
- **String taxonomy:** leptons = torus knots; quarks = chiral hyperbolic knots; bosons = unknots; neutrinos = linked knots.
- **Chirality:** ccw \leftrightarrow matter; cw \leftrightarrow antimatter via swirl-gravity coupling.

11 Canonical Checks (What to Verify in Every Paper)

1. Dimensional analysis on every new term/equation.
2. Limits: low-swirl $\|\omega_\odot\| \rightarrow 0$ recovers classical mechanics/EM; large-scale averages reproduce Newtonian gravity with G_{string} .
3. Numerics: provide prefactors using Canon constants; add any new constants to Sec. 4.
4. Explicit topology \leftrightarrow quantum mapping (which invariants, normalization).
5. Cite any non-original constructs (BibTeX keys).

12 Persona Prompts

Reviewer Persona

You are a peer reviewer for an SST paper. Use only the definitions and constants in the "SST Canon (v0.3.1)". Check dimensional consistency, limiting behavior, and numerical validation. Flag any use of non-canonical constants or equations unless equivalence is proved. Demand explicit mapping from knot invariants (linking, writhe, twist) to claimed quantum numbers.

Theorist Persona

You are a theoretical physicist specialized in Swirl String Theory (SST). Base all reasoning on the attached "SST Canon (v0.3.1)". Your task: derive the swirl-based Hamiltonian for [TARGET SYSTEM], use Sec. 9, and verify the Swirl Clock law (Sec. 8.4). Provide boxed equations, dimensional checks, and a short numerical evaluation using the Canon constants.

Bridging Persona (Compare to GR/SM)

Work strictly within SST Canon (v0.3.1). Compare [TARGET] to its GR/SM counterpart. Identify exact replacements (e.g., curvature \rightarrow swirl), and show which terms reduce to Newtonian/Maxwellian limits. Include a correspondence table and any constraints needed for equivalence.

13 Session Kickoff Checklist

1. Start new chat per task; attach this Canon first.
2. Paste a persona prompt (Sec. 12).
3. Attach only task-relevant papers/sources.
4. State any corrections explicitly (they persist in the session).
5. At end, record Canon deltas (if any) and bump version.

14 Appendix: Boxed Canon Equations (paste-ready)

1. **Energy:** $E_{\text{SST}} = \frac{4}{\alpha\varphi} \left(\frac{1}{2} \rho_f v_{\odot}^2 \right) V$

2. **Mass:** $M_{\text{SST}} = \frac{E_{\text{SST}}}{c^2}$

3. **G coupling:** $G_{\text{string}} = \frac{v_{\odot} c^5 t_p^2}{2 F_{\text{EM}}^{\text{max}} r_c^2}$

4. **Swirl Clock:** $\frac{dt_{\text{local}}}{dt_{\infty}} = \sqrt{1 - \|\boldsymbol{\omega}_{\odot}\|^2 r_c^2 / c^2} = \sqrt{1 - \|\mathbf{v}_{\odot}\|^2 / c^2}$

5. **Swirl profile:** $\Omega_{\text{swirl}}(r) = \frac{v_{\odot}}{r_c} e^{-r/r_c}$

15 Swirl Hamiltonian Density (Canonical Form)

With the effective fluid density ρ_f , the swirl vorticity $\boldsymbol{\omega}_{\odot} = \nabla \times \mathbf{v}_{\odot}$, and a Lagrange multiplier λ to enforce incompressibility, a Kelvin-compatible, dimensionally normalized Hamiltonian density is given by:

$$\mathcal{H}_{\text{SST}}[\mathbf{v}_{\odot}] = \frac{1}{2} \rho_f \|\mathbf{v}_{\odot}\|^2 + \frac{1}{2} \rho_f r_c^2 \|\boldsymbol{\omega}_{\odot}\|^2 + \frac{1}{2} \rho_f r_c^4 \|\nabla \boldsymbol{\omega}_{\odot}\|^2 + \lambda (\nabla \cdot \mathbf{v}_{\odot}). \quad (14)$$

All terms carry units of energy density ($\text{J} \cdot \text{m}^{-3}$). The first two terms represent the kinetic and rotational energy of the swirl, while the third term, proportional to the gradient of vorticity, corresponds to the energy associated with the curvature or "stiffness" of the swirl filaments. In the limit where the core radius $r_c \rightarrow 0$ or for spatially uniform vorticity, this expression reduces to the simpler forms.

16 Swirl Pressure Law (Euler Corollary)

For a steady, purely azimuthal drift velocity $v_{\theta}(r)$ with no radial flow, the radial component of the Euler momentum equation for an inviscid fluid provides a direct relationship for the swirl pressure gradient:

$$\frac{1}{\rho_f} \frac{dp_{\text{swirl}}}{dr} = \frac{v_{\theta}(r)^2}{r}. \quad (15)$$

This is a canonical theorem derived directly from first principles. For a system exhibiting an asymptotically flat rotation curve where $v_{\theta}(r) \rightarrow v_0$ for large r , the pressure profile is found by integration:

$$p_{\text{swirl}}(r) = p_0 + \rho_f v_0^2 \ln \left(\frac{r}{r_0} \right). \quad (16)$$

Here, p_0 is the pressure at a reference radius r_0 . The resulting outward-rising pressure creates an inward-pointing force ($-\nabla p_{\text{swirl}}$), providing the centripetal acceleration required to maintain the flat rotation curve.

17 Experimental Protocols (Canon-ready)

Universality of $v_{\odot} = f \Delta x$ (metrology across platforms)

From `ExperimentalValidationOfVortexCoreTangentialVelocity.tex`: measure a natural frequency f and a spatial step Δx from standing/propagating modes; verify

$$v_{\odot} = f \Delta x \approx 1.09385 \times 10^6 \text{ m/s} \quad (\text{X1})$$

Platforms: magnet/electret domains, laser interferometry on coil-bound modes, and acoustic analogues. Require ppm-level agreement; report mean and standard deviation across platforms.

Swirl gravitational potential

From `ExperimentalValidationOfGravitationalPotential.tex`: infer $p_{\text{swirl}}(r)$ from centripetal balance (§16) and compare predicted forces with measured thrust or buoyancy anomalies in shielded high-voltage/coil experiments (geometry: starship/Rodin coils). Ensure dimensional consistency and calibrate only via Canon constants.

References

- [1] H. von Helmholtz. über integrale der hydrodynamischen gleichungen, welche den wirbelbewegungen entsprechen. Journal für die reine und angewandte Mathematik, 55:25–55, 1858.
- [2] W. Thomson (Lord Kelvin). On vortex motion. Transactions of the Royal Society of Edinburgh, 25:217–260, 1869.
- [3] G. K. Batchelor. An Introduction to Fluid Dynamics. Cambridge University Press, 1967.
- [4] Hans Ertel. Ein neuer hydrodynamischer erhaltungssatz. Meteorologische Zeitschrift, 59:271–281, 1942.
- [5] L. D. Landau and E. M. Lifshitz. Fluid Mechanics. Course of Theoretical Physics, Vol. 6. Pergamon, 2nd edition, 1987.
- [6] H. K. Moffatt. The degree of knottedness of tangled vortex lines. Journal of Fluid Mechanics, 1969.
- [7] Philip G. Saffman. Vortex Dynamics. Cambridge University Press, 1992.
- [8] R. P. Feynman. Application of quantum mechanics to liquid helium. In C. J. Gorter, editor, Progress in Low Temperature Physics, volume 1, pages 17–53. North-Holland, 1955.
- [9] Russell J. Donnelly. Quantized Vortices in Helium II. Cambridge University Press, 1991.
- [10] L. D. Landau and E. M. Lifshitz. Fluid Mechanics. Course of Theoretical Physics, Vol. 6. Pergamon, 2nd edition, 1987.
- [11] John David Jackson. Classical Electrodynamics. Wiley, 3rd edition, 1999.
- [12] W. Thomson (Lord Kelvin). On vortex atoms. Proc. Royal Society of Edinburgh, 1867.
- [13] E. Schrödinger. An undulatory theory of the mechanics of atoms and molecules. Physical Review, 1926.

- [14] G. Călugăreanu. L'intégral de gauss et l'analyse des noeuds tridimensionnels. Rev. Math. Pures Appl., 4, 1959.
- [15] J. H. White. Self-linking and the gauss integral in higher dimensions. American Journal of Mathematics, 91, 1969.
- [16] F. B. Fuller. The writhing number of a space curve. Proc. Natl. Acad. Sci. USA, 68:815–819, 1971.
- [17] H. K. Moffatt and R. L. Ricca. Helicity and the călugăreanu invariant. Proc. R. Soc. Lond. A, 439:411–429, 1992.
- [18] H. Hopf. über die abbildungen der dreidimensionalen sphäre auf die kugelfläche. Mathematische Annalen, 104:637–665, 1931.
- [19] J. H. C. Whitehead. An expression of hopf's invariant as an integral. Proc. Natl. Acad. Sci. USA, 33:117–123, 1947.
- [20] I. E. Dzyaloshinskii. A thermodynamic theory of “weak” ferromagnetism of antiferromagnetics. J. Phys. Chem. Solids, 4:241–255, 1958.
- [21] T. Moriya. Anisotropic superexchange interaction and weak ferromagnetism. Phys. Rev., 120:91–98, 1960.
- [22] A. Aharoni. Introduction to the Theory of Ferromagnetism. Oxford University Press, 1996.
- [23] G. K. Batchelor. An Introduction to Fluid Dynamics. Cambridge University Press, 1967.
- [24] W. G. Unruh. Experimental black-hole evaporation? Physical Review Letters, 46:1351–1353, 1981.
- [25] M. Visser. Acoustic black holes: horizons, ergospheres and hawking radiation. Classical and Quantum Gravity, 15(6):1767–1791, 1998.
- [26] P. Painlevé. La mécanique classique et la théorie de la relativité. Comptes Rendus de l'Académie des Sciences, 173:677–680, 1921.
- [27] A. Gullstrand. Allgemeine lösung des statischen einkörperproblems in der einsteinschen gravitationstheorie. Arkiv för Matematik, Astronomi och Fysik, 16:1–15, 1922.
- [28] Dale Rolfsen. Knots and Links. Publish or Perish, 1976.
- [29] W. B. R. Lickorish. An Introduction to Knot Theory, volume 175 of Graduate Texts in Mathematics. Springer, 1997.
- [30] Kunio Murasugi. Knot Theory and Its Applications. Birkhäuser, 1996.
- [31] Jason Cantarella, Robert B. Kusner, and John M. Sullivan. On the minimum ropelength of knots and links. Inventiones mathematicae, 150(2):257–286, 2002.
- [32] Eric J. Rawdon. Approximating the thickness of a knot. In A. Stasiak, V. Katrich, and L. Kauffman, editors, Ideal Knots, pages 143–150. World Scientific, 1998.
- [33] Roy P. Kerr. Gravitational field of a spinning mass as an example of algebraically special metrics. Physical Review Letters, 11:237–238, 1963.
- [34] G. Călugăreanu. Sur les classes d'isotopie des noeuds tridimensionnels et leurs invariants. Czechoslovak Mathematical Journal, 1961.

- [35] W. Wien. Ueber die energieverteilung im emissionsspectrum eines schwarzen körpers. Annalen der Physik, 1894.
- [36] M. Planck. On the law of distribution of energy in the normal spectrum. Annalen der Physik, 1901.