




### • 一维高斯分布

$$N(x \mid \mu, \sigma) = rac{1}{\sigma \sqrt{2\pi}} e^{-rac{(x-\mu)^2}{2\sigma^2}}$$

## 聚类: k-means、GMM



#### • 一维高斯分布



$$N(x\mid \mu,\sigma) = rac{1}{\sigma\sqrt{2\pi}}e^{-rac{(x-\mu)^2}{2\sigma^2}}$$

### 聚类: k-means、GMM



- 二维高斯分布
  - D表示d维

$$N(x\mid \mu, \Sigma) = rac{1}{(2\pi)^{d/2}\sqrt{|\Sigma|}}exp(-rac{1}{2}(x-\mu)^T\Sigma^{-1}(x-\mu))$$


$$\Sigma$$
即 $\begin{pmatrix} \sigma_1^2 & \sigma_{12} \\ \sigma_{21} & \sigma_2^2 \end{pmatrix}$   
 $\Sigma^{-1}$ 为矩阵的逆矩阵  
 $|\Sigma|$ 为矩阵的行列式

# 聚类: k-means、GMM



### • 二维高斯分布

$$N(x\mid \mu, \Sigma) = rac{1}{(2\pi)^{d/2}\sqrt{|\Sigma|}}exp(-rac{1}{2}(x-\mu)^T\Sigma^{-1}(x-\mu))$$



$$N\left[egin{pmatrix} \mu_1 \ \mu_2 \end{pmatrix}, egin{pmatrix} \sigma_1^2 & \sigma_{12} \ \sigma_{21} & \sigma_2^2 \end{pmatrix}
ight]$$