

REVISED How blockchain-timestamped protocols could improve the trustworthiness of medical science [version 32; referees: 3 approved]

Greg Irving¹, John Holden²

Author affiliations

Grant information



This article is included in the [All trials matter](#) channel.

Abstract

Trust in scientific research is diminished by evidence that data are being manipulated. Outcome switching, data dredging and selective publication are some of the problems that undermine the integrity of published research. Methods for using blockchain to provide proof of pre-specified endpoints in clinical trial protocols were first reported by Carlisle. We wished to empirically test such an approach using a clinical trial protocol where outcome switching has previously been reported. Here we confirm the use of blockchain as a low cost, independently verifiable method to audit and confirm the reliability of scientific studies.

Corresponding author: Greg Irving

How to cite: Irving G and Holden J. How blockchain-timestamped protocols could improve the trustworthiness of medical science [version 32; referees: 3 approved]. *F1000Research* 20172016, 5:222

(doi: [10.12688/f1000research.8114.32](https://doi.org/10.12688/f1000research.8114.32)) Copyright: © 20172016 Irving G and Holden J. This is an open access article distributed under the terms of the [Creative Commons Attribution Licence](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. Data associated with the article are available under the terms of the [Creative Commons Zero "No rights reserved" data waiver](#) (CC0 1.0 Public domain dedication).

Competing interests:

No competing interests were disclosed.

First published: 26 Feb 2016, 5:222 (doi: [10.12688/f1000research.8114.1](https://doi.org/10.12688/f1000research.8114.1)) Latest published: 30 Mar

2017, 5:222 (doi: [10.12688/f1000research.8114.3](https://doi.org/10.12688/f1000research.8114.3))

REVISED Amendments from Version 21

To clarify our approach to verifying the original transaction we have included text to confirm that this was done using the same Strongcoin© account used to generate the original transaction. We have also replaced the .docx copy of the file used to verify the transaction with the original .txt file.

~~The method we tested here was first proposed by Carlisle in the grey literature. Clear reference to the previously described method (Reference 6) has been added throughout the revised article.~~

See referee responses

Editorial note:

Concerns have been raised about the overlap between Version 1 of this article and a previously published blog by Carlisle, who proposed the method 2 years earlier [Carlisle, Benjamin Gregory. “Proof of prespecified endpoints in medical research with the bitcoin blockchain”, 25 August 2014], and that the correction (Version 2) published soon after the original was not sufficient to rectify the overlap.

The case has since been discussed in a Committee of Publication Ethics (COPE) Forum, and COPE advised that the correction was sufficient to correct the scientific literature.

The case has been referred to the University of Cambridge for consideration.

Introduction

Trust in scientific research is diminished by evidence that data are being manipulated¹. Outcome switching, data dredging and selective publication are some of the problems that undermine the integrity of published research. The declaration of Helsinki states that every clinical trial must be registered in a publicly accessible database before recruitment of the first subject². Yet despite the creation of numerous trial registries problems such as differences between pre-specified and reported outcomes persist³⁻⁵. If readers doubt the trustworthiness of scientific research then it is largely valueless to them and those they influence. Here we confirm the use of blockchain as a low cost, independently verifiable method that could be widely and readily used to audit and confirm the reliability of scientific studies.

A blockchain is a distributed, tamper proof public ledger of timestamped transactions. It provides a method for establishing the existence of a transaction at a particular time that can be independently verified by any interested party. When someone wishes to add to it, participants in the network – all of whom have copies of the existing blockchain – run algorithms to evaluate and verify the proposed action. Once the majority of ‘nodes’ confirm that a transaction is valid i.e. matches the blockchain history then the new transaction will be approved and added to the chain. Once a block of data is recorded on a blockchain ledger it is extremely difficult to change or remove it as doing so would require changing the record on many thousands of computers worldwide. This prevents tampering or future revision of a submitted timestamped record. Such distributive version control has been increasingly used in fields such as software development, engineering and genetics. A method for using blockchain to provide proof of pre-specified endpoints in clinical trial protocols was first suggested by Carlisle in 2014⁶. We wished to empirically test such an approach using a clinical trial protocol where outcome switching has previously been reported.

Methods

In this study we used publically available documentation from a recently reported randomized control trial^{7,8}. A copy of the clinicaltrials.gov study protocol was prepared based on its pre-specified endpoints and planned analyses which was saved as an unformatted text file⁷. Following a method similar to that described by Carlisle the document’s SHA256 digest for the text was then calculated by entering text from the trial protocol into an SHA256 calculator (Xorbin©)⁶. This was then converted into a bitcoin private key and corresponding public key using a bitcoin wallet. To do this a new account was created in Strongcoin©⁹ and the SHA256 digest used as the account password [to generate a private key](#)(private key)⁶. From this

Strongcoin© automatically generated a corresponding Advanced Encryption Standard 256 bit public key⁶. An arbitrary amount of bitcoin was then sent to a corresponding bitcoin address. To verify the existence of the document a second researcher was sent the originally prepared unformatted document. An SHA256 digest was created as previously described. ~~The and a~~ corresponding private key, public key and bitcoin address were confirmed using the original Strongcoin© account and blockchain.info©⁶. ~~The generated~~. ~~The exact replication of the~~ bitcoin address (1AHjCz2oEUTH8js4S8vViC8NKph4zCACXH) was then used to prove the documents existence in the blockchain using blockchain.info©¹⁰. The protocol document was then edited to reflect any changes to pre-specified outcomes as reported by the COMPare group³. This was used to create a further SHA256 digest ~~which differed to that for the pre-specified protocol and would not allow the and corresponding public,~~ private key ~~to be unlocked in Strongcoin©3 and bitcoin address3~~.

Dataset 1. Unformatted text file.

Downloaded data do not display as expected?

Download the data

Results

Incorporating a transaction from the bitcoin wallet into the blockchain using a private key generated from the SHA256 digest of the trial protocol timestamped a record of the study protocol. The transaction took under five minutes to complete. The process cost was free as the nominal bitcoin transaction could be retrieved. Researchers were able to search for the transaction on the blockchain, confirm the date when the transaction occurred and verify the authenticity of the original protocol ~~by generating identical public and private keys~~. Any changes made to the original document generated ~~a different SHA 256 digest~~ ~~different public and private keys~~ indicating that protocol had been altered. This included assessment of an edited protocol reflecting pre-specified outcomes not reported or non-pre-specified outcomes reported in the final paper.

Discussion

Fraud in scientific methods erodes confidence in medicine as a whole which is essential to performing its function¹. This study demonstrates that the method described by Carlisle provides an immutable record of the existence, integrity and ownership of a specific trial protocol⁶. It is a simple and cheap way of allowing a third party to audit and externally validate outcomes and analyses specified *a-priori* with the findings reported *a-posteriori*. It prevents researchers from changing study endpoints or analyses after seeing their study results without reporting such changes⁶. Transaction codes could be recorded in scientific papers, reference databases or trial registries to facilitate external verification. As discussed in the CONSORT guidelines, switching of outcomes in trials is sometimes necessary for perfectly legitimate reasons but this should be disclosed in the final report¹¹. The use of blockchain timestamped protocols could facilitate trust in the reporting of this process by providing evidence of precisely when protocol changes took place. At the same time, fraudulent attempts to prepare multiple study protocols in advance would be technically possible but would also leave behind a publically available trail of evidence that could not be destroyed⁶. The blockchain offers a number of advantages over the current approaches used trial registries or publishing protocols. Firstly, the blockchain would not be confined to the validation of clinical trials. The approach could be used for a whole range of observational and experimental studies where registries do not currently exist. Secondly, the blockchain provides a real-time timestamped record of a protocol. Such precision is important given persistent problems with protocol registration after trial initiation¹². Thirdly, with over 30,000 trials currently published annually and rising, manual outcome verification is simply not possible¹³.

Conclusion

Blockchain-timestamped protocols can allow the exact wording and existence of a protocol at a given point in time to be verified. They have the potential to support automated, extremely robust verification of pre-specified with reported outcomes. This evidence should increase trust in medical science by diminishing suspicion in reported data and the conclusions that are drawn.