

ECE 457A TUTORIAL 08: GAME THEORY

06-Nov-2023

Danial Sadrian Zadeh,
Department of Electrical and Computer Engineering

Mixed Strategies (Example 1)

There is no NE in this game.

If we play this game, we should be unpredictable.

\implies we must randomize (mix) between strategies.

First-Order Condition

P2

	H α	T $(1-\alpha)$
H θ	1, -1	-1, 1
T $(1-\theta)$	-1, 1	1, -1

P1

$$\pi_{P1} = (\theta) \underbrace{((1)(\alpha) + (-1)(1-\alpha))}_{2\alpha-1} + (1-\theta) \underbrace{((-1)(\alpha) + (1)(1-\alpha))}_{-2\alpha+1} = 4\alpha\theta - 2\theta - 2\alpha + 1$$

$$\pi_{P2} = (\alpha) \underbrace{((-1)(\theta) + (1)(1-\theta))}_{-2\theta+1} + (1-\alpha) \underbrace{((1)(\theta) + (-1)(1-\theta))}_{2\theta-1} = -4\alpha\theta + 2\theta + 2\alpha - 1$$

$$\frac{\partial \pi_{p1}}{\partial \alpha} = 4\alpha - 2 = 0 \Rightarrow \boxed{\alpha = 0.5}$$

$$\frac{\partial \pi_{p2}}{\partial \alpha} = -4\theta + 2 = 0 \Rightarrow \boxed{\theta = 0.5}$$

$$\left. \begin{aligned} \frac{\partial^2 \pi_{p1}}{\partial \alpha^2} &= 0 \leq 0 \quad \checkmark \\ \frac{\partial^2 \pi_{p2}}{\partial \alpha^2} &= 0 \leq 0 \quad \checkmark \end{aligned} \right\}$$

Payoff-Equating Method

$$P1: 2\alpha - 1 = -2\alpha + 1 \Rightarrow \boxed{\alpha = 0.5}$$

$$P2: -2\theta + 1 = 2\theta - 1 \Rightarrow \boxed{\theta = 0.5}$$

Mixed Strategies (Example 2)

we need to get rid of as many strategies as we can.

For P1, M is completely dominated by U!!!

For P2, m is dominated by ℓ !!!

FOC

	ℓ	m	r
U	3, 2	2, 1	1, 3
M	2, 1	1, 5	0, 3
D	1, 3	4, 2	2, 2

	α	$(1-\alpha)$
θ	3, 2	1, 3
$(1-\theta)$	1, 3	2, 2

$$\pi_{p1} = \underbrace{\Theta((3)(\alpha) + (1)(1-\alpha))}_{2\alpha+1} + (1-\Theta) \underbrace{((1)(\alpha) + (2)(1-\alpha))}_{-\alpha+2} = 3\alpha\Theta - \alpha - \Theta + 2$$

$$\pi_{p2} = \alpha \underbrace{((2)(\Theta) + (3)(1-\Theta))}_{-\Theta+3} + (1-\alpha) \underbrace{((3)(\Theta) + (2)(1-\Theta))}_{\Theta+2} = -2\alpha\Theta + \alpha + \Theta + 2$$

$$\frac{\partial \pi_{p1}}{\partial \Theta} = 3\alpha - 1 = 0 \Rightarrow \boxed{\alpha = \frac{1}{3}} \quad \left\{ \begin{array}{l} \frac{\partial \pi_{p2}}{\partial \alpha} = -2\Theta + 1 = 0 \Rightarrow \boxed{\Theta = \frac{1}{2}} \\ \frac{\partial^2 \pi_{p1}}{\partial \Theta^2} = 0 \leq 0 \checkmark \\ \frac{\partial^2 \pi_{p2}}{\partial \alpha^2} = 0 \leq 0 \checkmark \end{array} \right.$$

$$\boxed{PEM} \quad \left\{ \begin{array}{l} P1: (2\alpha+1) = (-\alpha+2) \Rightarrow \boxed{\alpha = \frac{1}{3}} \checkmark \\ P2: (-\Theta+3) = (\Theta+2) \Rightarrow \boxed{\Theta = \frac{1}{2}} \checkmark \end{array} \right.$$

Continuous Strategies (Cournot and Stackelberg)

profit per quantity: $P(Q) = 9 - q_1 - q_2$

total output: $Q = q_1 + q_2$

marginal cost: $c(q) = 2q \longrightarrow \underline{c = 2}$

payoffs:
$$\begin{cases} \pi_1 = P(Q)q_1 - c(q_1) = (9 - q_1 - q_2)q_1 - 2q_1 \\ \pi_2 = P(Q)q_2 - c(q_2) = (9 - q_1 - q_2)q_2 - 2q_2 \end{cases}$$

Cournot

Reaction Functions

$$R_1 = \frac{\partial \pi_1}{\partial q_1} = 7 - 2q_1 - q_2 = 0$$

$$R_2 = \frac{\partial \pi_2}{\partial q_2} = 7 - q_1 - 2q_2 = 0$$

Cournot-Nash
Eq.



$$R_1 = R_2 \Rightarrow$$

$$q_1 = q_2$$

$$R_1 = 7 - 2q_1 - q_1 = 0 \Rightarrow q_1 = \frac{7}{3} = q_2 \quad \checkmark$$

$$\text{total market output} = Q = q_1 + q_2 = \frac{14}{3}$$

Stackelberg

Let us assume that player 2 moves first. So, player 2 is the Stackelberg leader, while player 1 is the Stackelberg follower.

Therefore, player 2 predicts the reaction function of player 1.

$$R_1 = \frac{\partial \pi_1}{\partial q_1} = 7 - 2q_1 - q_2 = 0 \Rightarrow q_1 = \frac{7 - q_2}{2} \quad (*)$$

$$\pi_2 = \left(9 + \frac{q_2 - 7}{2} - q_2\right)q_2 - 2q_2$$

$$\Rightarrow \frac{\partial \pi_2}{\partial q_2} = 7 - \frac{7}{2} - q_2 = 0 \Rightarrow q_2 = \frac{7}{2} \quad (*) \Rightarrow q_1 = \frac{7}{4}$$

substitution

$$Q = q_1 + q_2 = \frac{7}{4} + \frac{7}{2} = \frac{21}{4}$$

$$\left\{ \begin{array}{l} P1: \text{follower} \\ P2: \text{leader} \end{array} \right. \Rightarrow \left\{ \begin{array}{l} q_1 = \frac{7}{4} \\ q_2 = \frac{7}{2} \end{array} \right. \Rightarrow Q = \frac{21}{4}$$

$$\left\{ \begin{array}{l} P1: \text{leader} \\ P2: \text{follower} \end{array} \right. \Rightarrow \left\{ \begin{array}{l} q_1 = \frac{7}{2} \\ q_2 = \frac{7}{4} \end{array} \right. \Rightarrow Q = \frac{21}{4}$$

Cournot

$$q_1 = q_2 = \frac{7}{3} = 2.37$$

$$Q = q_1 + q_2 = \frac{14}{3} = 4.67$$

$$P(Q) = 4.33$$

Stackelberg

$$q_1 = \frac{7}{4} = 1.75$$

$$q_2 = \frac{7}{2} = 3.5$$

$$Q = q_1 + q_2 = \frac{21}{4} = 5.25$$

most total
output

$$P(Q) = 3.75$$

lowest price

Consumers' preference

Continuous Strategies (Bertrand)

Two firms setting price at the same time

$$\left\{ \begin{array}{l} q_1 = 72 - 3P_1 + 2P_2 \\ q_2 = 72 + 2P_1 - 3P_2 \end{array} \right. \quad \left\{ \begin{array}{l} \text{marginal cost} \\ c = 0 \text{ for both} \end{array} \right.$$

Payoffs

$$\left\{ \begin{array}{l} \pi_1 = q_1 (P_1 - c) = 72P_1 - 3P_1^2 + 2P_1P_2 \\ \pi_2 = q_2 (P_2 - c) = 72P_2 + 2P_1P_2 - 3P_2^2 \end{array} \right.$$

$$\left. \begin{aligned} R_1 &= \frac{\partial \pi_1}{\partial p_1} = 72 - 6p_1 + 2p_2 = 0 \\ R_2 &= \frac{\partial \pi_2}{\partial p_2} = 72 + 2p_1 - 6p_2 = 0 \end{aligned} \right\} \Rightarrow R_1 = R_2$$

$$\Rightarrow -6p_1 + 2p_2 = 2p_1 - 6p_2 \Rightarrow \boxed{p_1 = p_2}$$

$$\textcircled{K} \quad 72 - 6p_1 + 2p_1 = 0 \Rightarrow \boxed{p_1 = 18 = p_2} \checkmark$$

$$q_1 = q_2 = 54 \Rightarrow Q = q_1 + q_2 = \boxed{108} \checkmark$$

References

- <https://www.tayfunsonmez.net/wp-content/uploads/2013/10/E3o8SL7.pdf>
- https://eml.berkeley.edu/~webfac/dellavigna/e101a_sp08/fexam3solutions.pdf

