

Estimating the number of COVID-19 victims by using a Monte Carlo algorithm on a generalized logistic equation

Biagio Paparella

10.06.2020

Oberseminar - Institute for Numerical Simulation - Bonn

Goal of my recent work: influenced by recent events, try predicting the number of COVID-19 victims.

- (Possibly) the most reliable measurement, compare e.g. with infected (test problems, etc...);

Introduction and motivation

Goal of my recent work: influenced by recent events, try predicting the number of COVID-19 victims.

- (Possibly) the most reliable measurement, compare e.g. with infected (test problems, etc...);
- Studied time span: April to predict May (stable conditions);

Goal of my recent work: influenced by recent events, try predicting the number of COVID-19 victims.

- (Possibly) the most reliable measurement, compare e.g. with infected (test problems, etc...);
- Studied time span: April to predict May (stable conditions);
- Idea: if it works, repeat now to predict June/July...

Introduction and motivation

Goal of my recent work: influenced by recent events, try predicting the number of COVID-19 victims.

- (Possibly) the most reliable measurement, compare e.g. with infected (test problems, etc...);
- Studied time span: April to predict May (stable conditions);
- Idea: if it works, repeat now to predict June/July...
- Chosen Countries: Germany, France, Italy.

The model in theory: ODE description

$X(t)$: number of deaths in time. Modeled via a simple 1-d ODE with initial conditions X_0 . Generalized logistic map (S-shaped, exp initially, then flat):

$$X'(t) = \frac{q}{v} X(t) \left(1 - \left(\frac{X(t)}{Q} \right)^v \right) \quad (1)$$

- It has a close form solution (here not written);

The model in theory: ODE description

$X(t)$: number of deaths in time. Modeled via a simple 1-d ODE with initial conditions X_0 . Generalized logistic map (S-shaped, exp initially, then flat):

$$X'(t) = \frac{q}{v} X(t) \left(1 - \left(\frac{X(t)}{Q} \right)^v \right) \quad (1)$$

- It has a close form solution (here not written);
- Depends on the parameter $P = \{q, Q, v\}$ (more next slide).

The model in theory: ODE description

$X(t)$: number of deaths in time. Modeled via a simple 1-d ODE with initial conditions X_0 . Generalized logistic map (S-shaped, exp initially, then flat):

$$X'(t) = \frac{q}{v} X(t) \left(1 - \left(\frac{X(t)}{Q} \right)^v \right) \quad (1)$$

- It has a close form solution (here not written);
- Depends on the parameter $P = \{q, Q, v\}$ (more next slide).
- In others words: assume $\exists P \in \mathbb{R}^3$, s.t. $X^P(t)$ correctly describe the (past and future) number of victims. Want to find P .

The model in theory: parameters description

Do not need to remember the form of $X^P(t)$. It's worth understanding the role of the parameter P .

- $P = \{q, Q, v\}$ are all *strictly* positive.

The model in theory: parameters description

Do not need to remember the form of $X^P(t)$. It's worth understanding the role of the parameter P .

- $P = \{q, Q, v\}$ are all *strictly* positive.
- Q is the carrying capacity ($\lim_{t \rightarrow \infty} X(t) = Q$);

The model in theory: parameters description

Do not need to remember the form of $X^P(t)$. It's worth understanding the role of the parameter P .

- $P = \{q, Q, \nu\}$ are all *strictly* positive.
- Q is the carrying capacity ($\lim_{t \rightarrow \infty} X(t) = Q$);
- $\nu = 1$ gives the *simple* logistic model, while $\nu \rightarrow 0^+$ the *Gompertz* model. Both very famous and used in the literature;

The model in theory: parameters description

Do not need to remember the form of $X^P(t)$. It's worth understanding the role of the parameter P .

- $P = \{q, Q, v\}$ are all *strictly* positive.
- Q is the carrying capacity ($\lim_{t \rightarrow \infty} X(t) = Q$);
- $v = 1$ gives the *simple* logistic model, while $v \rightarrow 0^+$ the *Gompertz* model. Both very famous and used in the literature;

The model in theory: parameters description

Do not need to remember the form of $X^P(t)$. It's worth understanding the role of the parameter P .

- $P = \{q, Q, \nu\}$ are all *strictly* positive.
- Q is the carrying capacity ($\lim_{t \rightarrow \infty} X(t) = Q$);
- $\nu = 1$ gives the *simple* logistic model, while $\nu \rightarrow 0^+$ the *Gompertz* model. Both very famous and used in the literature;
- q is related to the "starting" exponential growth;

The model in theory: parameters description

Do not need to remember the form of $X^P(t)$. It's worth understanding the role of the parameter P .

- $P = \{q, Q, v\}$ are all *strictly* positive.
- Q is the carrying capacity ($\lim_{t \rightarrow \infty} X(t) = Q$);
- $v = 1$ gives the *simple* logistic model, while $v \rightarrow 0^+$ the *Gompertz* model. Both very famous and used in the literature;
- q is related to the "starting" exponential growth;
- **Intuition:** Q the "final" number of victims (10, 30, 50K?), we expect $0 < q \ll 1$ (warning: this comes from the simple logistic) and we prefer $0 < v < 1$ oscillating between the two common models (but no actual math. restriction).

The model in theory: parameters description

Do not need to remember the form of $X^P(t)$. It's worth understanding the role of the parameter P .

- $P = \{q, Q, v\}$ are all *strictly* positive.
- Q is the carrying capacity ($\lim_{t \rightarrow \infty} X(t) = Q$);
- $v = 1$ gives the *simple* logistic model, while $v \rightarrow 0^+$ the *Gompertz* model. Both very famous and used in the literature;
- q is related to the "starting" exponential growth;
- **Intuition:** Q the "final" number of victims (10, 30, 50K?), we expect $0 < q \ll 1$ (warning: this comes from the simple logistic) and we prefer $0 < v < 1$ oscillating between the two common models (but no actual math. restriction).
- Keep in mind: Q large, > 10.000 , v, q small, $\in [0, 1]$.

The model in practice: observed vector

The available dataset are "numbers of total deaths until day t_i ": discrete, limited, ODE trajectory of $X_{X_0}^P(t)$. Use them to infer P .

Fix $T + 1$ times $\{t_i\}_{i=0,\dots,T}$, e.g. 21 days. The observed vector $\mathbf{y} \in \mathbb{R}^{T+1}$ is the random variable defined componentwise as:

$$y_i(\omega) = X_{X_0}^P(t_i) + \eta_i(\omega) \quad (2)$$

where $\eta_i \sim \mathcal{N}(0, \sigma_i^2)$, $i \in \{0, \dots, T\}$.

The model in practice: observed vector

The available dataset are "numbers of total deaths until day t_i ": discrete, limited, ODE trajectory of $X_{X_0}^P(t)$. Use them to infer P .

Fix $T + 1$ times $\{t_i\}_{i=0,\dots,T}$, e.g. 21 days. The observed vector $\mathbf{y} \in \mathbb{R}^{T+1}$ is the random variable defined componentwise as:

$$y_i(\omega) = X_{X_0}^P(t_i) + \eta_i(\omega) \quad (2)$$

where $\eta_i \sim \mathcal{N}(0, \sigma_i^2)$, $i \in \{0, \dots, T\}$.

- In other words, \mathbf{y} is what we measure: "noised truth".

The model in practice: noise

To assume an error of the form

$$y_i(\omega) = X_{X_0}^P(t_i) + \eta_i(\omega) \quad (3)$$

is a **strong** assumption (I know). Need to decide the standard variations σ_i for each day. Arbitrarily.

Quantile formula: $\eta_i \sim N(0, \sigma_i^2) \implies \mathbb{P}[-2\sigma_i \leq \eta_i \leq 2\sigma_i] \geq 95\%$

Read like: "trust $0 \pm 2\sigma_i$ ". Adding the real number y_i : "trust $y_i \pm 2\sigma_i$ ". So we arbitrarily choose:

The model in practice: noise

To assume an error of the form

$$y_i(\omega) = X_{X_0}^P(t_i) + \eta_i(\omega) \quad (3)$$

is a **strong** assumption (I know). Need to decide the standard variations σ_i for each day. Arbitrarily.

Quantile formula: $\eta_i \sim N(0, \sigma_i^2) \implies \mathbb{P}[-2\sigma_i \leq \eta_i \leq 2\sigma_i] \geq 95\%$

Read like: "trust $0 \pm 2\sigma_i$ ". Adding the real number y_i : "trust $y_i \pm 2\sigma_i$ ". So we arbitrarily choose:

- **Case 1:** σ_i s.t. we trust $y_i \pm \frac{y_i}{10}$, i.e. "10% of error";

The model in practice: noise

To assume an error of the form

$$y_i(\omega) = X_{X_0}^P(t_i) + \eta_i(\omega) \quad (3)$$

is a **strong** assumption (I know). Need to decide the standard variations σ_i for each day. Arbitrarily.

Quantile formula: $\eta_i \sim N(0, \sigma_i^2) \implies \mathbb{P}[-2\sigma_i \leq \eta_i \leq 2\sigma_i] \geq 95\%$

Read like: "trust $0 \pm 2\sigma_i$ ". Adding the real number y_i : "trust $y_i \pm 2\sigma_i$ ". So we arbitrarily choose:

- **Case 1:** σ_i s.t. we trust $y_i \pm \frac{y_i}{10}$, i.e. "10% of error";
- **Case 2:** σ_i s.t. we trust $y_i \pm y_i$, i.e. "100% of error";

The model in practice: noise

To assume an error of the form

$$y_i(\omega) = X_{X_0}^P(t_i) + \eta_i(\omega) \quad (3)$$

is a **strong** assumption (I know). Need to decide the standard variations σ_i for each day. Arbitrarily.

Quantile formula: $\eta_i \sim N(0, \sigma_i^2) \implies \mathbb{P}[-2\sigma_i \leq \eta_i \leq 2\sigma_i] \geq 95\%$

Read like: "trust $0 \pm 2\sigma_i$ ". Adding the real number y_i : "trust $y_i \pm 2\sigma_i$ ". So we arbitrarily choose:

- **Case 1:** σ_i s.t. we trust $y_i \pm \frac{y_i}{10}$, i.e. "10% of error";
- **Case 2:** σ_i s.t. we trust $y_i \pm y_i$, i.e. "100% of error";
- **Warning:** the inference algorithm [not yet described] is destroyed for higher errors, but not with our cases - checked with toy model data.

Inference technique: MCMC

- We have the model $X^P(t)$, the noise structure and the data.
Last step: choose how to infer $P \in \mathbb{R}^3$.

Inference technique: MCMC

- We have the model $X^P(t)$, the noise structure and the data.
Last step: choose how to infer $P \in \mathbb{R}^3$.
- in general, I am mainly into the topic "Bayesian Inverse Problems", where a common technique is the "Preconditioned Monte Carlo" algorithm (pCN);

Inference technique: MCMC

- We have the model $X^P(t)$, the noise structure and the data.
Last step: choose how to infer $P \in \mathbb{R}^3$.
- in general, I am mainly into the topic "Bayesian Inverse Problems", where a common technique is the "Preconditioned Monte Carlo" algorithm (pCN);
- most suitable for infinite-dimensional cases, large data, inversion of PDEs...

Inference technique: MCMC

- We have the model $X^P(t)$, the noise structure and the data.
Last step: choose how to infer $P \in \mathbb{R}^3$.
- in general, I am mainly into the topic "Bayesian Inverse Problems", where a common technique is the "Preconditioned Monte Carlo" algorithm (pCN);
- most suitable for infinite-dimensional cases, large data, inversion of PDEs...
- here: only 20 days, 3 real parameters, a simple 1-dim ODE with close form solution.

Inference technique: MCMC

- We have the model $X^P(t)$, the noise structure and the data.
Last step: choose how to infer $P \in \mathbb{R}^3$.
- in general, I am mainly into the topic "Bayesian Inverse Problems", where a common technique is the "Preconditioned Monte Carlo" algorithm (pCN);
- most suitable for infinite-dimensional cases, large data, inversion of PDEs...
- here: only 20 days, 3 real parameters, a simple 1-dim ODE with close form solution.
- But I was curious and insisted: let's use the pCN algorithm anyway!

Inference technique: MCMC

- We have the model $X^P(t)$, the noise structure and the data.
Last step: choose how to infer $P \in \mathbb{R}^3$.
- in general, I am mainly into the topic "Bayesian Inverse Problems", where a common technique is the "Preconditioned Monte Carlo" algorithm (pCN);
- most suitable for infinite-dimensional cases, large data, inversion of PDEs...
- here: only 20 days, 3 real parameters, a simple 1-dim ODE with close form solution.
- But I was curious and insisted: let's use the pCN algorithm anyway!
- **In brief:** we infer P by using a Monte Carlo algorithm.

The Bayesian setting: how to infer P ?

- The idea is: *given* the data \mathbf{y} , find P . Equivalent: find $\mathbb{P}[P|\mathbf{y}]$.

The Bayesian setting: how to infer P ?

- The idea is: *given* the data \mathbf{y} , find P . Equivalent: find $\mathbb{P}[P|\mathbf{y}]$.
- Step 1: assume to have a probability density on \mathbb{R}^3 , called the **prior**. Blind guess about P , i.e. $\mathbb{P}[P \in dx]$. (more later);

The Bayesian setting: how to infer P ?

- The idea is: *given* the data \mathbf{y} , find P . Equivalent: find $\mathbb{P}[P|\mathbf{y}]$.
- Step 1: assume to have a probability density on \mathbb{R}^3 , called the **prior**. Blind guess about P , i.e. $\mathbb{P}[P \in dx]$. (more later);
- Step 2: interpret the read data as $\mathbb{P}[\mathbf{y}|P]$ (**likelihood**, obtain a complete formula by using the Gaussian noise, etc...)

The Bayesian setting: how to infer P ?

- The idea is: *given* the data \mathbf{y} , find P . Equivalent: find $\mathbb{P}[P|\mathbf{y}]$.
- Step 1: assume to have a probability density on \mathbb{R}^3 , called the **prior**. Blind guess about P , i.e. $\mathbb{P}[P \in dx]$. (more later);
- Step 2: interpret the read data as $\mathbb{P}[\mathbf{y}|P]$ (**likelihood**, obtain a complete formula by using the Gaussian noise, etc...)
- Step 3: conclude by using the Bayes formula:

$$\mathbb{P}[P|\mathbf{y}] \propto \mathbb{P}[\mathbf{y}|P] \times \mathbb{P}[P] \quad (4)$$

where $\mathbb{P}[P|\mathbf{y}]$ is the **posterior** measure.

The Bayesian setting: how to infer P ?

- The idea is: *given* the data \mathbf{y} , find P . Equivalent: find $\mathbb{P}[P|\mathbf{y}]$.
- Step 1: assume to have a probability density on \mathbb{R}^3 , called the **prior**. Blind guess about P , i.e. $\mathbb{P}[P \in dx]$. (more later);
- Step 2: interpret the read data as $\mathbb{P}[\mathbf{y}|P]$ (**likelihood**, obtain a complete formula by using the Gaussian noise, etc...)
- Step 3: conclude by using the Bayes formula:

$$\mathbb{P}[P|\mathbf{y}] \propto \mathbb{P}[\mathbf{y}|P] \times \mathbb{P}[P] \quad (4)$$

where $\mathbb{P}[P|\mathbf{y}]$ is the **posterior** measure.

- In brief: fix a belief about P , i.e. the prior. Read the data, obtain the likelihood. Multiply them to have the posterior probability, containing information about the "true" P given the data (details omitted on purpose).

The Bayesian setting: how to infer P ?

- The idea is: *given* the data \mathbf{y} , find P . Equivalent: find $\mathbb{P}[P|\mathbf{y}]$.
- Step 1: assume to have a probability density on \mathbb{R}^3 , called the **prior**. Blind guess about P , i.e. $\mathbb{P}[P \in dx]$. (more later);
- Step 2: interpret the read data as $\mathbb{P}[\mathbf{y}|P]$ (**likelihood**, obtain a complete formula by using the Gaussian noise, etc...)
- Step 3: conclude by using the Bayes formula:

$$\mathbb{P}[P|\mathbf{y}] \propto \mathbb{P}[\mathbf{y}|P] \times \mathbb{P}[P] \quad (4)$$

where $\mathbb{P}[P|\mathbf{y}]$ is the **posterior** measure.

- In brief: fix a belief about P , i.e. the prior. Read the data, obtain the likelihood. Multiply them to have the posterior probability, containing information about the "true" P given the data (details omitted on purpose).
- **Infer P = sample from the posterior**

The pCN Monte Carlo method

Let $\mathbb{P}[P|\mathbf{y}]$ be the posterior on \mathbb{R}^3 . Let $0 < \beta < 1$ the speed parameter, and the acceptance probability, for $v, u \in \mathbb{R}^3$:

$$a(u, v) \doteq \min\left\{1, \frac{\mathbb{P}(\mathbf{y}|v)}{\mathbb{P}(\mathbf{y}|u)}\right\} \quad (5)$$

To produce a **single sample**, construct a chain $\{x_i\}_{i \in \mathbb{N}}$ as follows:

1. set $x_0 \in \mathbb{R}^n$ **arbitrarily**. Then, for each $k > 0$:
2. sample a point $R \in \mathbb{R}^3$ from the **gaussian** prior $\mathbb{P}(P \in dx)$;
3. propose a candidate as $\hat{x}_k = \sqrt{(1 - \beta^2)}x_{k-1} + \beta R$;
4. accept it (i.e. set $x_k = \hat{x}_k$) with probability $a(x_{k-1}, \hat{x}_k)$;
5. (accepted or not) repeat from 2;

Run 120.000 multiple chains in parallel, different starting point, each stopped after 250.000 iterations. Beta around 0.05, s.t. acceptance rate $\sim 25\%$. **Last missing info: which prior?**

Prior measure on P : part 1

Before moving to the numerical results, we need to clarify the structure of the prior measure. It influences the results.

- The pCN Monte Carlo algorithm *requires* a centered Gaussian prior (on \mathbb{R}^3).

Prior measure on P : part 1

Before moving to the numerical results, we need to clarify the structure of the prior measure. It influences the results.

- The pCN Monte Carlo algorithm *requires* a centered Gaussian prior (on \mathbb{R}^3).
- Problematic: $P \in \mathbb{R}^3$ is strictly positive...so we just rejected the negative proposals...(pitfall?)...

Prior measure on P : part 1

Before moving to the numerical results, we need to clarify the structure of the prior measure. It influences the results.

- The pCN Monte Carlo algorithm *requires* a centered Gaussian prior (on \mathbb{R}^3).
- Problematic: $P \in \mathbb{R}^3$ is strictly positive...so we just rejected the negative proposals...(pitfall?)...
- and we assumed a prior structure with covariance matrix

$$\begin{pmatrix} \sigma_q^2 & 0 & 0 \\ 0 & \sigma_Q^2 & 0 \\ 0 & 0 & \sigma_v^2 \end{pmatrix} \quad (6)$$

Prior measure on P : part 1

Before moving to the numerical results, we need to clarify the structure of the prior measure. It influences the results.

- The pCN Monte Carlo algorithm *requires* a centered Gaussian prior (on \mathbb{R}^3).
- Problematic: $P \in \mathbb{R}^3$ is strictly positive...so we just rejected the negative proposals...(pitfall?)...
- and we assumed a prior structure with covariance matrix

$$\begin{pmatrix} \sigma_q^2 & 0 & 0 \\ 0 & \sigma_Q^2 & 0 \\ 0 & 0 & \sigma_v^2 \end{pmatrix} \quad (6)$$

- Therefore: we only need to choose the σ , and then we can move on the simulations!

Prior measure on P : part 2

Goal of this slide: explain how the values for σ_q , σ_Q and σ_v have been chosen.

- The idea is: the prior represents the "guess", the "default" choice. But if "nothing is done", the phenomenon propagates exponentially. Therefore we run an exponential interpolation, and register the number of victims in 1 Month: K . Then choose σ_Q in order to have the Gaussian 95% quantile just below this quantity;

Prior measure on P : part 2

Goal of this slide: explain how the values for σ_q , σ_Q and σ_v have been chosen.

- The idea is: the prior represents the "guess", the "default" choice. But if "nothing is done", the phenomenon propagates exponentially. Therefore we run an exponential interpolation, and register the number of victims in 1 Month: K . Then choose σ_Q in order to have the Gaussian 95% quantile just below this quantity;
- Dealing with σ_q and σ_v , they have been chosen in order to have that quantile between 0 and 1.

Prior measure on P : part 2

Goal of this slide: explain how the values for σ_q , σ_Q and σ_v have been chosen.

- The idea is: the prior represents the "guess", the "default" choice. But if "nothing is done", the phenomenon propagates exponentially. Therefore we run an exponential interpolation, and register the number of victims in 1 Month: K . Then choose σ_Q in order to have the Gaussian 95% quantile just below this quantity;
- Dealing with σ_q and σ_v , they have been chosen in order to have that quantile between 0 and 1.
- These idea formalize the intuition that we expect parameters in these range, without *forcing* any constraint.

How to read the results

- Now all the parameters, assumptions, hypothesis should be clear. In particular, we are using a possibly-not-so-good Monte Carlo algorithm with strong assumptions on the noise and the prior;

How to read the results

- Now all the parameters, assumptions, hypothesis should be clear. In particular, we are using a possibly-not-so-good Monte Carlo algorithm with strong assumptions on the noise and the prior;
- We remark: by reading the data we infer a 3-dimensional probability measure for $P...$

How to read the results

- Now all the parameters, assumptions, hypothesis should be clear. In particular, we are using a possibly-not-so-good Monte Carlo algorithm with strong assumptions on the noise and the prior;
- We remark: by reading the data we infer a 3-dimensional probability measure for $P...$
- we clustered it by the k-means algorithm and selected three set of parameters:


How to read the results

- Now all the parameters, assumptions, hypothesis should be clear. In particular, we are using a possibly-not-so-good Monte Carlo algorithm with strong assumptions on the noise and the prior;
- We remark: by reading the data we infer a 3-dimensional probability measure for P ...
- we clustered it by the k-means algorithm and selected three set of parameters:
- the **worst** case, i.e. the triple with the highest value for Q ; the **best** (converse) and the **expectation**.

How to read the results


- Now all the parameters, assumptions, hypothesis should be clear. In particular, we are using a possibly-not-so-good Monte Carlo algorithm with strong assumptions on the noise and the prior;
- We remark: by reading the data we infer a 3-dimensional probability measure for P ...
- we clustered it by the k-means algorithm and selected three set of parameters:
- the **worst** case, i.e. the triple with the highest value for Q ; the **best** (converse) and the **expectation**.
- We have plots for the "10%" error scenario, as well as for the "100%" (*meaning explained before*)

Results: Italy, assuming 10% of "error"




Results: France, assuming 10% of "error"

France: prediction until 01.06 using 3 weeks of data.




Results: Germany, assuming 10% of "error"

Germany: prediction until 01.06 using 3 weeks of data.




Results: Italy, assuming 100% of "error"




Results: France, assuming 100% of "error"

France: prediction until 01.06 using 3 weeks of data.



Results: Germany, assuming 100% of "error"

Germany: prediction until 01.06 using 3 weeks of data.



Conclusions

Despite various theoretical issues, the results give some degree of satisfaction. *But they do not seem to work with Spain, UK (I quickly tested).*

- A next step can be to use the last 20 days of data to predict for June/July;

Conclusions

Despite various theoretical issues, the results give some degree of satisfaction. *But they do not seem to work with Spain, UK (I quickly tested).*

- A next step can be to use the last 20 days of data to predict for June/July;
- Or try with another more appropriate Monte Carlo methods with better prior measures;

Conclusions

Despite various theoretical issues, the results give some degree of satisfaction. *But they do not seem to work with Spain, UK (I quickly tested).*

- A next step can be to use the last 20 days of data to predict for June/July;
- Or try with another more appropriate Monte Carlo methods with better prior measures;
- Or instead of the Bayesian approach, try maximizing the Likelihood (i.e. probabilistic least square);

Conclusions

Despite various theoretical issues, the results give some degree of satisfaction. *But they do not seem to work with Spain, UK (I quickly tested).*

- A next step can be to use the last 20 days of data to predict for June/July;
- Or try with another more appropriate Monte Carlo methods with better prior measures;
- Or instead of the Bayesian approach, try maximizing the Likelihood (i.e. probabilistic least square);
- Or simply to consider this example as a nice application of Monte Carlo Bayesian techniques, before moving back to the main PhD topic.

Thanks for the attention!