NUMERICAL ANALYSIS OF A LAX–WENDROFF TYPE LATTICE BOLTZMANN SCHEME FOR THE ADVECTION EQUATION

T. BELLOTTI[1], F. CAETANO[2], F. DUBOIS[3,2], AND B. GRAILLE[2]

Abstract. A linear monodimensional lattice Boltzmann scheme with three velocities tackling the advection equation is investigated. We prove that it is second-order accurate and provide necessary and sufficient stability conditions through an explicit formula involving the two relaxation parameters. Relying on existing results for finite difference schemes, we prove convergence for smooth initial data with initialization of the non-conserved moments at equilibrium. Several numerical simulations illustrate these theoretical findings.

Keywords: stability, Cohn-Schur algorithm, convergence
AMS2000: 65M12

1. Introduction

We investigate a three-velocity lattice Boltzmann scheme, see [38, 24] for a general introduction, designed to approximate the solution of the linear advection equation with second-order accuracy. The scheme under investigation fully aligns with the Lax-Wendroff method [30] for a specific choice of free parameters, and retains its centered character in full generality.

Finite-velocity models with three velocities have a long history in the literature; the earliest reference we found is by Broadwell [11], where a finite-velocity approximation of the Boltzmann equation (see [13] for an earlier example with two velocities)—reduced to a set of three equations under symmetry assumptions—is introduced. Almost three decades later, Qian, d'Humières, and Lallemand [33] proposed several lattice Boltzmann schemes, including a three-velocity model, to approximate the Navier-Stokes system. We note that the target equation we address here—the linear advection equation—is indeed simpler than those in the aforementioned references, which focused on either a discrete-velocity Boltzmann equation or the Navier-Stokes system.

In the case of a scalar equation and a three-velocity model considered here, the numerical scheme features four free parameters: two linear equilibria and two relaxation parameters. In this contribution, the first two are fixed to achieve second-order accuracy for the advection equation. The relaxation parameters remain free

[1] Université Paris-Saclay, CNRS, CentraleSupélec, Laboratoire EM2C & Fédération de Mathématiques de CentraleSupélec, 91190, Gif-sur-Yvette, France
[2] Université Paris-Saclay, CNRS, Laboratoire de mathématiques d'Orsay, 91405, Orsay, France
[3] Conservatoire National des Arts et Métiers, MACS laboratory, Paris, France
Email address: benjamin.graille@universite-paris-saclay.fr.
Date: October 15, 2026.

and solely influence stability, whereas accuracy is uniform in them. When they both equal one, we obtain a relaxation scheme which is the Lax-Wendroff method.

An essential question for any numerical scheme addressing evolutionary partial differential equations is its *stability*, and lattice Boltzmann methods are no exception. Most of the present work focuses on this aspect. Let us therefore review the main facets of and contributions to this topic.

A first approach relies on von Neumann stability analysis, and boils down to considering linear or linearized schemes, applying a spatial Fourier transform, and verifying whether the spectral radius of the resulting matrix is less than one for all frequencies. This method was adopted in [36, 29]. Given that the matrix size corresponds to the number of discrete velocities, the scheme may include numerous free parameters, and since all frequencies must be considered, this stability criterion is typically verified numerically. This approach has been widely used in subsequent works. Notably, Ginzburg and collaborators [22] derived partially explicit stability conditions—thus not solely numerically-checked—for schemes with two relaxation parameters whose sum equals two, using the Cohn-Schur algorithm, see [32, 37].

A second major research stream on the stability of lattice Boltzmann algorithms focuses on weighted L^2 stability and the so-called "stability structure" [3, 26, 34]. Although this concept is primarily designed for linearized schemes, it can also yield valuable insights in certain non-linear settings, as demonstrated in [25]. This approach involves constructing a weighted vector norm to evaluate the discrete distribution functions, ensuring that both the transport and—more critically—the relaxation steps of the algorithm become L^2-contractive under this norm. In this way, the elementary bound $\|A^n\| \leq \|A\|^n$, where A represents the numerical scheme, can be effectively employed, as $\|A\| \leq 1$.

A third set of contributions focuses on L^∞-stability and maximum principles, which are particularly effective for scalar target equations and generally more demanding than mere L^2 stability. This stringency enables their direct applicability to non-linear problems. This approach was pioneered by Dellacherie [16] in a linear framework and later extended to non-linear problems through various notions of monotonicity, as seen in [12, 4, 1, 2].

We note that the previously discussed stability notions sometimes lead to convergence proofs of the numerical solution to the target equation, as demonstrated in [25, 16, 12, 4, 1, 2]. This will also be one of the goals of the present contribution.

Our work conducts a comprehensive L^2 stability analysis of the considered algorithm by supplementing the von Neumann analysis with a discussion of eigenvalue multiplicities, in addition to the mere analysis of spectral radii (i.e., their moduli). We derive explicit necessary and sufficient stability conditions, obtained using the Cohn-Schur algorithm [32, 37], in terms of the Courant number and the two relaxation parameters of the scheme.

The article is structured as follows. In Section 2, we introduce the three-velocities lattice Boltzmann scheme, whose equilibria are fully determined in Section 3 to achieve second-order consistency with the target equation. Section 4 defines stability and recapitulates the necessary and sufficient stability conditions. Proofs, which constitute the bulk of the paper, are provided in Section 5. In Section 6, we use known results on finite difference methods to prove convergence of the conserved moment of the lattice Boltzmann scheme for smooth initial data and initialization of the non-conserved moments "at equilibrium". Additionally, we study discrete

wave propagation by the scheme for smooth data. Finally, Section 7 presents several numerical experiments that showcase the expected wave propagation patterns, convergence for smooth and non-smooth initial data, and empirical second-order convergence of an adaptation of the scheme to the (non-linear) inviscid Burgers' equation.

2. Description of the scheme

In this section, the investigated lattice Boltzmann scheme is presented. We tackle the approximation of the following Cauchy problem

$$\partial_t u(t,x) + V\partial_x u(t,x) = 0 \qquad \text{for} \quad t>0,\ x\in\mathbb{R}, \tag{1}$$

$$u(0,x) = u^\circ(x) \qquad \text{for} \quad x\in\mathbb{R}, \tag{2}$$

a linear scalar advection equation with constant velocity $V\in\mathbb{R}$. Space is discretized using a uniform grid $(j\Delta x)_{j\in\mathbb{Z}}$ with $\Delta x>0$, and time relying on $(n\Delta t)_{n\in\mathbb{N}}$ with $\Delta t>0$. We introduce the ratio $\lambda := \Delta x/\Delta t>0$, which is assumed given and fixed whatever the value of Δx.

In this paper, we consider a linear three-velocities monodimensional lattice Boltzmann scheme specified by three discrete velocities 0, λ, and $-\lambda$, the associated particle distribution functions f_0, f_+, and f_-, and three polynomials defining the three moments denoted by u, v, and w,

$$P_0 = 1, \qquad P_1 = X, \qquad P_2 = \tfrac{1}{2}X^2,$$
$$u = f_0+f_++f_-, \qquad v = \lambda f_+ - \lambda f_-, \qquad w = \tfrac{1}{2}\lambda^2 f_+ + \tfrac{1}{2}\lambda^2 f_-.$$

The three equilibrium values of the moments read

$$u^{\mathrm{eq}}(u) = u, \quad v^{\mathrm{eq}}(u) = c\lambda u, \quad w^{\mathrm{eq}}(u) = \tfrac{1}{2}d\lambda^2 u,$$

where c and d are real values. For the moment, c and d are kept free: their value to ensure consistency with (1) is discussed in Section 3. Finally, we introduce two relaxation parameters for the non-conserved moments s_v and s_w; and the centered relaxation parameters $t_v = s_v - 1$ and $t_w = s_w - 1$. We use vectorial notations by introducing the vector of the particle distribution functions $\boldsymbol{f} = (f_0, f_+, f_-)^T$ and the vector of the moments $\boldsymbol{m} = (u,v,w)^T$ linked by the matrix relation $\boldsymbol{m} = \boldsymbol{M}\boldsymbol{f}$ with

$$\boldsymbol{M} = \begin{pmatrix} 1 & 1 & 1 \\ 0 & \lambda & -\lambda \\ 0 & \frac{\lambda^2}{2} & \frac{\lambda^2}{2} \end{pmatrix}, \qquad \boldsymbol{M}^{-1} = \begin{pmatrix} 1 & 0 & -\frac{2}{\lambda^2} \\ 0 & \frac{1}{2\lambda} & \frac{1}{\lambda^2} \\ 0 & -\frac{1}{2\lambda} & \frac{1}{\lambda^2} \end{pmatrix}.$$

Then, the algorithm proceeds as follows. Given $(\boldsymbol{m}_j^0)_{j\in\mathbb{Z}}$ arbitrary (which however must be eventually linked to the initial datum u° to establish a connection with the Cauchy problem at hand), for $n\in\mathbb{N}$, compute

- Relaxation. For $j\in\mathbb{Z}$, $n\in\mathbb{N}$,

$$\boldsymbol{m}_j^{n,\star} = \mathbf{diag}(1, 1-s_v, 1-s_w)\,\boldsymbol{m}_j^n + \mathbf{diag}(0, s_v, s_w)\,\boldsymbol{m}^{\mathrm{eq}}(u_j^n). \tag{3}$$

- Transport. For $j\in\mathbb{Z}$, $n\in\mathbb{N}$,

$$f_{0,j}^{n+1} = f_{0,j}^{n,\star}, \qquad f_{\pm,j}^{n+1} = f_{\pm,j\mp\lambda\Delta t/\Delta x}^{n,\star} = f_{\pm,j\mp1}^{n,\star}. \tag{4}$$

As the problem is linear, it is convenient to introduce a discrete Fourier transform in space, see [37, Chapter 2] for the details. We denote any space-grid function having undergone a Fourier transformation by a hat, so that the scheme on the distribution functions reads

$$\hat{\boldsymbol{f}}^{n+1}(\xi) = \boldsymbol{G}(\xi)\hat{\boldsymbol{f}}^{n}, \qquad \forall \xi \in [-\pi, \pi],$$

where $\boldsymbol{G}$ is called "amplification matrix", and given by

$$\boldsymbol{G}(\xi) = \boldsymbol{T}(\xi)\boldsymbol{M}^{-1}\boldsymbol{R}\boldsymbol{M},$$

with

$$\boldsymbol{T}(\xi) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & e^{-\mathrm{i}\xi} & 0 \\ 0 & 0 & e^{\mathrm{i}\xi} \end{pmatrix}, \qquad \boldsymbol{R} = \begin{pmatrix} 1 & 0 & 0 \\ s_v c\lambda & 1-s_v & 0 \\ \frac{1}{2}s_w d\lambda^2 & 0 & 1-s_w \end{pmatrix}.$$

After algebra, we obtain

(5)
$$\boldsymbol{G} = \begin{pmatrix} 1-s_w d & s_w(1-d) & s_w(1-d) \\ \frac{e^{-\mathrm{i}\xi}}{2}(s_w d+s_v c) & \frac{e^{-\mathrm{i}\xi}}{2}(2-s_w(1-d)-s_v(1-c)) & \frac{e^{-\mathrm{i}\xi}}{2}(-s_w(1-d)+s_v(1+c)) \\ \frac{e^{\mathrm{i}\xi}}{2}(s_w d-s_v c) & \frac{e^{\mathrm{i}\xi}}{2}(-s_w(1-d)+s_v(1-c)) & \frac{e^{\mathrm{i}\xi}}{2}(2-s_w(1-d)-s_v(1+c)) \end{pmatrix}.$$

The matrix $\boldsymbol{G}$ can be rewritten in terms of the centered relaxation parameters t_v and t_w as

(6)
$$\boldsymbol{G} = \begin{pmatrix} 1-d-t_w d & (1+t_w)(1-d) & (1+t_w)(1-d) \\ \frac{e^{-\mathrm{i}\xi}}{2}(d+c+t_w d+t_v c) & \frac{e^{-\mathrm{i}\xi}}{2}(d+c-t_w(1-d)-t_v(1-c)) & \frac{e^{-\mathrm{i}\xi}}{2}(d+c-t_w(1-d)+t_v(1+c)) \\ \frac{e^{\mathrm{i}\xi}}{2}(d-c+t_w d-t_v c) & \frac{e^{\mathrm{i}\xi}}{2}(d-c-t_w(1-d)+t_v(1-c)) & \frac{e^{\mathrm{i}\xi}}{2}(d-c-t_w(1-d)-t_v(1+c)) \end{pmatrix}.$$

Remark 1. The amplification matrix does not depend on the choice of the second-order moment P_2 as long as two conditions are met. First, P_2 reads $\alpha X^2 + \beta$, for any real values $\alpha \neq 0$ and β. Second, the equilibrium values of the particle distribution functions are unchanged [20]. In particular, a classical choice for the second-order moment is $\tilde{P}_2 = 3X^2 - 2\lambda^2$ for which the corresponding matrix of the moments has orthogonal rows

$$\tilde{\boldsymbol{M}} = \begin{pmatrix} 1 & 1 & 1 \\ 0 & \lambda & -\lambda \\ -2\lambda^2 & \lambda^2 & \lambda^2 \end{pmatrix}, \qquad \tilde{\boldsymbol{M}}^{-1} = \begin{pmatrix} \frac{1}{3} & 0 & -\frac{2}{6\lambda^2} \\ \frac{1}{3} & \frac{1}{2\lambda} & \frac{1}{6\lambda^2} \\ \frac{1}{3} & -\frac{1}{2\lambda} & \frac{1}{6\lambda^2} \end{pmatrix}.$$

The amplification matrix $\boldsymbol{G}$ is then unchanged as long as $\tilde{w}^{\mathrm{eq}} = (3d-2)\lambda^2 u$.

3. Consistency

We now apply the algebraic framework developed by Bellotti et al. [9], which establishes an exact correspondence between a lattice Boltzmann scheme and an equivalent multistep finite-difference scheme. This correspondence enables the systematic derivation of the associated macroscopic equations and provides access to the classical finite-difference analysis framework.

Proposition 2 (Consistency)**.** *Assume that $s_v, s_w \neq 0$. Then, the modified equation of the scheme truncated at second order, cf.* [5, Theorem 3.7]*, is given by*

$$\partial_t u + \lambda c \partial_x u - \lambda^2 \Delta t\left(\tfrac{1}{2} - \tfrac{1}{s_v}\right)(d-c^2)\partial_{xx} u = \mathcal{O}(\Delta t^2). \tag{7}$$

Therefore, the scheme is first-order accurate with respect to (1) *upon selecting* $\lambda c = V$. *Moreover, the scheme becomes second-order accurate regardless of* s_v *by taking* $d = c^2$. *Under this choice, the modified equation truncated at third order is*

$$\partial_t u + V\partial_x u = -\lambda^3 \Delta t^2 c(1-c^2)\big(\tfrac{1}{12} - \big(\tfrac{1}{s_v}-\tfrac{1}{2}\big)\big(\tfrac{1}{s_w}-\tfrac{1}{2}\big)\big)\partial_{xxx} u + \mathcal{O}(\Delta t^3). \tag{8}$$

Proof. Equation (7) is given by [5, Theorem 3.7]. Equation (8) can be obtained by analogous computations at the next order. □

From now on in the paper, we adopt the choice $d = c^2$, so that d is no longer a free parameter. The product $(\frac{1}{s_v}-\frac{1}{2})(\frac{1}{s_w}-\frac{1}{2})$ is the so-called "magic parameter" [17], and the fact of achieving third-order accuracy by taking it equal to $1/12$ is well-known within the lattice Boltzmann community, see [21, 28].

For the previous consistency analysis, we have assumed that $s_v, s_w \neq 0$. When one or both vanish, the study must be adapted accordingly. When $s_v = 0$, the scheme is not consistent with the target equation. Assuming $s_v \neq 0$, when $s_w = 0$, it is only first-order accurate despite having taken $d = c^2$, unless $s_v = 2$, in which case the scheme becomes second-order accurate. This fact can be seen by following the procedure by [5] for the specific scheme at hand, or by performing the "equivalent equation" analysis following the guidelines of [18, 19].

We shall come back to consistency from the standpoint of Fourier analysis in Sec. 6, dedicated to convergence. Indeed, upon building a suitable link between $\boldsymbol{m}^0$ and the initial datum u° of the Cauchy problem, see (2), we expect and hope that u_j^n be a "good approximation" of the solution $u(n\Delta t, j\Delta x)$.

4. Stability

This section aims to provide an explicit relation between the relaxation parameters ensuring stability of the scheme. This condition is both necessary and sufficient. We first recall some definitions and properties of polynomials with complex coefficients.

4.1. Reminder on complex polynomials.

We begin by introducing definitions and notation for several classes of polynomials relevant to the stability analysis, which are characterized by the location of their zeros with respect to the unit circle. This section is based on [32, 31, 37]. We emphasize that these tools have already been used in the stability analysis of lattice Boltzmann schemes; see, for instance, [22], where the stability condition reduces to locating the roots of a second-degree polynomial.

Let $\varphi \in \mathbb{C}[z]$ be a polynomial with complex coefficients of degree d,

$$\varphi = a_d z^d + \ldots + a_0 = \sum_{\ell=0}^{d} a_\ell z^\ell.$$

We say that φ is of exact degree d if a_d is not zero.

Definition 3 (Schur polynomial)**.** The polynomial φ is a *Schur polynomial* if all its roots r satisfy $|r| < 1$. We denote by $\mathscr{S}$ the set of the Schur polynomials.

Definition 4 (von Neumann polynomial)**.** The polynomial φ is a *von Neumann polynomial* if all its roots r satisfy $|r| \leq 1$. We denote by $\mathscr{N}$ the set of the von Neumann polynomials.

Definition 5 (simple von Neumann polynomial)**.** The polynomial φ is a *simple von Neumann polynomial* if φ is a von Neumann polynomial and its roots on the unit circle are simple roots. We denote by $\mathscr{N}^s$ the set of the simple von Neumann polynomials.

We finally define the Cohn-Schur operator $\mathscr{C} : \mathbb{C}[z] \to \mathbb{C}[z]$ an application that decreases the degree. For any polynomial φ of exact degree d we first define the polynomial $\varphi^\star$ by

$$\varphi^\star = \bar{a}_0 z^d + \ldots + \bar{a}_d = \sum_{\ell=0}^{d} \bar{a}_{d-\ell}\, z^\ell,$$

where the bar on the coefficients of φ denotes the complex conjugate and then

$$\mathscr{C}(\varphi) = \frac{\varphi^\star(0)\,\varphi - \varphi(0)\,\varphi^\star}{z}.$$

Remark that $\mathscr{C}(\varphi)$ is indeed a polynomial as the constant term of $\varphi^\star(0)\varphi - \varphi(0)\varphi^\star$ vanishes. It is easy to see that the degree of $\mathscr{C}(\varphi)$ is less than that of φ. We also denote by a prime φ' the derivative of the polynomial φ with respect to its determinate.

The next three theorems give recursive tests for Schur, von Neumann and simple von Neumann polynomials [32].

Theorem 6 (Miller 6.2 page 404)**.** *The polynomial φ is a Schur polynomial of exact degree d if, and only if, $|\varphi(0)| < |\varphi^\star(0)|$ and $\mathscr{C}(\varphi)$ is a Schur polynomial of exact degree $d-1$.*

Theorem 7 (Miller 6.1 page 403)**.** *The polynomial φ is a von Neumann polynomial of exact degree d if, and only if, either*

(a) *$|\varphi(0)| < |\varphi^\star(0)|$ and $\mathscr{C}(\varphi)$ is a von Neumann polynomial of exact degree $d-1$ or*
(b) *$\mathscr{C}(\varphi)$ is identically zero and φ' is a von Neumann polynomial of exact degree $d-1$.*

Theorem 8 (Miller 6.3 page 404)**.** *The polynomial φ is a simple von Neumann polynomial of exact degree d if, and only if, either*

(a) *$|\varphi(0)| < |\varphi^\star(0)|$ and $\mathscr{C}(\varphi)$ is a simple von Neumann polynomial of exact degree $d-1$ or*
(b) *$\mathscr{C}(\varphi)$ is identically zero and φ' is a Schur polynomial of exact degree $d-1$.*

The proofs of these three theorems are lying on Rouché's theorem from complex analysis [35].

Definition 9. We denote

$$\mathscr{I} = \Big\{\varphi \in \mathbb{C}[z] \text{ s.t. } |\varphi(0)| < |\varphi^\star(0)|\Big\} \quad \text{and} \quad \mathscr{E} = \Big\{\varphi \in \mathbb{C}[z] \text{ s.t. } \mathscr{C}(\varphi) = 0\Big\}.$$

With these notations, theorems 6, 7, and 8 read

$$\varphi \in \mathscr{S} \iff \varphi \in \mathscr{I} \text{ and } \mathscr{C}(\varphi) \in \mathscr{S}, \tag{9}$$

$$\varphi \in \mathscr{N} \iff \Big(\varphi \in \mathscr{I} \text{ and } \mathscr{C}(\varphi) \in \mathscr{N}\Big) \text{ or } \Big(\varphi \in \mathscr{E} \text{ and } \varphi' \in \mathscr{N}\Big), \tag{10}$$

$$\varphi \in \mathscr{N}^s \iff \Big(\varphi \in \mathscr{I} \text{ and } \mathscr{C}(\varphi) \in \mathscr{N}^s\Big) \text{ or } \Big(\varphi \in \mathscr{E} \text{ and } \varphi' \in \mathscr{S}\Big). \tag{11}$$

Note that we cannot have simultaneously $\varphi \in \mathscr{I} \cap \mathscr{E}$. Indeed, if φ is of exact degree d and belongs to $\mathscr{I}$, the polynomial $\mathscr{C}(\varphi)$ is of exact degree $d-1$ as its term of degree $d-1$ is equal to $|\varphi^\star(0)|^2 - |\varphi(0)|^2 > 0$.

4.2. **Definition of the stability.** We consider in this contribution the ℓ^2-stability. As the amplification matrix $\boldsymbol{G}$ given in (6) does not depend on the time step Δt or the space step Δx, the definition for the stability reads as follow.

Definition 10 (Stability)**.** The lattice Boltzmann scheme characterized by the amplification matrix $\boldsymbol{G}$ is said stable if there exists one constant $C > 0$ such that

$$\forall n \in \mathbb{N}, \quad \forall \xi \in [-\pi, \pi] \qquad |\boldsymbol{G}(\xi)^n| \leq C.$$

The fact that this definition be fulfilled is equivalent to the ℓ^2 control

$$\Delta x \sum_{j \in \mathbb{Z}} |\boldsymbol{f}_j^n|^2 \leq C^2 \Delta x \sum_{j \in \mathbb{Z}} |\boldsymbol{f}_j^0|^2$$

for all $n \in \mathbb{N}$ and for all $\boldsymbol{f}_j^0$ thanks to the Parseval's relation. The following result gives a necessary condition for stability.Its sole necessary character comes from the fact that is does not ensure a bound uniformly in ξ.

Proposition 11. *If the lattice Boltzmann scheme characterized by the amplification matrix $\boldsymbol{G}$ is stable, then the minimal polynomial of $\boldsymbol{G}$ is a simple von Neumann polynomial for any $\xi \in [-\pi, \pi]$.*

Proof. The proof can be found in [15, Lemma 3]. For readibility reasons, we present it here briefly. The key point is the Dunford-Jordan decomposition of the complex square matrices. The matrix $\boldsymbol{G}$ is equivalent to a block diagonal matrix that reads

$$\begin{pmatrix} J_{\lambda_1} & & 0 \\ & \ddots & \\ 0 & & J_{\lambda_b} \end{pmatrix}, \text{ where } J_{\lambda_k} = \overbrace{\begin{pmatrix} \lambda_k & 1 & & 0 \\ & \ddots & \ddots & \\ & & \ddots & 1 \\ 0 & & & \lambda_k \end{pmatrix}}^{\mu_k \text{ columns}} \Bigg\} \mu_k \text{ rows}, \ 1 \leq k \leq b.$$

The complex values $\lambda_1, \ldots, \lambda_b$ are the eigenvalues of $\boldsymbol{G}$ which are not necessarily distinct. The sequel of matrices $(\boldsymbol{G}^n)_{n \in \mathbb{N}}$ is then bounded if, and only if, we have first $|\lambda_k| \leq 1$, for $1 \leq k \leq b$, and second $\mu_k = 1$ if $|\lambda_k| = 1$. This condition is therefore exactly that the minimal polynomial of $\boldsymbol{G}$ is a simple von Neumann polynomial. □

A sufficient condition, that is indeed useful in most of the configurations considered in the paper, comes from [8, Proposition 3 and 4].

Proposition 12. *If the characteristic polynomial of $\boldsymbol{G}$ is a simple von Neumann polynomial for any $\xi \in [-\pi, \pi]$, then the lattice Boltzmann scheme characterized by the amplification matrix $\boldsymbol{G}$ is stable.*

We hence see that gaps between necessary and sufficient conditions come from those between minimal and characteristic polynomials.

Notation 13 (minimal and characteristic polynomials)**.** The characteristic polynomial of the matrix $\boldsymbol{G}$ defined in (6) is denoted by $\varphi_3(X; \xi, c, t_v, t_w)$. Moreover, its minimal polynomial is denoted by $\mu(X; \xi, c, t_v, t_w)$.

As the minimal polynomial of a complex matrix is not easy to determine and its degree may vary with ξ, we propose to compute the characteristic polynomial φ_3 of the amplification matrix $\boldsymbol{G}$. Then, if $\varphi_3 \in \mathscr{N}^s$, the scheme is stable thanks to Prop. 12; if $\varphi_3 \notin \mathscr{N}$, the minimal polynomial μ is not a von Neumann polynomial and the scheme is unstable; and if $\varphi_3 \in \mathscr{N} \backslash \mathscr{N}^s$, the case has to be investigated more precisely. Fortunately, the first two cases are generic and the last one is not.

The characteristic polynomial of the amplification matrix $\boldsymbol{G}$ reads

$$\varphi_3(X;\xi,c,t_v,t_w) = \alpha_{3,0} + \alpha_{3,1}X + \alpha_{3,2}X^2 + \alpha_{3,3}X^3, \tag{12}$$

with

$$\begin{aligned}
\alpha_{3,0} &= -t_v t_w,\\
\mathrm{Re}(\alpha_{3,1}) &= t_v t_w - (1-\cos\xi)\, t_v\,(1+t_w)\, c^2 - \cos\xi\,(t_v+t_w),\\
\mathrm{Im}(\alpha_{3,1}) &= c\, t_w\,(1+t_v)\sin\xi,\\
\mathrm{Re}(\alpha_{3,2}) &= -1 + c^2(1+t_w)(1-\cos\xi) + \cos\xi\,(t_v+t_w),\\
\mathrm{Im}(\alpha_{3,2}) &= c(1+t_v)\sin\xi,\\
\alpha_{3,3} &= 1,
\end{aligned}$$

where $\mathrm{Re}(\cdot)$ and $\mathrm{Im}(\cdot)$ refer to the real and imaginary parts of a complex number.

Remark 14 (Corresponding finite difference scheme). As emphasized at the end of Sec. 3, we look at u_j^n as an approximation of the solution of the target PDE. One can therefore aim at recasting the discrete dynamics of u_j^n only in terms of itself. This can be done in full generality using the characteristic polynomial φ_3 from (12). This is a consequence of the Cayley-Hamilton theorem. More precisely,

(13) if $(u_j^n)_{j\in\mathbb{Z}}^{n\in\mathbb{N}}$ fulfills (3) and (4), it also fulfills, for $\xi \in [-\pi,\pi]$

$$\begin{cases}
\hat{u}^1(\xi) = (\boldsymbol{M}\boldsymbol{T}(\xi)\boldsymbol{M}^{-1}\boldsymbol{R}\,\hat{\boldsymbol{m}}^0(\xi))_1,\\
\hat{u}^2(\xi) = ((\boldsymbol{M}\boldsymbol{T}(\xi)\boldsymbol{M}^{-1}\boldsymbol{R})^2\,\hat{\boldsymbol{m}}^0(\xi))_1,\\
\hat{u}^{n+1}(\xi) = -\alpha_{3,2}(\xi)\hat{u}^n(\xi) - \alpha_{3,1}(\xi)\hat{u}^{n-1}(\xi) - \alpha_{3,0}(\xi)\hat{u}^{n-2}(\xi), \quad n \geq 2.
\end{cases}$$

This means that u_j^n also fulfills a multi-step finite difference scheme, whose "amplification polynomial" is φ_3, with specific initialization schemes, linked to the choice of $\boldsymbol{m}^0$, that provide u_j^1 and u_j^2. The stability of the multi-step finite difference scheme, namely the fact that

$$\sum_{j\in\mathbb{Z}} \Delta x |z_j^n|^2 \leq C^2 \Big(\sum_{j\in\mathbb{Z}} \Delta x |z_j^0|^2 + \sum_{j\in\mathbb{Z}} \Delta x |z_j^1|^2 + \sum_{j\in\mathbb{Z}} \Delta x |z_j^2|^2\Big), \quad n \geq 3,$$

where

$$\hat{z}^{n+1}(\xi) = -\alpha_{3,2}(\xi)\hat{z}^n(\xi) - \alpha_{3,1}(\xi)\hat{z}^{n-1}(\xi) - \alpha_{3,0}(\xi)\hat{z}^{n-2}(\xi), \quad n \geq 2,$$

is easy to ascertain.

Contrarily to the lattice Boltzmann setting given in Prop. 12, the stability of the multi-step finite difference scheme is equivalent to $\varphi_3(X;\xi,c,t_v,t_w) \in \mathscr{N}^s$ for all $\xi \in [-\pi,\pi]$ [37, Chapter 4]. Note that the initial data z_j^0, z_j^1, and z_j^2 are arbitrary. In particular, a counter-example is provided in [7], where the lattice Boltzmann scheme is stable, whereas the multi-step finite difference scheme that is obtained using the characteristic polynomial is not.

4.3. **Main result: the stability condition.** In this section, the stability condition is investigated and a necessary and sufficient condition is given.

The first result is usefull to reduce the complexity of the proof. The parameters that ensure the scheme stability have to be searched such that conditions $-1 \leq t_v, t_w \leq 1$ and $|c| \leq 1$ are satisfied (the case $t_v = t_w = -1$ being fully treated).

Proposition 15. *Assuming that the scheme is stable in the sense of Def. 10, first the centered relaxation parameters verify*

$$|t_v| \leq 1, \qquad \textit{and} \qquad |t_w| \leq 1;$$

second the equilibrium value c satisfies

$$|c| \leq 1 \qquad \textit{if} \qquad (t_v, t_w) \neq (-1, -1).$$

Moreover, if $(t_v, t_w) = (-1, -1)$, the scheme is stable.

Proof. First, for $\xi = 0$, we have

$$\varphi_3(X; \xi = 0, c, t_v, t_w) = (X-1)(X+t_v)(X+t_w).$$

If the scheme is stable, the polynomial φ_3 has to be a von Neumann polynomial in particular for $\xi = 0$: the condition $-1 \leq t_v, t_w \leq 1$ is then immediately satisfied.

To prove that the second condition is also a necessary condition for the stability, the characteristic polynomial φ_3 is studied for $\xi = \pi$ and $\xi = \pi/2$. First we assume that $t_w \in (-1, 1]$:

$$\varphi_3(X; \xi = \pi, c, t_v, t_w) = (X - t_v)(X^2 + X(2c^2 - 1)(t_w + 1) + t_w).$$

Assuming then that $|c| > 1$, we note $Q_3^\pi = X^2 + X(2c^2-1)(t_w+1) + t_w$. This polynomial has two real roots as its discriminant Δ is positive:

$$\Delta = (2c^2-1)^2(t_w+1)^2 - 4t_w = 4c^2(c^2-1)(1+t_w)^2 + (1-t_w)^2 > 0.$$

As Q_3^π is convex and $Q_3^\pi(-1) = 2(1-c^2)(1+t_w) < 0$, one of its roots is less than -1. Second, we assume that $t_w = -1$ and $t_v \in (-1, 1]$:

$$\varphi_3(X; \xi = \pi/2, c, t_v, -1) = (X-1)(X^2 + ic(1+t_v)X - t_v).$$

Assuming then that $|c| > 1$, we note $Q_3^{\pi/2} = X^2 + ic(1+t_v)X - t_v$. This polynomial has two purely imaginary roots as its discriminant Δ is negative:

$$\Delta = -c^2(1+t_v)^2 + 4t_v = -(c^2-1)(1+t_v)^2 - (1-t_v)^2 < 0.$$

We define $\tilde{Q}_3^{\pi/2}(X) = Q_3^{\pi/2}(iX) = -X^2 - c(1+t_v)X - t_v$, $\tilde{Q}_3^{\pi/2}$ is concave. If $c > 1$, $\tilde{Q}_3^{\pi/2}(-1) = (1+t_v)(c-1) > 0$, one of its roots is less than -1. If $c < -1$, $\tilde{Q}_3^{\pi/2}(1) = (1+t_v)(-c-1) > 0$, one of its roots is greater than 1. That ends the proof.

The case $t_v = -1$ and $t_w = -1$ is fully degenerated as the matrix $\boldsymbol{G}$ reads

$$\boldsymbol{G} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & e^{-i\xi} & 0 \\ 0 & 0 & e^{i\xi} \end{pmatrix}$$

and its minimal polynomial is clearly a simple von Neumann polynomial for any $\xi \in [-\pi, \pi]$ and for any $c \in \mathbb{R}$. □

The following theorem is the main result of the contribution and gives the necessary and sufficient condition for the D1Q3 scheme to be stable. In order to be more readble for regular users of lattice Boltzmann scheme written in the MRT formalism, the condition is written according to the relaxation parameters s_v, s_w and not the centered ones t_v, t_w.

Theorem 16. *The D1Q3 scheme is stable if, and only if the relaxation parameters verify the following properties according to the value of the velocity c:*

- $0 \leq s_v, s_w \leq 2$ *and* $(s_v, s_w) \notin \{(0,2),(2,2)\}$ *if* $c = 0$;
- $0 \leq s_v < 2$ *and* $0 \leq s_w \leq s_w^\star(s_v)$ *if* $0 < |c| < 1$;
- $0 \leq s_v, s_w \leq 2$ *and* $(s_v, s_w) \notin \{(0,2),(2,0),(2,2)\}$ *if* $|c| = 1$;
- $s_v = s_w = 0$ *if* $|c| > 1$.

The limit value $s_w^\star : [0,2] \to [0,2]$ reads as a bijective decreasing function given by

$$s_w^\star(s) = 1 + \frac{s^2 - 8s + 8}{\sqrt{(s^2 - 8s + 8)^2 + 4s^2(2-s)^2} + 2s(2-s)}. \tag{14}$$

Fig. 1 shows the stable area according to the relaxation parameters s_v and s_w. This area does not depend on the velocity c as long as it verifies $0 < |c| < 1$.

Proof of Thm. 16. The proof simply consists of applying,

- Prop. 32 for $s_v = 0$ and $s_w = 0$, Prop. 33 for $s_v = 2$ and $s_w = 0$, Prop. 34 for $s_v = 0$ and $s_w = 2$, Prop. 35 for $s_v = 2$ and $s_w = 2$;
- Prop. 28 for $0 < s_v < 2$ and $s_w = 0$, Prop. 29 for $s_v = 0$ and $0 < s_w < 2$, Prop. 30 for $0 < s_v < 2$ and $s_w = 2$, Prop. 31 for $s_v = 2$ and $0 < s_w < 2$;
- Prop. 27 for $0 < s_v < 2$ and $0 < s_w < 2$;

keeping in mind that $s_v = t_v + 1$ and $s_w = t_w + 1$. □

Corollary 17. *The single relaxation time scheme obtained by fixing $s_w = s_v$ is stable if the unique relaxation parameter $s = s_v = s_w$ satisfies*

- $0 \leq s \leq s^\star = 6 - 2\sqrt{6} \gtrapprox 1.1010$, *if* $0 < |c| < 1$;
- $0 \leq s < 2$, *if* $|c| \in \{0, 1\}$.

Proof of Cor. 17. The limit relaxation parameter $s^\star$ is defined as the unique solution of the fixed point problem $s = s_w^\star(s)$ on $[0,2]$ or equivalently $s^\star = t^\star + 1$ with $t^\star$ the unique solution of $t = t_w^\star(t)$ on $[-1,1]$. As $t = 3 - 2\sqrt{2}$ (the unique root of $t^2 - 6t + 1$ in $[-1,1]$) is not a fixed point of $t_w^\star$, we have

$$\begin{aligned}
t = t_w^\star(t) &\iff \begin{cases} t(t^2 - 6t + 1) = -2(1 - t^2) + \sqrt{(t^2 - 6t + 1)^2 + 4(1 - t^2)^2} \\ \text{and } -1 \leq t \leq 1 \end{cases} \\
&\iff \begin{cases} (t(t^2 - 6t + 1) + 2(1 - t^2))^2 = (t^2 - 6t + 1)^2 + 4(1 - t^2)^2 \\ \text{and } -1 \leq t \leq 1 \text{ and } t(t^2 - 6t + 1) + 2(1 - t^2) \geq 0 \end{cases} \\
&\iff \begin{cases} 4(1 - t^2)t(t^2 - 6t + 1) = (1 - t^2)(t^2 - 6t + 1)^2 \\ \text{and } -1 \leq t \leq 1 \text{ and } t^3 - 8t^2 + t + 2 \geq 0. \end{cases}
\end{aligned}$$

As both $t = \pm 1$ do not satisfy the inequality $t^3 - 8t^2 + t + 2 \geq 0$, we can simplify the first equality by $1 - t^2$ and also by $t^2 - 6t + 1$ (as $3 - 2\sqrt{2}$ is not a solution)




FIGURE 1. Representation of the stability condition of Thm. 16: green means stable while purple means unstable

that yields

$$t = t_w^\star(t) \iff \left(t^2 - 10t + 1 = 0 \text{ and } -1 \leq t \leq 1 \text{ and } t^3 - 8t^2 + t + 2 \geq 0\right).$$
$$\iff t = 5 - 2\sqrt{6}.$$

That ends the proof. □

Corollary 18. *The lattice Boltzmann scheme introduced in* [17] *with the linked relaxation parameters* $s_w = 2 - s_v$ *is stable if*

- $0 \leq s_v < 2$ *if* $0 < |c| < 1$;
- $0 < s_v < 2$ *if* $c \in \{0, 1\}$.

Proof of Cor. 18. Thanks to Thm. 16, we only have to prove that $2 - s \leq s_w^\star(s)$ for $0 \leq s \leq 2$ or equivalently that $t + t_w^\star(t) \geq 0$ for $-1 \leq t \leq 1$. We first consider the obvious case $t = 3 - 2\sqrt{2} > 0$ with $t_w^\star(t) = 0$. For $t \neq 3 - 2\sqrt{2}$, we rewrite $t + t_w^\star(t)$

as

$$t + t_w^\star(t) = \sqrt{1+x^2} + 1 - x \quad \text{with} \quad x = 2(1-t^2)/(t^2-6t+1).$$

The function $x \mapsto \sqrt{1+x^2}+1-x$ is clearly decreasing and positive on $\mathbb{R}$, what ends the proof. □

5. Proof of the stability condition

This section is devoted to the study of the main theorem. We first restrict the domain in which the parameters (t_v, t_w, c) lie thanks to symmetry properties and we introduce some notations to facilitate a complete proof. The computations were carried out with the help of SageMath [39]. The computer-assisted calculations accompanying this work are available at [**?**].

We recall that we have defined φ_3 as the characteristic polynomial of the amplification matrix $\boldsymbol{G}$ given in (5). The subscript 3 indicates that the degree of this polynomial is 3. Then we note $\varphi_2^\circ = \mathscr{C}(\varphi_3)$ the polynomial obtained after one step of the Cohn-Schur operator and $\varphi_2^\bullet = \mathrm{d}\varphi_3/\mathrm{d}X$ the polynomial obtained after the derivative operator. The subscript 2 indicates that the degree of these polynomials is 2. We continue this process by defining polynomials of degree 1 $\varphi_1^{\circ\circ} = \mathscr{C}(\varphi_2^\circ)$, $\varphi_1^{\circ\bullet} = \mathrm{d}\varphi_2^\circ/\mathrm{d}X$, $\varphi_1^{\bullet\circ} = \mathscr{C}(\varphi_2^\bullet)$, and $\varphi_1^{\bullet\bullet} = \mathrm{d}\varphi_2^\bullet/\mathrm{d}X$. The expression of each of these polynomials is given in the appendix A.

5.1. Decomposition of the studied domain.

Thanks to theorem 15, we assume in this section that

$$|t_v| \le 1, \quad |t_w| \le 1, \quad |c| \le 1.$$

The following property is usefull to reduce the complexity of the proof.

Property 19. *The matrix $\boldsymbol{G}$ defined in (6) depends on the t_v, t_w, ξ, and c variables and satisfies the symmetry properties:*

$$\boldsymbol{G}(\xi,-c,t_v,t_w) = \boldsymbol{P}\overline{\boldsymbol{G}(\xi,c,t_v,t_w)}\boldsymbol{P}, \quad \boldsymbol{G}(-\xi,c,t_v,t_w) = \overline{\boldsymbol{G}(\xi,c,t_v,t_w)}$$

with

$$\boldsymbol{P} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}$$

for $-\pi \le \xi \le \pi$, $-1 \le c \le 1$, and $-1 \le t_v, t_w \le 1$. The characteristic polynomial φ_3 and the minimal polynomial μ satisfy

$$\varphi_3(X;\xi,-c,t_v,t_w) = \overline{\varphi_3(X;\xi,c,t_v,t_w)}, \quad \varphi_3(X;-\xi,c,t_v,t_w) = \overline{\varphi_3(X;\xi,c,t_v,t_w)}$$
$$\mu(X;\xi,-c,t_v,t_w) = \overline{\mu(X;\xi,c,t_v,t_w)}, \qquad \mu(X;-\xi,c,t_v,t_w) = \overline{\mu(X;\xi,c,t_v,t_w)},$$

for $-\pi \le \xi \le \pi$, $-1 \le c \le 1$, and $-1 \le t_v, t_w \le 1$.

We then investigate the stability condition assuming that the parameters lie in the hypercube

$$-1 \le t_v \le 1, \quad -1 \le t_w \le 1, \quad 0 \le c \le 1, \quad 0 \le \xi \le \pi. \tag{15}$$

In order to split the proof, we decompose the square domain

$$\mathcal{D}_{t_v t_w} = [-1,1]\times[-1,1] \tag{16}$$

where the centered relaxation parameters t_v and t_w are taken into its interior $\mathcal{I}_{t_v t_w}$, its open edges $\mathcal{E}_{t_v t_w}$, and its vertices $\mathcal{V}_{t_v t_w}$ (see Fig. 2):

$$\mathcal{D}_{t_v t_w} = \mathcal{I}_{t_v t_w} \cup \mathcal{E}_{t_v t_w} \cup \mathcal{V}_{t_v t_w} \tag{17}$$

where

$$\mathcal{I}_{t_v t_w} = (-1,1)\times(-1,1),$$

$$\begin{aligned}
&\mathcal{E}_{t_v t_w} = \mathcal{E}^b_{t_v t_w} \cup \mathcal{E}^t_{t_v t_w} \cup \mathcal{E}^l_{t_v t_w} \cup \mathcal{E}^r_{t_v t_w}, && \mathcal{V}_{t_v t_w} = \mathcal{V}^{lb}_{t_v t_w} \cup \mathcal{V}^{rb}_{t_v t_w} \cup \mathcal{V}^{lt}_{t_v t_w} \cup \mathcal{V}^{rt}_{t_v t_w},\\
&\mathcal{E}^b_{t_v t_w} = (-1,1)\times\{-1\}, && \mathcal{V}^{lb}_{t_v t_w} = \{-1\}\times\{-1\},\\
&\mathcal{E}^t_{t_v t_w} = (-1,1)\times\{+1\}, && \mathcal{V}^{rb}_{t_v t_w} = \{+1\}\times\{-1\},\\
&\mathcal{E}^l_{t_v t_w} = \{-1\}\times(-1,1), && \mathcal{V}^{lt}_{t_v t_w} = \{-1\}\times\{+1\},\\
&\mathcal{E}^r_{t_v t_w} = \{+1\}\times(-1,1), && \mathcal{V}^{rt}_{t_v t_w} = \{+1\}\times\{+1\}.
\end{aligned}$$




FIGURE 2. Decomposition of the square $\mathcal{D}_{t_v t_w} = [-1,1]\times[-1,1]$.

We also decompose the rectangular domain

$$\mathcal{D}_{c\xi} = [0,1]\times[0,\pi] \tag{18}$$

where the velocity c and the wave number ξ are taken into its interior $\mathcal{I}_{c\xi}$, its open edges $\mathcal{E}_{c\xi}$, and its vertices $\mathcal{V}_{c\xi}$ (see Fig. 3):

$$\mathcal{D}_{c\xi} = \mathcal{I}_{c\xi} \cup \mathcal{E}_{c\xi} \cup \mathcal{V}_{c\xi} \tag{19}$$

where

$$\mathcal{I}_{c\xi} = (0,1)\times(0,\pi),$$

$$\begin{aligned}
&\mathcal{E}_{c\xi} = \mathcal{E}^b_{c\xi} \cup \mathcal{E}^t_{c\xi} \cup \mathcal{E}^l_{c\xi} \cup \mathcal{E}^r_{c\xi}, && \mathcal{V}_{c\xi} = \mathcal{V}^{lb}_{c\xi} \cup \mathcal{V}^{rb}_{c\xi} \cup \mathcal{V}^{lt}_{c\xi} \cup \mathcal{V}^{rt}_{c\xi},\\
&\mathcal{E}^b_{c\xi} = (0,1)\times\{0\}, && \mathcal{V}^{lb}_{c\xi} = \{0\}\times\{0\},\\
&\mathcal{E}^t_{c\xi} = (0,1)\times\{\pi\}, && \mathcal{V}^{rb}_{c\xi} = \{1\}\times\{0\},\\
&\mathcal{E}^l_{c\xi} = \{0\}\times(0,\pi), && \mathcal{V}^{lt}_{c\xi} = \{0\}\times\{\pi\},\\
&\mathcal{E}^r_{c\xi} = \{1\}\times(0,\pi), && \mathcal{V}^{rt}_{c\xi} = \{1\}\times\{\pi\}.
\end{aligned}$$




FIGURE 3. Decomposition of the square $\mathcal{D}_{c\xi} = [0,1]\times[0,\pi]$.

5.2. **Technical lemmas.** In this section, we prove some technical lemmas that are used in the sequel to investigate the stability area. In the first two lemmas, we characterize the membership properties of the polynomial φ_3 in the sets $\mathscr{I}$ and $\mathscr{E}$.

Lemma 20. *For $(t_v,t_w)\in\mathcal{D}_{t_vt_w}$ and $(c,\xi)\in\mathcal{D}_{c\xi}$, we have*

$$\varphi_3(\cdot;\xi,c,t_v,t_w)\in\mathscr{I} \iff (t_v,t_w)\in\mathcal{I}_{t_vt_w}\cup\mathcal{E}_{t_vt_w}.$$

Proof of lemma 20. $\varphi_3\in\mathscr{I}$ if, and only if, $|\alpha_{3,0}|<|\alpha_{3,3}|$ that reads $|t_vt_w|<1$. □

Lemma 21. *For $(t_v,t_w)\in\mathcal{D}_{t_vt_w}$ and $(c,\xi)\in\mathcal{D}_{c\xi}$, we have*

$$\varphi_3(\cdot;\xi,c,t_v,t_w)\in\mathscr{E} \iff (t_v,t_w)\in\mathcal{V}_{t_vt_w}.$$

Proof of lemma 21. By lemma 20, and since the intersection $\mathscr{I}\cap\mathscr{E}$ is empty, it follows that

$$\varphi_3(\cdot;\xi,c,t_v,t_w)\in\mathscr{E} \implies (t_v,t_w)\in\mathcal{V}_{t_vt_w}.$$

Denoting $\varphi_2^\circ=\mathscr{C}(\varphi_3)$, we have $\varphi_2^\circ(\cdot;\xi,c,t_v,t_w)\equiv 0$ if, and only if,

$$\alpha_{3,0}\,\overline{\alpha}_{3,0}=\alpha_{3,3}\,\overline{\alpha}_{3,3},\quad \overline{\alpha}_{3,3}\,\alpha_{3,1}=\alpha_{3,0}\,\overline{\alpha}_{3,2}\quad\text{and}\quad \overline{\alpha}_{3,3}\,\alpha_{3,2}=\alpha_{3,0}\,\overline{\alpha}_{3,1}.$$

It is obvious that two of these three relations imply the third. The previous equalities yield as $\alpha_{3,3}=1$

$$\alpha_{3,0}\,\overline{\alpha}_{3,0}=1\quad\text{and}\quad \alpha_{3,1}=\alpha_{3,0}\,\overline{\alpha}_{3,2}.$$

The condition $|\alpha_{3,0}|=1$ reads $|t_vt_w|=1$. First, we assume that $t_vt_w=1$. As $(t_v,t_w)\in[-1,1]^2$, we have $t_v=t_w$ and

$$\alpha_{3,1}-\alpha_{3,0}\,\overline{\alpha}_{3,2}=c(t_w-t_v)\big(c(1-\cos\xi)+i\sin\xi\big)=0.$$

Second, we assume that $t_vt_w=-1$. As $(t_v,t_w)\in[-1,1]^2$, we have $t_v=-t_w$ and

$$\alpha_{3,1}-\alpha_{3,0}\,\overline{\alpha}_{3,2}=-(t_w+t_v)\big(c^2+(2-c^2)\cos\xi-ic\sin\xi\big)=0.$$

That ends the proof. □

The next two lemmas are dedicated to the characterization of membership for the polynomial $\varphi_2^\circ=\mathscr{C}(\varphi_3)$ obtained after one step of the Cohn-Schur operator.

Lemma 22. *For $(t_v,t_w)\in\mathcal{I}_{t_vt_w}\cup\mathcal{E}_{t_vt_w}$ and $(c,\xi)\in\mathcal{D}_{c\xi}$, we have*

$$\varphi_2^\circ(\cdot;\xi,c,t_v,t_w)\in\mathscr{I}\iff\begin{cases}(t_v,t_w)\in\mathcal{I}_{t_vt_w},\\ \text{or}\\ (t_v,t_w)\in\mathcal{E}^l_{t_vt_w}\cup\mathcal{E}^r_{t_vt_w}\ \text{and}\ (c,\xi)\in\mathcal{I}_{c\xi}\cup\mathcal{E}^l_{c\xi}\cup\mathcal{E}^t_{c\xi},\\ \text{or}\\ (t_v,t_w)\in\mathcal{E}^b_{t_vt_w}\cup\mathcal{E}^t_{t_vt_w}\ \text{and}\ (c,\xi)\in\mathcal{I}_{c\xi}\cup\mathcal{E}^l_{c\xi}.\end{cases}$$

Proof of lemma 22. The polynomial $\varphi_2^\circ=\mathscr{C}(\varphi_3)$ is given in (25). Then, the condition $\varphi_2^\circ(\cdot;\xi,c,t_v,t_w)\in\mathscr{I}$ reads

$$\psi_2^\circ(\xi,c,t_v,t_w)=\beta^\circ_{2,0}+\beta^\circ_{2,1}c^2+\beta^\circ_{2,2}c^4>0,$$

with

$$\begin{aligned}\beta^\circ_{2,0}&=(1-t_vt_w)^2\big((1+t_vt_w)^2-\cos^2(\xi)(t_v+t_w)^2\big),\\ \beta^\circ_{2,1}&=(\cos\xi-1)\big(t_w^2(1-t_v^2)^2+\cos\xi\big(t_v^4t_w^2+2t_v^3t_w^3+2t_v^2t_w^4-2t_v^3t_w-6t_v^2t_w^2\\ &\qquad\qquad-2t_vt_w^3+2t_v^2+2t_vt_w+t_w^2\big)\big),\\ \beta^\circ_{2,2}&=-(1-t_w^2)^2t_v^2(1-\cos\xi)^2.\end{aligned}$$

First, we consider $(t_v,t_w)\in\mathcal{I}_{t_vt_w}$, that is $|t_v|<1$ and $|t_w|<1$. We observe that ψ_2° can be seen as a polynomial of degree 2 in the indeterminate c^2. As $\beta^\circ_{2,2}\le0$, this polynomial is concave. We then compute its value in $c^2=0$ and $c^2=1$:

$$\begin{aligned}\psi_2^\circ(\xi,0,t_v,t_w)&=\beta^\circ_{2,0}=(1-t_vt_w)^2\big((1-t_v^2)(1-t_w^2)+(1-\cos^2(\xi))(t_v+t_w)^2\big),\\ \psi_2^\circ(\xi,1,t_v,t_w)&=\beta^\circ_{2,0}+\beta^\circ_{2,1}+\beta^\circ_{2,2}\\ &=(1-t_v^2)(1-t_w^2)\big((1-\cos\xi t_vt_w)^2+(1-\cos^2(\xi))t_v^2t_w^2\big).\end{aligned}$$

We have $\psi_2^\circ(\xi,0,t_v,t_w)>0$ and $\psi_2^\circ(\xi,1,t_v,t_w)>0$, the condition $\psi_2^\circ(\xi,c,t_v,t_w)>0$ is then satisfied for $(c,\xi)\in\mathcal{D}_{c\xi}$.

Second, we consider $(t_v,t_w)\in\mathcal{E}^l_{t_vt_w}\cup\mathcal{E}^r_{t_vt_w}$, that is $|t_v|=1$ and $|t_w|<1$. We have

$$\psi_2^\circ(\xi,c,\pm1,t_w)=(1-\cos\xi)(1-c^2)(1-t_w^2)^2(1+\cos\xi+c^2(1-\cos\xi)).$$

The condition $\psi_2^\circ(\xi,c,\pm1,t_w)>0$ is then satisfied for $(c,\xi)\in\mathcal{I}_{c\xi}\cup\mathcal{E}^l_{c\xi}\cup\mathcal{E}^t_{c\xi}$.

Third, we consider $(t_v,t_w)\in\mathcal{E}^b_{t_vt_w}\cup\mathcal{E}^t_{t_vt_w}$, that is $|t_v|<1$ and $|t_w|=1$. We have

$$\psi_2^\circ(\xi,c,t_v,\pm1)=(1-\cos^2\xi)(1-c^2)(1-t_v^2)^2.$$

The condition $\psi_2^\circ(\xi,c,t_v,\pm1)>0$ is then satisfied for $(c,\xi)\in\mathcal{I}_{c\xi}\cup\mathcal{E}^l_{c\xi}$. □

Lemma 23. *For $(t_v,t_w)\in\mathcal{I}_{t_vt_w}\cup\mathcal{E}_{t_vt_w}$ and $(c,\xi)\in\mathcal{D}_{c\xi}$, we have*

$$\varphi_2^\circ(\cdot;\xi,c,t_v,t_w)\in\mathscr{E}\iff\begin{cases}(t_v,t_w)\in\mathcal{E}^l_{t_vt_w}\cup\mathcal{E}^r_{t_vt_w}\\ \qquad\text{and}\ (c,\xi)\in\mathcal{E}^b_{c\xi}\cup\mathcal{E}^r_{c\xi}\cup\mathcal{V}_{c\xi},\\ \text{or}\\ (t_v,t_w)\in\mathcal{E}^b_{t_vt_w}\cup\mathcal{E}^t_{t_vt_w}\\ \qquad\text{and}\ (c,\xi)\in\mathcal{E}^b_{c\xi}\cup\mathcal{E}^r_{c\xi}\cup\mathcal{E}^t_{c\xi}\cup\mathcal{V}_{c\xi}.\end{cases}$$

Proof of lemma 23. We first reduce the investigation to $|t_v| = 1$ or $|t_w| = 1$. Indeed, as the intersection $\mathscr{I} \cap \mathscr{E}$ is empty, we necessarely have

$$\varphi_2^\circ(\cdot; \xi, c, t_v, t_w) \in \mathscr{E} \implies (t_v, t_w) \in \mathcal{E}_{t_v t_w} \cup \mathcal{V}_{t_v t_w}$$

thanks to the previous lemma 22. Note that we assume $|t_v t_w| < 1$, then both equalities $|t_v| = 1$ and $|t_w| = 1$ are not allowed simultaneously in this proof. The expression of the polynomial $\varphi_1^{\circ\circ}$ is given in (27).

Considering $t_v = -1$, the coefficients $\alpha_{1,0}^{\circ\circ}$ and $\alpha_{1,1}^{\circ\circ}$ admit these factorizations:

$$\alpha_{1,0}^{\circ\circ} = (1-\cos\xi)(1-c^2)(1-t_w^2)^2(c^2(1-\cos\xi) - (1+\cos\xi)),$$
$$\alpha_{1,1}^{\circ\circ} = (1-\cos\xi)(1-c^2)(1-t_w^2)^2(c^2(1-\cos\xi) + (1+\cos\xi)).$$

As $|t_w| < 1$, the equalities $\alpha_{1,0}^{\circ\circ} = \alpha_{1,1}^{\circ\circ} = 0$ are then equivalent to

$$\cos\xi = 1 \text{ or } c^2 = 1 \text{ or } (c = 0 \text{ and } \cos\xi = -1).$$

Considering $t_v = 1$, the coefficients $\alpha_{1,0}^{\circ\circ}$ and $\alpha_{1,1}^{\circ\circ}$ admit these factorizations:

$$\alpha_{1,1}^{\circ\circ} = -\operatorname{Re}(\alpha_{1,0}^{\circ\circ}) = (1-\cos\xi)(1-c^2)(1-t_w^2)^2(1+\cos\xi+c^2(1-\cos\xi)),$$
$$\operatorname{Im}(\alpha_{1,0}^{\circ\circ}) = 2\sin\xi(1-\cos\xi)c(1-c^2)(1-t_w^2)^2.$$

As $|t_w| < 1$, the equalities $\alpha_{1,0}^{\circ\circ} = \alpha_{1,1}^{\circ\circ} = 0$ are then equivalent to

$$\cos\xi = 1 \text{ or } c^2 = 1 \text{ or } (c = 0 \text{ and } \cos\xi = -1).$$

Considering now $t_w = -1$, the coefficients $\alpha_{1,0}^{\circ\circ}$ and $\alpha_{1,1}^{\circ\circ}$ admit these factorizations:

$$\alpha_{1,1}^{\circ\circ} = -\alpha_{1,0}^{\circ\circ} = (1-\cos^2\xi)(1-c^2)(1-t_v^2)^2.$$

As $|t_v| < 1$, the equalities $\alpha_{1,0}^{\circ\circ} = \alpha_{1,1}^{\circ\circ} = 0$ are then equivalent to

$$\cos^2\xi = 1 \text{ or } c^2 = 1.$$

Considering finally $t_w = 1$, the coefficients $\alpha_{1,0}^{\circ\circ}$ and $\alpha_{1,1}^{\circ\circ}$ admit these factorizations:

$$\alpha_{1,1}^{\circ\circ} = -\operatorname{Re}(\alpha_{1,0}^{\circ\circ}) = (1-\cos^2\xi)(1-c^2)(1-t_v^2)^2,$$
$$\operatorname{Im}(\alpha_{1,0}^{\circ\circ}) = 2\sin\xi(1-\cos\xi)c(1-c^2)(1-t_v^2)^2.$$

As $|t_v| < 1$, the equalities $\alpha_{1,0}^{\circ\circ} = \alpha_{1,1}^{\circ\circ} = 0$ are then equivalent to $\cos^2\xi = 1$ or $c^2 = 1$. □

5.3. **Stability on $\mathcal{I}_{t_v t_w}$.** In this section, we consider $(t_v, t_w) \in \mathcal{I}_{t_v t_w}$, that is $|t_v| < 1$ and $|t_w| < 1$. We prove in Prop. 27 that (see Fig. 4):

- for $|t_v| < 1$ and $|t_w| < 1$, the scheme is stable for $c \in \{0, 1\}$;
- for $|t_v| < 1$, $|t_w| < 1$, and $0 < c < 1$, the scheme is stable if, and only if $-1 < t_w \leq t_w^\star(t_v)$, the function $t_w^\star$ being defined in the sequel.

We first give some technical results used to describe the stability area. We introduce a polynomial of degre two of the variables t_v and t_w and we investigate its sign. We will see in the sequel that this sign precisely describes the stability area restricted to the open square $(t_v, t_w) \in \mathcal{I}_{t_v t_w}$. Let's be

$$\Upsilon(t, s) = -t^2 s^2 + 6ts^2 + 4t^2 s - s^2 + t^2 - 4s - 6t + 1.$$




FIGURE 4. Stability for $(t_v, t_w) \in \mathcal{I}_{t_v t_w}$. Green means stable while purple means unstable.




FIGURE 5. Representation of the functions $t_w^\star$ and $t_v^\star$

5.3.1. *Parametrization according to t_v.* In this first paragraph, we consider Υ as an element of $\mathbb{R}[t][s]$. We introduce two polynomial functions $\kappa, \eta : [-1, 1] \to \mathbb{R}$ by

$$\kappa(t) = 1 - 6t + t^2, \qquad \eta(t) = 2(1 - t^2),$$

in order to have

$$\Upsilon(t, s) = \kappa(t)(1 - s^2) - 2\eta(t)s.$$

We immediately have the following table of signs

t	-1		$3 - 2\sqrt{2}$		1
$\kappa(t)$	⋮	$+$	0	$-$	⋮
$\eta(t)$	0	$+$	⋮	$+$	0

We then define two functions that are the roots of Υ parametrized by the first variable t. The first one $t_w^\star : [-1, 1] \to \mathbb{R}$ and the second one $t_w^\diamond : [-1, 1] \backslash \{3 - 2\sqrt{2}\} \to \mathbb{R}$ read

$$t_w^\star(t) = \frac{\kappa(t)}{\sqrt{\eta(t)^2 + \kappa(t)^2} + \eta(t)}, \quad t_w^\diamond(t) = -\frac{\sqrt{\eta(t)^2 + \kappa(t)^2} + \eta(t)}{\kappa(t)}. \tag{20}$$

We have the following table of variations proved in Prop. 24.

t	-1		$3-2\sqrt{2}$		1
$t_w^\star(t)$	1	$\searrow$	0	$\searrow$	-1
$t_w^\diamond(t)$	-1	$\searrow$	$-\infty \,\|\, +\infty$	$\searrow$	1

Proposition 24. *The functions $t_w^\star$ and $t_w^\diamond$ defined in (20) satisfy the three following properties:*

- *the function $t_w^\star$ is decreasing and bijective from $[-1,1]$ into $[-1,1]$;*
- *the function $t_w^\diamond$ is decreasing and bijective from $[-1,3-\sqrt{2}[$ into $]-\infty,-1]$;*
- *the function $t_w^\diamond$ is decreasing and bijective from $]3-\sqrt{2},1]$ into $[1,+\infty[$.*

Proof of Prop. 24. As $\eta(-1)=\eta(1)=0$, $\kappa(-1)>0$, and $\kappa(1)<0$, we first have

$$t_w^\star(-1)=1, \quad t_w^\star(1)=-1, \quad t_w^\diamond(-1)=-1, \quad t_w^\diamond(1)=1.$$

For $t\in(-1,1)$, the function $t_w^\star$ reads

$$t_w^\star(t)=\varphi\Big(\frac{\kappa}{\eta}(t)\Big) \qquad \text{with } \varphi(s)=\frac{s}{\sqrt{1+s^2}+1}.$$

The function φ is defined for $s\in\mathbb{R}$ and verifies

$$\varphi'(s)=\frac{1}{\sqrt{1+s^2}(1+\sqrt{1+s^2})}>0 \quad s\in\mathbb{R},$$

the function φ is then an increasing function. Moreover we have

$$\left(\frac{\kappa}{\eta}\right)'(t)=\frac{\kappa'(t)\eta(t)-\kappa(t)\eta'(t)}{\eta^2(t)}=-\frac{(3t-1)^2+8}{3(1-t^2)^2}<0,$$

the function κ/η is then a decreasing function, what proves the first propertie. The second and third properties are immediately obtained after remarking that $t_w^\diamond(t)=-1/t_w^\star(t)$. □

For a given $t\in[-1,1]$, the roots of the polynomial $\Upsilon(t,s)$, viewed as a quadratic polynomial in s with coefficients that are themselves quadratic polynomials in t, are exactly $t_w^\star(t)$ and $t_w^\diamond(t)$ (for $t=3-2\sqrt{2}$, the only existing root is $t_w^\star(t)$).

5.3.2. *Parametrization according to t_w.* In this second paragraph, we consider Υ as an element of $\mathbb{R}[s][t]$. We introduce two polynomial functions $\mu,\nu:[-1,1]\to\mathbb{R}$ by

$$\mu(s)=1+4s-s^2, \qquad \nu(s)=3(1-s^2),$$

in order to have

$$\Upsilon(t,s)=\mu(s)t^2-2\nu(s)t+\mu(-s).$$

We immediately have the following table of signs

s	-1		$2-\sqrt{5}$		$\sqrt{5}-2$		1
$\mu(s)$	⋮	$-$	0	$+$	⋮	$+$	⋮
$\mu(-s)$	⋮	$+$	⋮	$+$	0	$-$	⋮
$\nu(s)$	0	$+$	⋮	$+$	⋮	$+$	0

We then define two functions that are the roots of Υ parametrized by the second variable s. The first one $t_v^\star : [-1,1] \to \mathbb{R}$ and the second one $t_v^\diamond : [-1,1]\backslash\{2-\sqrt{5}\} \to \mathbb{R}$ read

$$t_v^\star(s) = \frac{\mu(-s)}{\sqrt{\nu(s)^2-\mu(s)\mu(-s)}+\nu(s)}, \quad t_v^\diamond(t) = \frac{\sqrt{\nu(s)^2-\mu(s)\mu(-s)}+\nu(s)}{\mu(s)}. \tag{21}$$

We have the following table of variations proved in Prop. 25.

t	-1		$\sqrt{5}-2$		1
$t_v^\star(s)$	1	$\searrow$	0	$\searrow$	-1
$t_v^\diamond(s)$	-1	$\searrow$	$-\infty$ ‖ $+\infty$	$\searrow$	1

Proposition 25. *The functions $t_v^\star$ and $t_v^\diamond$ defined in (21) satisfy the three following properties:*

- *the function $t_v^\star$ is decreasing and bijective from $[-1,1]$ into $[-1,1]$;*
- *the function $t_v^\diamond$ is decreasing and bijective from $[-1,\sqrt{5}-2[$ into $]-\infty,-1]$;*
- *the function $t_v^\diamond$ is decreasing and bijective from $]\sqrt{5}-2,1]$ into $[1,+\infty[$.*

Proof of Prop. 25. As $\nu(-1)=\nu(1)=0$, $\mu(-1)=-4$, and $\mu(1)=4$, we first have

$$t_v^\star(-1)=1, \quad t_v^\star(1)=-1, \quad t_v^\diamond(-1)=-1, \quad t_v^\diamond(1)=1.$$

For $s\in(-1,1)$, we have

$$t_v^{\star\prime}(s) = -\frac{12(1-s+s^2)}{(\sqrt{8+8s^4}+3(1-s^2))^2} - \frac{16(1-s^2)(2-s+2s^2)}{(\sqrt{8+8s^4}+3(1-s^2))^2\sqrt{8+8s^4}} < 0,$$

what proves the first property. The second and third properties are immediately obtained after remarking that $t_v^\diamond(s) = 1/t_v^\star(-s)$. □

5.3.3. *Description of the stability area in $\mathcal{I}_{t_v t_w}$.* The following proposition concerns the caracterization of the sign of $\Upsilon(t,s)$.

Proposition 26. *The function $\Upsilon : [-1,1]\times[-1,1] \to \mathbb{R}$ defined by*

$$\Upsilon(t,s) = -t^2s^2 + 6ts^2 + 4t^2s - s^2 + t^2 - 4s - 6t + 1,$$

satisfies the two properties

$$\Upsilon(t,s) \geq 0 \iff -1 \leq s \leq t_w^\star(t) \iff -1 \leq t \leq t_v^\star(s),$$
$$\Upsilon(t,s) > 0 \iff -1 \leq s < t_w^\star(t) \iff -1 \leq t < t_v^\star(s).$$

Proof of Prop. 26. A direct investigation of the quadratic polynomial $\Upsilon_t : s \to \Upsilon(t,s)$ for $t\in[-1,1]$ yields to three properties:

- Υ_t is strictly concave for $t\in[-1,3-2\sqrt{2}[$, linear decreasing for $t=3-2\sqrt{2}$, and strictly convex for $t\in]3-2\sqrt{2},1]$;
- if $t\neq 3-2\sqrt{2}$, the two roots of Υ_t are $t_w^\star(t)$ and $t_w^\diamond(t)$;
- if $t=3-2\sqrt{2}$, the only root of Υ_t is $t_w^\star(t)=0$.

Combining these properties, we obtain first that, for $-1\leq t<3-2\sqrt{2}$, $\Upsilon_t(s)\geq 0$ if, and only if, $t_w^\diamond(t)\leq s\leq t_w^\star(t)$, second that, for $t=3-2\sqrt{2}$, $\Upsilon_t(s)\geq 0$ if, and only if, $s\leq t_w^\star(t)$, and third that, for $3-2\sqrt{2}<t\leq 1$, $\Upsilon_t(s)\geq 0$ if, and only if, $s\leq t_w^\star(t)$ or $s\geq t_w^\diamond(t)$. The table of variations of $t_w^\star$ and $t_w^\diamond$ finally ends the proof for the inequalities between s and $t_w^\star(t)$. Composing these inequalities with the decreasing function $t_v^\star$ yields to the second ones as $t_v^\star\circ t_w^\star$ is the identity function of $[-1,1]$. □

Proposition 27 (stability on $\mathcal{I}_{t_v t_w}$)**.** *Assuming $t_v \in (-1,1)$ and $t_w \in (-1,1)$, we have*

- *for $c \in \{0,1\}$, the lattice Boltzmann scheme is stable;*
- *for $0 < c < 1$, the lattice Boltzmann scheme is stable if, and only if the centered relaxation parameters satisfy $-1 < t_w \leq t_w^\star(t_v)$ (or the equivalent relation $-1 < t_v \leq t_v^\star(t_w)$).*

Proof of Prop. 27. In this proof, we assume that $|t_v| < 1$ and $|t_w| < 1$, Lemma 20 implies that $\varphi_3(\cdot;\xi,c,t_v,t_w) \in \mathscr{I}$ and Lemma 22 implies that $\varphi_2^\circ(\cdot;\xi,c,t_v,t_w) \in \mathscr{I}$. Then Theorems 7 and 8 reduce to

$$\varphi_3(\cdot;\xi,c,t_v,t_w) \in \mathscr{N}^s \iff \varphi_2^\circ(\cdot;\xi,c,t_v,t_w) \in \mathscr{N}^s \iff \varphi_1^{\circ\circ}(\cdot;\xi,c,t_v,t_w) \in \mathscr{N}^s,$$
$$\varphi_3(\cdot;\xi,c,t_v,t_w) \in \mathscr{N} \iff \varphi_2^\circ(\cdot;\xi,c,t_v,t_w) \in \mathscr{N} \iff \varphi_1^{\circ\circ}(\cdot;\xi,c,t_v,t_w) \in \mathscr{N}.$$

As $\varphi_1^{\circ\circ}$ is a polynomial of degree one, being a simple von Neumann polynomial or a von Neumann polynomial is exactly the same property. Then we have proven that

$$\varphi_3(\cdot;\xi,c,t_v,t_w) \in \mathscr{N}^s \iff \varphi_3(\cdot;\xi,c,t_v,t_w) \in \mathscr{N} \iff \varphi_1^{\circ\circ}(\cdot;\xi,c,t_v,t_w) \in \mathscr{N},$$
$$(t_v,t_w) \in \mathcal{I}_{t_v t_w},\ (c,\xi) \in \mathcal{D}_{c\xi}.$$

We now investigate the property $\varphi_1^{\circ\circ}(\cdot;\xi,c,t_v,t_w) \in \mathscr{N}$. The expression of the polynomial $\varphi_1^{\circ\circ}$ is given in (27). We then evaluate $\psi_1^{\circ\circ}(\xi,c,t_v,t_w) = |\alpha_{1,1}^{\circ\circ}|^2 - |\alpha_{1,0}^{\circ\circ}|^2$ and $\varphi_1^{\circ\circ} \in \mathscr{N}$ if, and only if, $\psi_1^{\circ\circ} \geq 0$. This polynomial admits the factorization

$$\psi_1^{\circ\circ}(\xi,c,t_v,t_w) = c^2(1-c^2)(1+t_w)(1-t_v^2t_w^2)(1-\cos\xi)^2\theta_1^{\circ\circ}(\xi,c,t_v,t_w),$$

where $\theta_1^{\circ\circ}$ is a polynomial of degree two according to the variable $\cos\xi$. It reads

$$\theta_1^{\circ\circ}(\xi,c,t_v,t_w) = \alpha_0(1-\cos^2\xi) + \tfrac{1}{2}\alpha_-(1-\cos\xi) + \tfrac{1}{2}\alpha_+(1+\cos\xi),$$

with

$$\begin{aligned}
\alpha_0 &= (c^2t_v^2(1-t_w^2)^2 - (t_v+t_w)^2(1-t_vt_w)^2) \\
&\qquad ((1+t_w)(1-t_v^2)(1-c^2) + 4t_v(1-t_w)), \\
\alpha_+ &= (1-t_w)(1-t_v^2)(1-t_vt_w)^2\Upsilon(t_v,t_w), \\
\alpha_- &= (1+t_w)(1-t_v^2)(1-t_w)^2((1-t_v)(1-t_vt_w) + 2c^2t_v(1+t_w))^2,
\end{aligned}$$

First, we obtain that, for $c \in \{0,1\}$, we have $\psi_1^{\circ\circ}(\xi,c,t_v,t_w) = 0$ and that the scheme is stable. Second, for $0 < c < 1$, we have

$$\psi_1^{\circ\circ}(\xi,c,t_v,t_w) \geq 0 \iff \xi = 0 \text{ or } \theta_1^{\circ\circ}(\xi,c,t_v,t_w) \geq 0,$$

as $|t_v| < 1$ and $|t_w| < 1$. Observing that the target condition $-1 < t_w \leq t_w^\star(t_v)$ does not depend on c, we investigate the sign of $\theta_1^{\circ\circ}(\xi,c,t_v,t_w)$ first in the case $c = 0$ even if this case is already covered for the stability. For $c = 0$, $\theta_1^{\circ\circ}(\xi,0,t_v,t_w)$ reads

$$\theta_1^{\circ\circ}(\xi,0,t_v,t_w) = (1-t_vt_w)^2(\overline{\gamma}_{1,0}^{\circ\circ} + \overline{\gamma}_{1,1}^{\circ\circ}\cos\xi)(\gamma_{1,0}^{\circ\circ} + \gamma_{1,1}^{\circ\circ}\cos\xi),$$

where $\overline{\gamma}_{1,0}^{\circ\circ}$, $\overline{\gamma}_{1,1}^{\circ\circ}$, $\gamma_{1,0}^{\circ\circ}$, and $\gamma_{1,1}^{\circ\circ}$ only depend on t_v and t_w. The term $\overline{\gamma}_{1,0}^{\circ\circ} + \overline{\gamma}_{1,1}^{\circ\circ}\cos\xi$ is positive for $\xi \in [0,\pi]$ by linearity as

$$\overline{\gamma}_{1,0}^{\circ\circ} - \overline{\gamma}_{1,1}^{\circ\circ} = (1+t_v)(1+t_w) > 0, \qquad \overline{\gamma}_{1,0}^{\circ\circ} + \overline{\gamma}_{1,1}^{\circ\circ} = (1-t_v)(1-t_w) > 0.$$

Moreover, we have

$$\gamma_{1,0}^{\circ\circ} - \gamma_{1,1}^{\circ\circ} = (1-t_v)^3(1-t_w)^2 > 0, \qquad \gamma_{1,0}^{\circ\circ} + \gamma_{1,1}^{\circ\circ} = (1+t_v)\Upsilon(t_v,t_w),$$

where the function Υ is the one defined in Prop. 26. As an intermediate result, we obtain for $(t_v, t_w) \in \mathcal{I}_{t_v t_w}$ that

$$\forall \xi \in [0,\pi]\ \theta_1^{\circ\circ}(\xi, 0, t_v, t_w) \geq 0 \iff \Upsilon(t_v, t_w) \geq 0 \iff -1 < t_w \leq t_w^\star(t_v).$$

Finally, we have to prove that the result remains the same for all the values of $c \in [0,1]$.

We begin the proof by assuming that $t_w < t_w^\star(t_v)$. We use the fact that the function $\theta_1^{\circ\circ}(\xi, c, t_v, t_w)$ is a polynomial of order 2 in the variable $\cos\xi$ which coefficients are polynomials of order 2 in the variable c^2. For $c = 0$, $\theta_1^{\circ\circ}(\xi, 0, t_v, t_w)$ has two real roots in $(-\infty, -1) \cup (1, +\infty)$. We prove that the discriminant of $\theta_1^{\circ\circ}(\xi, c, t_v, t_w)$ is non negative for $0 \leq c \leq 1$ and that $\theta_1^{\circ\circ}(0, c, t_v, t_w) > 0$ and $\theta_1^{\circ\circ}(\pi, c, t_v, t_w) > 0$ for $0 \leq c \leq 1$. So it implies that, being given a value $0 \leq c \leq 1$, the two roots of $\theta_1^{\circ\circ}(\xi, c, t_v, t_w)$ are real and remain in $(-\infty, -1) \cup (1, +\infty)$ by using the continuity. First, asuming $-1 < t_v < 1$ and $-1 < t_w < t_w^\star(t_v)$,

$$\begin{aligned}
\theta_1^{\circ\circ}(0, c, t_v, t_w) &= (1 - t_w)(1 - t_v^2)(1 - t_v t_w)^2 \Upsilon(t_v, t_w) > 0,\\
\theta_1^{\circ\circ}(\pi, c, t_v, t_w) &= (1 - t_v^2)(1 + t_w)(1 - t_w)^2\\
&\qquad ((1 - t_v)(1 - t_v t_w)(1 - c^2) + (1 + t_v)(1 + t_v t_w)c^2)^2 > 0.
\end{aligned}$$

Second, the discriminant Δ of $\theta_1^{\circ\circ}(\xi, c, t_v, t_w)$ reads $\Delta = \Delta_1^2 \Delta_2$ with

$$\begin{aligned}
\Delta_1 &= 2(t_v^3 t_w + 2t_v^2 t_w^2 - t_v^2 t_w + t_v t_w - 2t_v - t_w),\\
\Delta_2 &= \delta_{20} + \delta_{22} c^2 + \delta_{24} c^4.
\end{aligned}$$

We then prove that $\Delta_2 \geq 0$ in order to conclude that $\Delta \geq 0$. We investigate the sign of Δ_2 as a polynomial of degre 2 in the variable c^2. We observe that the value of $\Delta_2(c^2)$ is bounded by the chord at $c^2 = 0$ and $c^2 = 1$ and the two tangents to the curve at the same points. That is

$$\Delta_2(c^2) \geq \min\big(\delta_{20}, \delta_{20} + \delta_{22} + \delta_{24}, \delta_{20} + \tfrac{1}{2}\delta_{22}\big).$$

We have

$$\begin{aligned}
\delta_{20} &= 4(t_v + t_w)^2 (1 - t_v t_w)^4 \geq 0,\\
\delta_{20} + \delta_{22} + \delta_{24} &= (1 - t_v^2)^2 (1 - t_w)^2 (1 + t_v t_w^2)^2 \geq 0,\\
2\delta_{20} + \delta_{22} &= (1 - t_w)(1 - t_v^2)(1 - t_v t_w)^2 \Lambda(t_v, t_w),
\end{aligned}$$

where Λ reads as a polynomial of degré 2 in the variable t_v

$$\Lambda(t_v, t_w) = (t_w^3 - t_w^2 + 3t_w + 1)t_v^2 - 2(1 + t_w)(t_w^2 - 4t_w + 1)t_v + (t_w^3 + 3t_w^2 - t_w + 1).$$

We prove that the polynomial Λ is non negative under the assumptions $-1 < t_v < 1$ and $-1 < t_w < t_w^\star(t_v)$. We compute the discriminant of Λ

$$\Delta_\Lambda = -16 t_w (1 - t_w)^2 (2 + 3t_w + 2t_w^2).$$

Then, Δ_Λ has the opposite sign to t_w. First case, $0 \leq t_w < 1$: $\Delta_\Lambda \leq 0$ implies that $\Lambda(t_v, t_w)$ is non negative as $\Lambda(0, t_w) = 1 - t_w + t_w^2(3 + t_w) > 0$. Second case, $-1 < t_w < 0$: as $\Delta_\Lambda > 0$, the polynomial Λ has two distinct real roots. As

$$\begin{aligned}
\Lambda(-1, t_w) &= 4(1 + t_w)(1 - t_w)^2 > 0,\\
\Lambda(1, t_w) &= 8 t_w (1 + t_w) < 0,
\end{aligned}$$

there is exactly one sign change of the polynomial Λ on the interval $[-1,1]$. We write $t_v^{\square}(t_w)$ for the zero of Λ that lies in $(-1,1)$ whenever $-1<t_w<0$. Thus, it suffices to show that

$$\Lambda(t_v^\star(t_w),t_w)\geq 0,$$

which in turn implies that Λ is strictly positive on the interval we are concerned with, namely $t_v\in(-1,t_v^\star(t_w))$. A direct calculation yields

$$\Lambda(t_v^\star(t_w),t_w)=\frac{4(1+t_w)^3}{\left(\sqrt{8+8t_w^4}+3(1-t_w^2)\right)^2}\left(3t_w^4+4t_w^3-2t_w^2-4t_w+3-\sqrt{8+8t_w^4}(1-t_w-t_w^2)\right).$$

We remark that

$$3t_w^4+4t_w^3-2t_w^2-4t_w+3=1+t_w^4+2(1-t_w-t_w^2)^2>0.$$

Then, we conclude that $\Lambda(t_v^\star(t_w),t_w)\geq 0$ as

$$(3t_w^4+4t_w^3-2t_w^2-4t_w+3)^2-8(1+t_w^4)(1-t_w-t_w^2)^2=(1-t_w^2)^2(1+4t_w-t_w^2)^2\geq 0.$$

At this point, we prove that

$$\theta_1^{\circ\circ}(\xi,c,t_v,t_w)\geq 0,\quad 0\leq\xi\leq\pi,\ 0\leq c\leq 1,\ -1<t_v<1,\ -1<t_w<t_w^\star(t_v).$$

For $t_w=t_w^\star(t_v)$, the continuity of the function $\theta_1^{\circ\circ}$ according to all its variables immediately implies that the inquality remains true for $-1<t_w\leq t_w^\star(t_v)$. We end the proof by assuming that $t_w>t_w^\star(t_v)$. Then $\Upsilon(t_v,t_w)<0$ implies

$$\theta_1^{\circ\circ}(0,c,t_v,t_w)=(1-t_w)(1-t_v^2)(1-t_vt_w)^2\Upsilon(t_v,t_w)<0.$$

As an final result, we obtain for $(t_v,t_w)\in\mathcal{I}_{t_vt_w}$ and $0<c<1$ that

$$\forall\xi\in[0,\pi]\ \theta_1^{\circ\circ}(\xi,c,t_v,t_w)\geq 0\iff\Upsilon(t_v,t_w)\geq 0\iff -1<t_w\leq t_w^\star(t_v),$$

relation which ends the proof. □

5.4. **Stability on $\mathcal{E}_{t_vt_w}$.** In this section, we consider the four edges $\mathcal{E}^b_{t_vt_w}$, $\mathcal{E}^t_{t_vt_w}$, $\mathcal{E}^l_{t_vt_w}$, and $\mathcal{E}^r_{t_vt_w}$. We prove that (see Fig. 6):

- for $|t_v|<1$ and $t_w=-1$, the scheme is stable for $c\in[0,1]$ (see Prop. 28);
- for $t_v=-1$ and $|t_w|<1$, the scheme is stable for $c\in[0,1]$ (see Prop. 29);
- for $|t_v|<1$ and $t_w=1$, the scheme is stable for $c=0$ and $c=1$ and is unstable for $c\in(0,1)$ (see Prop. 30);
- for $t_v=1$ and $|t_w|<1$, the scheme is stable for $c=0$ and $c=1$ and is unstable for $c\in(0,1)$ (see Prop. 31).




FIGURE 6. Stability for $(t_v,t_w)\in\mathcal{E}_{t_vt_w}$. Green means stable while purple means unstable.

Proposition 28 (stability on $\mathcal{E}^b_{t_v t_w}$)**.** *Considering $t_v \in (-1,1)$ and $t_w = -1$, the lattice Boltzmann scheme is stable for every value $c \in [0,1]$.*

Proof of Prop. 28. In this proof, we assume that $|t_v| < 1$ and $t_w = -1$. Lemma 20 then implies that $\varphi_3(\cdot;\xi,c,t_v,-1) \in \mathscr{I}$ for $\xi \in [0,\pi]$, $c \in [0,1]$. Then, applying Th. 8, $\varphi_3(\cdot;\xi,c,t_v,-1) \in \mathscr{N}^s$ if, and only if $\varphi_2^\circ(\cdot;\xi,c,t_v,-1) \in \mathscr{N}^s$. Lemmas 22 and 23 read then

$$\varphi_2^\circ(\cdot;\xi,c,t_v,-1) \in \mathscr{I} \iff (c,\xi) \in \mathcal{I}_{c\xi} \cup \mathcal{E}^l_{c\xi},$$
$$\varphi_2^\circ(\cdot;\xi,c,t_v,-1) \in \mathscr{E} \iff (c,\xi) \in \mathcal{E}^b_{c\xi} \cup \mathcal{E}^r_{c\xi} \cup \mathcal{E}^t_{c\xi} \cup \mathcal{V}_{c\xi}.$$

First, we consider $(c,\xi) \in \mathcal{I}_{c\xi} \cup \mathcal{E}^l_{c\xi}$, we have

$$\varphi_1^{\circ\circ}(X;\xi,c,t_v,-1) = (X-1)(1-t_v^2)^2(1-\cos^2\xi)(1-c^2) \in \mathscr{N}^s.$$

Second, we consider $(c,\xi) \in \mathcal{E}^b_{c\xi} \cup \mathcal{E}^r_{c\xi} \cup \mathcal{E}^t_{c\xi} \cup \mathcal{V}_{c\xi}$, we have

$$\varphi_1^{\circ\bullet}(X;\xi,c,t_v,-1) = (2X - 1 - \cos\xi + \mathrm{i}c\sin\xi)(1-t_v^2) \in \mathscr{N}^s.$$

We conclude that $\varphi_3(\cdot;\xi,c,t_v,-1) \in \mathscr{N}^s$ for every $(c,\xi) \in \mathcal{D}_{c\xi}$ and then that the scheme is stable for $c \in [0,1]$. □

Proposition 29 (stability on $\mathcal{E}^l_{t_v t_w}$)**.** *Considering $t_v = -1$ and $t_w \in (-1,1)$, the lattice Boltzmann scheme is stable for every value $c \in [0,1]$.*

Proof of Prop. 29. In this proof, we assume that $t_v = -1$ and $|t_w| < 1$. Lemma 20 then implies that $\varphi_3(\cdot;\xi,c,-1,t_w) \in \mathscr{I}$ for $\xi \in [0,\pi]$, $c \in [0,1]$. Then, applying Th. 8, $\varphi_3(\cdot;\xi,c,-1,t_w) \in \mathscr{N}^s$ if, and only if $\varphi_2^\circ(\cdot;\xi,c,-1,t_w) \in \mathscr{N}^s$. Lemmas 22 and 23 read then

$$\varphi_2^\circ(\cdot;\xi,c,-1,t_w) \in \mathscr{I} \iff (c,\xi) \in \mathcal{I}_{c\xi} \cup \mathcal{E}^l_{c\xi} \cup \mathcal{E}^t_{c\xi},$$
$$\varphi_2^\circ(\cdot;\xi,c,-1,t_w) \in \mathscr{E} \iff (c,\xi) \in \mathcal{E}^b_{c\xi} \cup \mathcal{E}^r_{c\xi} \cup \mathcal{V}_{c\xi}.$$

First, we consider $(c,\xi) \in \mathcal{I}_{c\xi} \cup \mathcal{E}^l_{c\xi} \cup \mathcal{E}^t_{c\xi}$, we have

$$\varphi_1^{\circ\circ}(X;c,\xi,-1,t_w) = (1-\cos\xi)(1-c^4)(1-t_w^2)^2(\alpha_1^{\circ\circ} - X\beta_1^{\circ\circ})$$

with $\mu = (1-c^2)/(1+c^2) \in (0,1]$, $\alpha_1^{\circ\circ} = \mu + \cos\xi$, and $\beta_1^{\circ\circ} = 1 + \mu\cos\xi \geq 0$. Then $\beta_1^{\circ\circ} \geq |\alpha_1^{\circ\circ}|$ as $(1-\mu)(1-\cos\xi) \geq 0$ and $(1+\mu)(1+\cos\xi) \geq 0$, which implies $\varphi_1^{\circ\circ}(X;c,\xi,-1,t_w) \in \mathscr{N}^s$. Second, we consider $(c,\xi) \in \mathcal{E}^b_{c\xi} \cup \mathcal{E}^r_{c\xi} \cup \mathcal{V}_{c\xi}$, we have

$$\varphi_1^{\circ\bullet}(X;c,\xi,-1,t_w) = (1-t_w^2)(2X - 1 + c^2 - (1+c^2)\cos\xi).$$

As $-2 \leq -2c^2 \leq 1-c^2+(1+c^2)\cos\xi \leq 2$, we obtain that $\varphi_1^{\circ\bullet}(X;c,\xi,-1,t_w) \in \mathscr{N}^s$. We conclude that $\varphi_3(\cdot;\xi,c,-1,t_w) \in \mathscr{N}^s$ for every $(c,\xi) \in \mathcal{D}_{c\xi}$ and then that the scheme is stable for $c \in [0,1]$. □

Proposition 30 (stability on $\mathcal{E}^t_{t_v t_w}$)**.** *Considering $t_v \in (-1,1)$ and $t_w = 1$, the lattice Boltzmann scheme is stable for $c = 0$ and $c = 1$ but unstable for every other values $c \in (0,1)$ (see Fig. 7).*

Proof of Prop. 30. In this proof, we assume that $|t_v| < 1$ and $t_w = 1$. Lemma 20 then implies that $\varphi_3(\cdot;\xi,c,t_v,1) \in \mathscr{I}$ for $\xi \in [0,\pi]$, $c \in [0,1]$. Then, applying Th. 7,




FIGURE 7. Stability for $(t_v, t_w) \in \mathcal{E}^t_{t_v t_w}$.

$\varphi_3(\cdot;\xi,c,t_v,1) \in \mathscr{N}$ if, and only if $\varphi_2^\circ(\cdot;\xi,c,t_v,1) \in \mathscr{N}$. Lemmas 22 and 23 read then

$$\varphi_2^\circ(\cdot;\xi,c,t_v,1) \in \mathscr{I} \iff (c,\xi) \in \mathcal{I}_{c\xi} \cup \mathcal{E}^l_{c\xi},$$
$$\varphi_2^\circ(\cdot;\xi,c,t_v,1) \in \mathscr{E} \iff (c,\xi) \in \mathcal{E}^b_{c\xi} \cup \mathcal{E}^r_{c\xi} \cup \mathcal{E}^t_{c\xi} \cup \mathcal{V}_{c\xi}.$$

First, we consider $(c,\xi) \in \mathcal{I}_{c\xi} \cup \mathcal{E}^l_{c\xi}$, we have

$$\varphi_1^{\circ\circ}(X;\xi,c,t_v,1) = (1-t_v^2)^2(1-c^2)(1-\cos\xi)\big(X(1+\cos\xi) - (1+\cos\xi - 2\mathrm{i}c\sin\xi)\big)$$

As $\sin\xi \neq 0$ and $\cos\xi \neq -1$ for $(c,\xi) \in \mathcal{I}_{c\xi}$, the root $1-2\mathrm{i}c\sin\xi/(1+\cos\xi)$ is located outside the closed unit circle and $\varphi_1^{\circ\circ}(\cdot;\xi,c,t_v,1) \notin \mathscr{N}$. However, $\varphi_1^{\circ\circ}(\cdot;\xi,0,t_v,1) \in \mathscr{N}^s$. Second, we consider $(c,\xi) \in \mathcal{E}^b_{c\xi} \cup \mathcal{E}^r_{c\xi} \cup \mathcal{E}^t_{c\xi} \cup \mathcal{V}_{c\xi}$, we have

$$\varphi_1^{\circ\bullet}(X;0,c,t_v,1) = 2X(1-t_v^2) \in \mathscr{N}^s, \qquad c \in [0,1],$$
$$\varphi_1^{\circ\bullet}(X;\pi,c,t_v,1) = 2(1-t_v^2)\big(X(2c^2-1)+1\big) \in \mathscr{N}^s, \qquad c \in [0,1],$$
$$\varphi_1^{\circ\bullet}(X;\xi,1,t_v,1) = (1-t_v^2)(2X+1-\cos\xi+\mathrm{i}\sin\xi) \in \mathscr{N}^s, \qquad \xi \in [0,\pi].$$

We conclude that $\varphi_3(\cdot;\xi,c,t_v,1) \in \mathscr{N}^s$ for every $(c,\xi) \in \mathcal{E}_{c\xi} \cup \mathcal{V}_{c\xi}$, and that $\varphi_3(\cdot;\xi,c,t_v,1) \notin \mathscr{N}$ for $(c,\xi) \in \mathcal{I}_{c\xi}$. Finally, the scheme is stable for $c=0$ and $c=1$ and is unstable for $c \in (0,1)$. □

Proposition 31 (stability on $\mathcal{E}^r_{t_v t_w}$)**.** *Considering $t_v = 1$ and $t_w \in (-1,1)$, the lattice Boltzmann scheme is stable for $c=0$ and $c=1$ but unstable for every other values $c \in (0,1)$ (see Fig. 8).*

Proof of Prop. 31. We assume in this proof that $t_v = 1$ and $|t_w| < 1$. Lemma 20 then implies that $\varphi_3(\cdot;\xi,c,1,t_w) \in \mathscr{I}$ for $\xi \in [0,\pi]$, $c \in [0,1]$. Then, applying Th. 8, $\varphi_3(\cdot;\xi,c,1,t_w) \in \mathscr{N}^s$ if, and only if $\varphi_2^\circ(\cdot;\xi,c,1,t_w) \in \mathscr{N}^s$. Lemmas 22 and 23 read then

$$\varphi_2^\circ(\cdot;\xi,c,1,t_w) \in \mathscr{I} \iff (c,\xi) \in \mathcal{I}_{c\xi} \cup \mathcal{E}^l_{c\xi} \cup \mathcal{E}^t_{c\xi},$$
$$\varphi_2^\circ(\cdot;\xi,c,1,t_w) \in \mathscr{E} \iff (c,\xi) \in \mathcal{E}^b_{c\xi} \cup \mathcal{E}^r_{c\xi} \cup \mathcal{V}_{c\xi}.$$




FIGURE 8. Stability for $(t_v, t_w) \in \mathcal{E}^r_{t_v t_w}$.

First, we consider $(c,\xi) \in \mathcal{I}_{c\xi} \cup \mathcal{E}^l_{c\xi} \cup \mathcal{E}^t_{c\xi}$, we have

$$\begin{aligned}
\varphi_1^{\circ\circ}(X;c,\xi,1,t_w) &= (1-\cos\xi)(1-c^2)(1-t_w^2)^2 \\
&\qquad \big((1+\cos\xi+c^2(1-\cos\xi))(X-1)+2\mathrm{i}c\sin\xi\big) \notin \mathscr{N}, \quad (c,\xi)\in\mathcal{I}_{c\xi},\\
\varphi_1^{\circ\circ}(X;0,\xi,1,t_w) &= (1-\cos\xi)(1-t_w^2)^2(1+\cos\xi)(X-1) \in \mathscr{N}^s, \quad \xi\in(0,\pi),\\
\varphi_1^{\circ\circ}(X;c,\pi,1,t_w) &= 4c^2(1-c^2)(1-t_w^2)^2(X-1) \in \mathscr{N}^s, \quad c\in(0,1).
\end{aligned}$$

Second, we consider $(c,\xi) \in \mathcal{E}^b_{c\xi} \cup \mathcal{E}^r_{c\xi} \cup \mathcal{V}_{c\xi}$, we have

$$\varphi_1^{\circ\bullet}(X;c,\xi,1,t_w) = (1-t_w^2)(2X-(1-c^2)(1-\cos\xi)+2\mathrm{i}c\sin\xi).$$

Then

$$\begin{aligned}
\varphi_1^{\circ\bullet}(X;c,0,1,t_w) &= 2(1-t_w^2)X \in \mathscr{N}^s, \quad c\in[0,1],\\
\varphi_1^{\circ\bullet}(X;1,\xi,1,t_w) &= 2(1-t_w^2)(X+\mathrm{i}\sin\xi) \in \mathscr{N}^s, \quad \xi\in[0,\pi],\\
\varphi_1^{\circ\bullet}(X;0,\pi,1,t_w) &= 2(1-t_w^2)(X-1) \in \mathscr{N}^s.
\end{aligned}$$

We conclude that $\varphi_3(\cdot;\xi,c,1,t_w) \in \mathscr{N}^s$ for every $(c,\xi) \in \mathcal{E}_{c\xi} \cup \mathcal{V}_{c\xi}$ and that $\varphi_3(\cdot;\xi,c,1,t_w) \notin \mathscr{N}$ for $(c,\xi) \in \mathcal{I}_{c\xi}$. Finally, the scheme is stable for $c=0$ and $c=1$ and is unstable for $c\in(0,1)$. □

5.5. **Stability in $\mathcal{V}_{t_v t_w}$.** In this section, we consider the four vertices $\mathcal{V}^{lb}_{t_v t_w}$, $\mathcal{V}^{rb}_{t_v t_w}$, $\mathcal{V}^{lt}_{t_v t_w}$, and $\mathcal{V}^{rt}_{t_v t_w}$. We prove that (see Fig. 9):

- for $t_v=-1$ and $t_w=-1$, the scheme is stable for $c\in[0,1]$ (see Prop. 32);
- for $t_v=1$ and $t_w=-1$, the scheme is stable for $c=0$ and unstable for $c\in(0,1]$ (see Prop. 33);
- for $t_v=-1$ and $t_w=1$, the scheme is stable for $c\in(0,1)$ and unstable for $c\in\{0,1\}$ (see Prop. 34);
- for $t_v=1$ and $t_w=1$, the scheme is unstable for $c\in[0,1]$ (see Prop. 35).

Proposition 32 (stability on $\mathcal{V}^{lb}_{t_v t_w}$)**.** *Considering $t_v=-1$ and $t_w=-1$, the lattice Boltzmann scheme is stable for every value $c\in[0,1]$.*




FIGURE 9. Stability for $(t_v, t_w) \in \mathcal{V}_{t_v t_w}$. Green means stable while purple means unstable.

Proof of Prop. 32. Assuming $t_v = -1$ and $t_w = -1$, the amplification matrix $\boldsymbol{G}$ reads

$$\boldsymbol{G} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & e^{-\mathrm{i}\xi} & 0 \\ 0 & 0 & e^{\mathrm{i}\xi} \end{pmatrix}.$$

The scheme is then clearly stable as the matrix $\boldsymbol{G}$ is diagonal for every $\xi \in [0, \pi]$ with eigenvalues on the unit circle. □

Proposition 33 (stability on $\mathcal{V}^{rb}_{t_v t_w}$)**.** *Considering $t_v = 1$ and $t_w = -1$, the lattice Boltzmann scheme is stable for $c = 0$ and unstable for $c \in (0, 1]$ (see Fig. 10).*




FIGURE 10. Stability for $(t_v, t_w) \in \mathcal{V}^{rb}_{t_v t_w}$.

Proof of Prop. 33. Assuming $t_v = 1$ and $t_w = -1$, the characteristic polynomial given in (12) yields to

$$\begin{aligned} \varphi_3(X; \xi, c, 1, -1) &= X^3 - X^2(1 - 2\mathrm{i}c\sin\xi) - X(1 + 2\mathrm{i}c\sin\xi) + 1 \\ &= (X - 1)(X^2 + 2\mathrm{i}c\sin\xi X - 1) \end{aligned}$$

for which the roots are 1, $\sqrt{1-c^2\sin^2\xi}+\mathrm{i}c\sin\xi$, and $-\sqrt{1-c^2\sin^2\xi}+\mathrm{i}c\sin\xi$. Then $\varphi_3 \in \mathscr{N}$ for $(c,\xi) \in \mathcal{D}_{c\xi}$ and

$$\varphi_3 \in \mathscr{N}^s \iff (c,\xi) \in \mathcal{I}_{c\xi} \cup \mathcal{E}^r_{c\xi} \backslash \{(1,\pi/2)\}.$$

For $(c,\xi) \in \mathcal{E}^l_{c\xi} \cup \mathcal{V}^{lt}_{c\xi} \cup \mathcal{V}^{lb}_{c\xi}$, that is $c=0$ and $0 \le \xi \le \pi$, the minimal polynomial of the amplification matrix

$$\boldsymbol{G} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & e^{-\mathrm{i}\xi} \\ 0 & e^{\mathrm{i}\xi} & 0 \end{pmatrix}$$

reads $\mu = X^2 - 1 \in \mathscr{N}^s$. The scheme is then stable for $c=0$. For $(c,\xi) \in \mathcal{E}^b_{c\xi} \cup \mathcal{V}^{lb}_{c\xi} \cup \mathcal{V}^{rb}_{c\xi}$, that is $0 \le c \le 1$ and $\xi = 0$, the minimal polynomial of the amplification matrix

$$\boldsymbol{G} = \begin{pmatrix} 1 & 0 & 0 \\ c & c & 1+c \\ -c & 1-c & -c \end{pmatrix}$$

reads also $\mu = X^2 - 1 \in \mathscr{N}^s$. For $(c,\xi) \in \mathcal{E}^t_{c\xi} \cup \mathcal{V}^{rt}_{c\xi}$, that is $0 < c \le 1$ and $\xi = \pi$, the minimal polynomial of the amplification matrix

$$\boldsymbol{G} = \begin{pmatrix} 1 & 0 & 0 \\ -c & -c & -1-c \\ c & -1+c & c \end{pmatrix}$$

reads $\mu = (X-1)(X^2-1) \notin \mathscr{N}^s$. The last case is then $c=1$ and $\xi = \pi/2$ for which the minimal polynomial of the matrix

$$\boldsymbol{G} = \begin{pmatrix} 1 & 0 & 0 \\ -\mathrm{i} & -\mathrm{i} & -2\mathrm{i} \\ -\mathrm{i} & 0 & -\mathrm{i} \end{pmatrix}$$

reads $\mu = (X-1)(X+\mathrm{i})^2 \notin \mathscr{N}^s$ as $\boldsymbol{G}^2$ is not equal to $(1-\mathrm{i})\boldsymbol{G} + \mathrm{i}\boldsymbol{I}$. The scheme is then unstable for $c > 0$. □

Proposition 34 (stability on $\mathcal{V}^{lt}_{t_v t_w}$)**.** *Considering $t_v = -1$ and $t_w = 1$, the lattice Boltzmann scheme is stable for every value $c \in (0,1)$ and unstable for $c \in \{0,1\}$ (see Fig. 11).*

Proof of Prop. 34. Assuming $t_v = -1$ and $t_w = 1$, the characteristic polynomial given in (12) yields to

$$\begin{aligned}\varphi_3(X;\xi,c,-1,1) &= X^3 + X^2(-1+2c^2(1-\cos\xi)) + X(-1+2c^2(1-\cos\xi)) + 1 \\ &= (1+X)(1-2X(1-c^2(1-\cos\xi)) + X^2)\end{aligned}$$

for which the roots are -1, $\alpha + \mathrm{i}\sqrt{1-\alpha^2}$ and $\alpha - \mathrm{i}\sqrt{1-\alpha^2}$ with $\alpha = 1-c^2(1-\cos\xi)$. Then $\varphi_3 \in \mathscr{N}$ for $(c,\xi) \in \mathcal{D}_{c\xi}$ and

$$\varphi_3 \in \mathscr{N}^s \iff \alpha^2 \neq 1 \iff (c,\xi) \in \mathcal{I}_{c\xi} \cup \mathcal{E}^t_{c\xi} \cup \mathcal{E}^r_{c\xi}.$$

We first investigate the stability for $c=0$. For $(c,\xi) \in \mathcal{E}^l_{c\xi} \cup \mathcal{V}^{lt}_{c\xi}$, that is $c=0$ and $0 < \xi \le \pi$, the minimal polynomial of the amplification matrix

$$\boldsymbol{G} = \begin{pmatrix} 1 & 2 & 2 \\ 0 & 0 & -e^{-\mathrm{i}\xi} \\ 0 & -e^{\mathrm{i}\xi} & 0 \end{pmatrix}$$




FIGURE 11. Stability for $(t_v, t_w) \in \mathcal{V}^{lt}_{t_v t_w}$.

reads $\mu = (X-1)^2(X+1)$ as $\boldsymbol{G}^2$ is not equal to $\boldsymbol{I}$:

$$\boldsymbol{G}^2 = \begin{pmatrix} 1 & 2(1-e^{\mathrm{i}\xi}) & 2(1-e^{-\mathrm{i}\xi}) \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

This shows that the scheme is unstable for $c = 0$. We remark that the matrix $\boldsymbol{G}$ reduces to identity for $\xi = 0$ but it is evidently not enough to impose the stability of the scheme.

Second, we investigate the stability for $c = 1$. The necessary condition involving the characteristic polynomial is fulfilled for $0 < \xi < \pi$ ($(c, \xi) \in \mathcal{E}^{r}_{c\xi}$). We remark first that for $(c, \xi) \in \mathcal{V}^{rb}_{c\xi}$, that is $c = 1$ and $\xi = 0$, the minimal polynomial of the amplification matrix

$$\boldsymbol{G} = \begin{pmatrix} -1 & 0 & 0 \\ 1 & 1 & 0 \\ 1 & 0 & 1 \end{pmatrix}$$

reads $\mu = X^2 - 1 \in \mathscr{N}^s$ and second that for $(c, \xi) \in \mathcal{V}^{rt}_{c\xi}$, that is $c = 1$ and $\xi = \pi$, the minimal polynomial of the amplification matrix

$$\boldsymbol{G} = \begin{pmatrix} -1 & 0 & 0 \\ -1 & -1 & 0 \\ -1 & 0 & -1 \end{pmatrix}$$

reads $\mu = (X+1)^2 \notin \mathscr{N}^s$, hence the scheme is unstable for $c = 1$.

Finally, we investigate the stability for $0 < c < 1$. For $(c, \xi) \in \mathcal{E}^{b}_{c\xi}$, that is $0 < c < 1$ and $\xi = 0$, the minimal polynomial of the amplification matrix

$$\boldsymbol{G} = \begin{pmatrix} 1-2c^2 & 2(1-c^2) & 2(1-c^2) \\ c^2 & c^2 & c^2-1 \\ c^2 & c^2-1 & c^2 \end{pmatrix}$$

reads $\mu = X^2 - 1 \in \mathscr{N}^s$ as $\boldsymbol{G}^2 = \boldsymbol{I}$. This means that the necessary condition for stability is fulfilled. To show that the scheme is indeed stable, we argue by a direct

computation of $\boldsymbol{G}^n$ for $\xi \in [0,\pi]$.

$$\boldsymbol{G} = \begin{pmatrix} c^2(\cos\xi - \mathrm{i}\sin\xi) & c^2(\cos\xi - \mathrm{i}\sin\xi) & -(1-c^2)(\cos\xi - \mathrm{i}\sin\xi) \\ c^2(\cos\xi + \mathrm{i}\sin\xi) & -(1-c^2)(\cos\xi + \mathrm{i}\sin\xi) & c^2(\cos\xi + \mathrm{i}\sin\xi) \end{pmatrix}.$$

This matrix is diagonalisable on $\mathbb{C}$ for every $\xi \in [0,\pi]$ with eigenvalues $\lambda_0 = -1$, $\lambda_\pm = \alpha \pm \mathrm{i}\sqrt{1-\alpha^2}$, with $\alpha = 1 - c^2(1-\cos\xi) \in (-1,1]$. Using half and double angle formulas for sine and cosine, we have

$$\alpha = 1 - 2c^2\sin(\xi/2)^2, \qquad \lambda_\pm = 1 - 2c^2\sin(\xi/2)^2 \pm 2\mathrm{i}c\sin(\xi/2)\sqrt{1-c^2\sin(\xi/2)^2}.$$

Eigenvectors corresponding to these eigenvalues are given by

$$V_0 = \begin{pmatrix} -1-e^{-\mathrm{i}\xi} \\ e^{-\mathrm{i}\xi} \\ 1 \end{pmatrix}, \quad V_\pm = \begin{pmatrix} 2(1-c^2) \\ ce^{-\mathrm{i}\xi/2}(c\sin(\xi/2) \mp \sqrt{1-c^2\sin(\xi/2)^2}) \\ ce^{-\mathrm{i}\xi/2}(c\sin(\xi/2) \pm \sqrt{1-c^2\sin(\xi/2)^2}) \end{pmatrix}.$$

Defining $\boldsymbol{P}$ the change of basis and $\boldsymbol{\Lambda}$ the diagonal matrix of the eigenvalues

$$\boldsymbol{P} = \left(\begin{array}{c|c|c} V_0 & V_- & V_+ \end{array}\right) \qquad \boldsymbol{\Lambda} = \begin{pmatrix} \lambda_0 & 0 & 0 \\ 0 & \lambda_- & 0 \\ 0 & 0 & \lambda_+ \end{pmatrix},$$

we have

$$\boldsymbol{G}^n = \frac{1}{\det \boldsymbol{P}} \boldsymbol{P}\boldsymbol{\Lambda}^n \operatorname{Com}(\boldsymbol{P})^T.$$

The important point is that

$$\det \boldsymbol{P} = -8c(1-c^2\sin(\xi/2)^2)^{3/2}e^{-\mathrm{i}\xi/2} \implies |\det \boldsymbol{P}| \geq 8c(1-c^2)^{3/2} > 0,$$

as $0 < c < 1$. This proves that the iteration matrix is uniformly power bounded, and concludes this part of the proof. □

Proposition 35 (stability on $\mathcal{V}^{rt}_{t_v t_w}$)**.** *Considering $t_v = 1$ and $t_w = 1$, the lattice Boltzmann scheme is unstable for $c \in [0,1]$ (see Fig. 12).*




FIGURE 12. Stability for $(t_v, t_w) \in \mathcal{V}^{rt}_{t_v t_w}$.

Proof of Prop. 35. Assuming $t_v = 1$ and $t_w = 1$, the characteristic polynomial given in (12) yields to

$$\varphi_3(X;\xi,c,1,1) = X^3 + \alpha X^2 - \bar{\alpha}X - 1,$$

with

$$\alpha = -1 + 2c^2(1-\cos\xi) + 2\cos\xi + 2\mathrm{i}c\sin\xi.$$

As $\varphi_2^\circ = 0$ ($\varphi_3 \in \mathscr{E}$), we have $\varphi_3 \in \mathscr{N} \iff \varphi_2^\bullet \in \mathscr{N}$ thanks to Th. 7. Computing $\varphi_2^\bullet = 3X^2 + 2\alpha X - \bar{\alpha}$, we obtain that

$$\mathscr{C}(\varphi_2^\bullet) = \varphi_1^{\bullet\circ} = (9 - |\alpha|^2)X + 2(3\alpha + \bar{\alpha}^2).$$

We then reduce

$$\varphi_3 \in \mathscr{N} \iff \varphi_2^\bullet \in \mathscr{N} \iff \begin{cases} |\alpha|^2 < 9 \text{ and } 2|3\alpha + \bar{\alpha}^2| \le 9 - |\alpha|^2 \\ \text{or} \\ |\alpha|^2 = 9 \text{ and } 3\alpha + \bar{\alpha}^2 = 0, \end{cases}$$
$$\iff |\alpha|^2 \le 9 \text{ and } 2|3\alpha + \bar{\alpha}^2| \le 9 - |\alpha|^2.$$

Thanks to lemma 36, the inequality $|\alpha|^2 \le 9$ is always true. We obtain

$$(9 - |\alpha|^2)^2 - 4|3\alpha + \bar{\alpha}^2|^2 = 48(1-\cos\xi)\beta(c^2, \cos\xi)$$

with

$$\begin{aligned}\beta(c^2, C) = {} & (1-c^2)(c^6 + 5c^4 - 8c^2 + 1) + C(1-c^2)(-3c^6 - 5c^4 + 6c^2 + 3) \\ & + C^2(-3c^8 + 8c^6 - 13c^4 + 6c^2 + 3) + C^3(1 + c - c^2)^2(1 - c - c^2)^2.\end{aligned}$$

Fixing $c \in [0,1]$, we consider $\beta(c^2, C)$ as a polynomial of degree 3 in terms of C.

First, fixing $c = 0$, we have $\beta(0, C) = 1 + 3C + 3C^2 + C^3 = (1+C)^3 \ge 0$, then $\varphi_3 \in \mathscr{N}$. To conclude the case $\mathcal{E}^l_{c\xi} \cup \mathcal{V}^{lb}_{c\xi} \cup \mathcal{V}^{lt}_{c\xi}$, we write the amplification matrix $\boldsymbol{G}$:

$$\boldsymbol{G} = \begin{pmatrix} 1 & 2 & 2 \\ 0 & -e^{-\mathrm{i}\xi} & 0 \\ 0 & 0 & -e^{\mathrm{i}\xi} \end{pmatrix}.$$

For $\xi \in [0, \pi]$, the minimal polynomial of the amplification matrix reads

$$\begin{aligned} &\mu(X;0,0,1,1) = (X-1)(X+1), \\ &\mu(X;\xi,0,1,1) = (X-1)(X+e^{-\mathrm{i}\xi})(X+e^{\mathrm{i}\xi}), \qquad 0 < \xi < \pi, \\ &\mu(X;\pi,0,1,1) = (X-1)^2. \end{aligned}$$

It is then a simple von Neumann polynomial only for $\xi \in [0, \pi)$. The scheme is then clearly unstable for $c = 0$.

Second, fixing $c = 1$, we have $\beta(1, C) = C^2(1+C) \ge 0$, then $\varphi_3 \in \mathscr{N}$. To conclude the case $\mathcal{E}^r_{c\xi} \cup \mathcal{V}^{rb}_{c\xi} \cup \mathcal{V}^{rt}_{c\xi}$, we write the amplification matrix $\boldsymbol{G}$:

$$\boldsymbol{G} = \begin{pmatrix} -1 & 0 & 0 \\ 2e^{-\mathrm{i}\xi} & e^{-\mathrm{i}\xi} & 2e^{-\mathrm{i}\xi} \\ 0 & 0 & -e^{\mathrm{i}\xi} \end{pmatrix}.$$

The minimal polynomial of the amplification matrix reads

$$\mu(X;\xi,1,1,1) = (X+1)(X-e^{-\mathrm{i}\xi})(X+e^{\mathrm{i}\xi})$$

for $\xi \in (0, \pi]$ and $\mu(X;0,1,1,1) = (X-1)(X+1)$. It is then a simple von Neumann polynomial only for $\xi \notin \{\pi/2, \pi\}$. The scheme is then clearly unstable for $c = 1$.

Third, fixing $\xi = 0$, the equality $\cos\xi = 1$ implies immediately that $\varphi_3 \in \mathscr{N}$. To conclude the case $\mathcal{E}^b_{c\xi}$, we write the amplification matrix $\boldsymbol{G}$:

$$\boldsymbol{G} = \begin{pmatrix} 1-2c^2 & 2(1-c^2) & 2(1-c^2) \\ c(1+c) & -1+c+c^2 & c(1+c) \\ -c(1-c) & -c(1-c) & -1-c+c^2 \end{pmatrix}.$$

The characteristic polynomial reads, as $\alpha = 1$, $\varphi_3(X;0,c,1,1) = (X+1)^2(X-1)$. As $\boldsymbol{G}^2$ is the identity matrix for $c \in [0,1]$, the minimal polynomial of the amplification matrix is a simple von Neumann polynomial.

Fourth, fixing $\xi = \pi$, we have $\beta(c^2,-1) = 8c^6(1-c^2) \geq 0$, then $\varphi_3 \in \mathscr{N}$. To conclude the case $\mathcal{E}^t_{c\xi}$, we write the amplification matrix $\boldsymbol{G}$:

$$\boldsymbol{G} = \begin{pmatrix} 1-2c^2 & 2(1-c^2) & 2(1-c^2) \\ -c(1+c) & 1-c-c^2 & -c(1+c) \\ c(1-c) & c(1-c) & 1+c-c^2 \end{pmatrix}.$$

The characteristic polynomial reads, as $\alpha = -4(1-c^2)$, $\varphi_3(X;\pi,c,1,1) = (X-1)(X^2 - 2(1-2c^2)X + 1)$. The polynomial $X^2 - 2(1-2c^2)X + 1$ is positive if $c \in (0,1)$. Then φ_3 is a simple von Neumann polynomial for $0 < c < 1$.

Finally, we treat the case $(c,\xi) \in \mathcal{I}_{c\xi}$. For a fixed value of c, the function $\beta(c^2,C)$ is a polynomial of degree 3 in the C variable with a non negative coefficient of higher degree. We could write the roots and observe that exactly two of them are localized in $(-1,1)$, the third one being in $(-\infty,-1)$. The area where β is negative is given in Fig. 12 for illustration purposes. In order to prove that the scheme is unstable for every $c \in (0,1)$, we only prove that there exists a value of $\xi \in (0,\pi)$ defined by $\cos\xi = -1+c^2$ such that $\beta(c^2,-1+c^2) < 0$ for every $c \in (0,1)$. We have

$$\begin{aligned} \beta(c^2,-1+c^2) &= -c^4(1-c^2)(c^8 - 11c^6 + 38c^4 - 54c^2 + 27), \\ &= -c^4(1-c^2)\left(c^8 + 11c^4(1-c^2) + 27(1-c^2)^2\right) < 0, \end{aligned}$$

what ends the proof. □

Lemma 36 (technical lemma)**.** *Let's $(c,\xi) \in \mathcal{D}_{c\xi}$ defined in (18). In the complex plane, the parameter α defined by*

$$\alpha = -1 + 2c^2(1-\cos\xi) + 2\cos\xi + 2ic\sin\xi,$$

lies in the rectangular triangle defined by its three vertices -3, 1 and $1+2i$ (see Fig. 13). As a consequence, $|\alpha| \leq 3$ and

$$|\alpha| = 3 \iff \alpha = -3 \iff (c,\xi) \in \mathcal{V}^{lt}_{c\xi} \ (c = 0 \text{ and } \xi = \pi)$$

Proof. The parameter α is in the given rectangular triangle if, and only if

$$-3 \leq \mathrm{Re}(\alpha) \leq 1, \quad 0 \leq \mathrm{Im}(\alpha) \leq 2, \quad \text{and} \quad 2\,\mathrm{Im}(\alpha) \leq 3 + \mathrm{Re}(\alpha).$$

Rewritting $\mathrm{Re}(\alpha) = 1 - 2(1-c^2)(1-\cos\xi)$ implies $-3 \leq \mathrm{Re}(\alpha) \leq 1$. Moreover, $\mathrm{Im}(\alpha) = 2c\sin\xi$ implies $0 \leq \mathrm{Im}(\alpha) \leq 2$ as $0 \leq c \leq 1$ and $0 \leq \xi \leq \pi$. Finally, we compute

$$3 + \mathrm{Re}(\alpha) - 2\,\mathrm{Im}(\alpha) = 2\left(\sqrt{1+\cos\xi} - c\sqrt{1-\cos\xi}\right)^2 \geq 0.$$

To conclude the proof, we observe that there is only one point in the triangle that intersects the circle of radius 3: the point -3. Then $|\alpha| = 3$ is equivalent to $\alpha = -3$. Solving the system $\mathrm{Re}(\alpha) = -3$ and $\mathrm{Im}(\alpha) = 0$ yields to $c = 0$ and $\xi = \pi$. □




FIGURE 13. Graphical illustration of lemma 36: the parameter α lies in a rectangular triangle included in the circle of radius 3.

6. Convergence and discrete waves

In order to avoid dealing with borderline cases, we here consider $(s_v, s_w) \in (0,2)^2$. The fact of having $s_v \neq 0$ and $s_w \neq 0$ ensures second-order consistency regardless of the specific values of these parameters. We also consider parameters where the finite difference scheme whose amplification polynomial is the characteristic polynomial of $\boldsymbol{G}$ is L^2-stable. In this way, we can easily re-use known result on this one, see Remark 14. This gives the following result.

Proposition 37 (Theorem 10.6.1 in [37])**.** *Let $|c| \leq 1$, $0 < s_v < 2$, and $0 < s_w \leq s_w^\star(s_v)$. Then, there exists a unique eigenvalue $z_\varphi(\xi)$ defined for $|\xi| \leq \xi_0$ for some $\xi_0 > 0$ such that $z_\varphi(\xi) = 1 - ic\xi + o(\xi)$. Moreover, there exists a constant $C > 0$ such that*

$$\left| \frac{e^{-ic\xi\Delta x} - z_\varphi(\xi\Delta x)}{\Delta t} \right| \leq C\Delta x^2 (1+|\xi|)^3. \tag{22}$$

Proof. All except the power of $1+|\xi|$ is Theorem 10.6.1 in [37]. This power is obtained by (23). □

Equation (22) is a way of restating that the scheme is second-order accurate. Moreover, the power in ξ on the right-hand side of (22) dictates the minimal smoothness of the initial datum to observe full second-order convergence. The following convergence theorem is adapted from finite difference, and based on the fact that the scheme behaves essentially as a one-step scheme with symbol $z_\varphi(\xi)$, as the other eigenvalues are mastered by stability.

Theorem 38 (Convergence)**.** *Let $|c| \leq 1$, $0 < s_v < 2$, and $0 < s_w \leq s_w^\star(s_v)$. Let $u^\circ \in H^\sigma(\mathbb{R})$ in (2) with $\sigma \geq 3$ and $u_j^0 = u^\circ(j\Delta x)$—which is possible thanks to the smoothness assumption—and initial non-conserved moments at equilibrium, namely*

$$v_j^0 = \lambda c\, u_j^0, \qquad w_j^0 = \tfrac{1}{2}\lambda^2 c^2\, u_j^0.$$

Then, for all $T > 0$, there exists $C_T > 0$ such that

$$\Big(\Delta x \sum_{j\in\mathbb{Z}} \big|u(n\Delta t, j\Delta x) - u_j^n\big|^2\Big)^{1/2} \leq C_T \Delta x^2 \|u^\circ\|_{H^3}, \qquad 0 \leq n\Delta t \leq T,$$

where u is the solution of (1) and (2).

Proof. This is [37, Theorem 10.6.2] using Prop. 37 and the fact that initializations at equilibrium yield first-order accurate (see [6]) initialization schemes, given in (13) from Remark 14. □

In term of expected order of convergence for non-smooth initial data, if $u^\circ \in H^{\sigma-}(\mathbb{R})$ for the Sobolev regularity, and $u^\circ \in B^\sigma_{2,\infty}(\mathbb{R})$ for the Besov regularity, from [37] for the Sobolev regularity (up to "invisible" logarithmic terms) and from [10], we expect the empirical order of convergence with respect to the L^2 norm in space be $\mathcal{O}(\Delta x^{\min(2,2/3\sigma)})$.

We finish on wave propagation in the scheme by considering that—if the three roots of the characteristic polynomial that we now call $z_\varphi(\xi)$, $z_{\sigma,1}(\xi)$, and $z_{\sigma,2}(\xi)$ are distinct for a given ξ—any moment (or distribution function)[1] can be written

$$\hat{u}^n(\xi) = z_\varphi(\xi)^n C_\varphi(\xi) + z_{\sigma,1}(\xi)^n C_{\sigma,1}(\xi) + z_{\sigma,2}(\xi)^n C_{\sigma,2}(\xi),$$

where $C_\varphi(\xi)$, $C_{\sigma,1}(\xi)$, and $C_{\sigma,2}(\xi)$ are determined by the initialization, with the last two being generally small at least close to $\xi = 0$. This formula expresses that one can interpret the numerical solution as a superposition of the action of three pseudo-schemes, and is therefore interesting to see how each one propagates and damps information, especially for spatially smooth solutions, i.e. in a low-frequency limit.

One can solve the cubic equation with the help of formal computations and perform Taylor expansions for $\xi \to 0$. However, this is cumbersome with the free parameters c, s_v, and s_w. A more elegant way—up to first order—would boil down to employ the first-order perturbation theory of simple eigenvalues, see [23, Theorem 1]. For second-order result, we could also turn to the procedure by [27, Chapter II]. What we actually employ is injecting a power series expansion in ξ of the eigenvalues into the expansion of the characteristic equation and solve order-by-order, which yields

$$\begin{aligned}
(23)\quad z_\varphi(\xi) &= 1 - ic\xi - \tfrac{1}{2}c^2\xi^2 + ic\Big((1-c^2)\big((\tfrac{1}{2}-\tfrac{1}{s_v})(\tfrac{1}{2}-\tfrac{1}{s_w})-\tfrac{1}{12}\big)+\tfrac{c^2}{6}\Big)\xi^3 + \mathcal{O}(\xi^4),\\
z_{\sigma,1}(\xi) &= (1-s_v)\Big(1 + ic\xi + \tfrac{2c^2(s_w-1)-s_v-s_w+2}{2(s_v-s_w)}\xi^2 + \mathcal{O}(\xi^3)\Big), \quad \text{and}\\
z_{\sigma,2}(\xi) &= (1-s_w)\Big(1 + \tfrac{(1-c^2)(-s_v-s_w+2)}{2(s_v-s_w)}\xi^2 + \mathcal{O}(\xi^3)\Big).
\end{aligned}$$

The first equation shows that the method is second-order accurate and can become third-order accurate when $(\frac{1}{2}-\frac{1}{s_v})(\frac{1}{2}-\frac{1}{s_w}) = \frac{1}{12}$, which is however forbidden by stability reasons. Moreover, we see that the regular part of the solution carried by the first parasitic mode is geometrically damped if $0 < s_v < 2$ and propagates at dimensionless velocity $-c$. The second parasitic mode is damped if $0 < s_w < 2$ and does not propagate. Notice that taking $s_w = 2 - s_v$, we obtain $z_{\sigma,2} \equiv 1 - s_w = s_v - 1$: this mode can only carries constants.

[1] Here, we illustrate things for the conserved moment u.

7. Numerical experiments




Figure 14. Conserved moment of the scheme as function of space and time for different relaxation parameters. In (a), propagation of the mode brought by $z_{\sigma,1}$ at speed $-c$ is clearly visible (right-to-left beam), as its damping is weak. In (b), the damping of $z_{\sigma,1}$ and $z_{\sigma,2}$ is significant, thus only the physical mode z_φ is observed (left-to-right beam), featuring dispersion. In (c), wee see that the solution is very inaccurate as s_v is very close to zero. Finally, in (d), which is an unstable configuration, we are able to observe the zero-velocity mode brought by $z_{\sigma,2}$ (vertical beam).

7.1. **Discrete waves.** We now conduct simulations on the domain $[-3, 3]$ using 500 grid-points, endowed with periodic boundary conditions. We take $\lambda = 1$ and $c = 4/5$. The initial datum is smooth and given by $u^\circ(x) = \cos(\pi x)^{10} \mathbb{1}_{(0,1/2)}(|x|)$, with non-conserved moments at equilibrium. We consider different relaxation parameters and the results are in Fig. 14. For the case (a), we see the physical wave propagating at speed c and undergoing dispersion (the milky shadow in the direction opposite to propagation). A parasitic wave is brought by $z_{\sigma,1}(\xi)$ propagating at speed $-c$, and experiencing temporal damping, which is slow enough to make it observable using a logarithmic color-scale. For case (b), we do not observe anything except the physical wave, since the relaxation parameters are pretty close to one. This entails a quick damping of the parasitic modes and the scheme behaves essentially as a Lax-Wendroff one. For (c), we see the effect of taking s_v very close to zero. Eventually, configuration (d) is unstable. Here, the spurious mode brought by $z_{\sigma,1}(\xi)$ damps away quickly, whereas the one brought by $z_{\sigma,2}(\xi)$, which does not propagate, is visible for a long time, as s_w is quite close to two.

7.2. **Convergence and order.** Under the same setting as Sec. 7.1 and a final time equal to one, we consider initial data $u^\circ(x) = \cos(\pi x)^{\sigma - 1/2} \mathbb{1}_{(0,1/2)}(|x|)$, so that




FIGURE 15. Empirical order of convergence for the L^2 error at the final time, as function of the smoothness of the initial datum.

$u^\circ \in H^{\sigma-}(\mathbb{R})$ for the Sobolev regularity, and $u^\circ \in B^{\sigma}_{2,\infty}(\mathbb{R})$ for the Besov regularity. We estimate this order by two simulations, one with 2000 grid-points and the other with 4000, yielding the results in Fig. 15. These are in very good agreement with the theoretical behavior, so that the method achieves full second-order accuracy starting from $\sigma \geq 3$. Although we present results only for $s_v = s_w = \frac{11}{10}$, analogous trends of convergence have been obtained for $s_v = \frac{49}{25}$ or $s_v = \frac{1}{25}$, and $s_w = 2 - s_v$.




FIGURE 16. L^2 errors at the final time for the Burgers' equation.

7.3. **Empirical convergence for the Burgers' equation.** If we abandon the setting of a linear problem as (1), by taking non-linear equilibria, we can also

obtain a formally second-order scheme for the inviscid Burgers' equation

$$\partial_t u + \partial_x(\tfrac{1}{2}u^2) = 0.$$

Notice that we do not say anything about its stability, and the scheme can be unstable even if the linearization is stable. By [5, Theorem 3.7], the modified equation of the scheme reads

$$\partial_t u + \partial_x v^{\mathrm{eq}}(u) - \Delta t\Big(\frac{1}{2} - \frac{1}{s_v}\Big)\partial_x\Big(2\partial_x w^{\mathrm{eq}}(u) - \frac{\mathrm{d}v^{\mathrm{eq}}(u)}{\mathrm{d}u}\partial_x v^{\mathrm{eq}}(u)\Big) = \mathcal{O}(\Delta t^2).$$

Formally, we obtain

$$2\partial_x w^{\mathrm{eq}}(u) - \frac{\mathrm{d}v^{\mathrm{eq}}(u)}{\mathrm{d}u}\partial_x v^{\mathrm{eq}}(u) = \Big(2\frac{\mathrm{d}w^{\mathrm{eq}}(u)}{\mathrm{d}u} - \Big(\frac{\mathrm{d}v^{\mathrm{eq}}(u)}{\mathrm{d}u}\Big)^2\Big)\partial_x u$$

so that formal second-order consistency with a conservation law with flux $v^{\mathrm{eq}}(u)$ can be achieved selecting

$$w^{\mathrm{eq}}(u) = \frac{1}{2}\int^u \Big(\frac{\mathrm{d}v^{\mathrm{eq}}(\eta)}{\mathrm{d}\eta}\Big)^2 \mathrm{d}\eta.$$

For the Burgers' equation, this boils down to $v^{\mathrm{eq}}(u) = u^2/2$ and $w^{\mathrm{eq}}(u) = u^3/6$ This scheme is already known in the literature: [14] calls it "Scheme 3" or "D1Q3 Lax-Wendroff".

We now conduct simulations on the domain $[-3, 3]$ endowed with periodic boundary conditions. We take $\lambda = 1$. The initial datum is $u^\circ(x) = \cos(\pi x/4)^{10}\mathbb{1}_{(0,2)}(|x|)$, with non-conserved moments at equilibrium. The final time is $1/2$, which is slightly before the blowup time. Results are provided in Fig. 16 and confirm empirical second-order convergence for all considered values of relaxation parameters. Moreover, we see that the choice $s_v = s_w = \frac{11}{10}$ yields asymptotically smaller errors compared to other choices, in particular against the Lax-Wendroff scheme, obtained when the relaxation is a projection on the equilibrium.

8. Conclusion

In this work, we have conducted a comprehensive stability analysis of a three-velocity lattice Boltzmann scheme for the linear advection equation. By choosing the equilibria to ensure second-order consistency, we have shown that the two relaxation parameters do not affect the accuracy of the scheme, but determine its stability properties. The Lax-Wendroff method is recovered as a particular case, corresponding to unit relaxation parameters, thereby establishing a direct connection between the lattice Boltzmann formulation and a classical finite difference scheme.

Our main contribution is a complete characterization, in terms of the Courant number and the two relaxation parameters, of the conditions under which the amplification matrix is uniformly power-bounded with respect to the Fourier frequency. We derive explicit necessary and sufficient conditions by combining von Neumann analysis with a detailed investigation of the eigenvalue structure of the amplification matrix. In particular, our analysis goes beyond the location of the eigenvalues in the closed unit disk by accounting for their multiplicities and the associated constraints required for power-boundedness. The use of the Cohn-Schur algorithm makes it possible to obtain an explicit characterization of the admissible parameter set, providing a rigorous and systematic account of the stability properties of the scheme.

We have further established convergence of the conserved moment towards the solution of the advection equation for smooth initial data, under equilibrium initialization of the non-conserved moments. The analysis also provides insight into the propagation of discrete waves and the relationship between the lattice Boltzmann dynamics and the underlying macroscopic equation. Numerical experiments support the theoretical findings, illustrate the expected propagation patterns, and confirm second-order convergence for smooth initial data. Additional experiments involving non-smooth data and an adaptation to the inviscid Burgers equation suggest the potential relevance of the proposed framework beyond the linear setting considered in the analysis.

Several directions for future research naturally follow from this work. A first objective is to extend the present analysis to lattice Boltzmann schemes with more velocities and additional relaxation parameters. Further extensions to higher spatial dimensions and more complex target equations would provide a broader framework for investigating the stability properties of lattice Boltzmann methods and their applicability to a wider range of transport phenomena.

Appendix A. Expression of the polynomials

The characteristic polynomial of the amplification matrix $\boldsymbol{G}$ reads

$$\varphi_3(X;\xi,c,t_v,t_w) = \alpha_{3,0} + \alpha_{3,1}X + \alpha_{3,2}X^2 + \alpha_{3,3}X^3, \tag{24}$$

with

$$\begin{aligned}
\alpha_{3,0} &= -t_v t_w,\\
\mathrm{Re}(\alpha_{3,1}) &= t_v t_w - (1-\cos\xi)t_v(1+t_w)c^2 - \cos\xi(t_v+t_w),\\
\mathrm{Im}(\alpha_{3,1}) &= c t_w(1+t_v)\sin\xi,\\
\mathrm{Re}(\alpha_{3,2}) &= -1 + c^2(1+t_w)(1-\cos\xi) + \cos\xi(t_v+t_w),\\
\mathrm{Im}(\alpha_{3,2}) &= c(1+t_v)\sin\xi,\\
\alpha_{3,3} &= 1.
\end{aligned}$$

Applying the Cohn-Schur algorithm on φ_3 yields:

$$\mathscr{C}(\varphi_3) = \varphi_2^\circ(X;\xi,c,t_v,t_w) = \alpha_{2,0}^\circ + \alpha_{2,1}^\circ X + \alpha_{2,2}^\circ X^2, \tag{25}$$

with

$$\begin{aligned}
\mathrm{Re}(\alpha_{2,0}^\circ) &= -c^2 t_v(1-t_w^2)(1-\cos(\xi)) - \cos(\xi)(t_v+t_w)(1-t_v t_w),\\
\mathrm{Im}(\alpha_{2,0}^\circ) &= c t_w(1-t_v^2)\sin(\xi),\\
\mathrm{Re}(\alpha_{2,1}^\circ) &= -(1-t_v^2 t_w^2) + c^2(1+t_w)(1-\cos\xi)(1-t_v^2 t_w)\\
&\qquad + \big(t_v(1-t_w^2) + t_w(1-t_v^2)\big)\cos\xi,\\
\mathrm{Im}(\alpha_{2,1}^\circ) &= c(1-t_v t_w^2)(1+t_v)\sin(\xi),\\
\alpha_{2,2}^\circ &= 1 - t_v^2 t_w^2.
\end{aligned}$$

The first derivative of φ_3 reads:

$$\mathrm{d}\varphi_3/\mathrm{d}X = \varphi_2^\bullet(X;\xi,c,t_v,t_w) = \alpha_{2,0}^\bullet + \alpha_{2,1}^\bullet X + \alpha_{2,2}^\bullet X^2, \tag{26}$$

with

$$\begin{aligned}
\mathrm{Re}(\alpha^{\bullet}_{2,0}) &= t_v t_w - (1-\cos\xi)t_v(1+t_w)c^2 - \cos\xi(t_v+t_w),\\
\mathrm{Im}(\alpha^{\bullet}_{2,0}) &= c t_w(1+t_v)\sin\xi,\\
\mathrm{Re}(\alpha^{\bullet}_{2,1}) &= -2 + 2c^2(1+t_w)(1-\cos\xi) + 2\cos\xi(t_v+t_w),\\
\mathrm{Im}(\alpha^{\bullet}_{2,1}) &= 2c(1+t_v)\sin\xi,\\
\alpha^{\bullet}_{2,2} &= 3.
\end{aligned}$$

Applying the Cohn-Schur algorithm on φ°_2 yields:

$$\mathscr{C}(\varphi^{\circ}_2) = \varphi^{\circ\circ}_1(X;\xi,c,t_v,t_w) = \alpha^{\circ\circ}_{1,0} + \alpha^{\circ\circ}_{1,1}X, \tag{27}$$

with

$$\begin{aligned}
\mathrm{Re}(\alpha^{\circ\circ}_{1,0}) = &-(1-t_vt_w)^2\big((1+t_vt_w)^2 - \cos^2(\xi)(t_v+t_w)^2\big)\\
&+ c^2(1-\cos\xi)\Big[(1-t_v)\big((1-t_vt_w)(1+t_v^2t_w^3) + t_vt_w(t_w-t_v)(1+t_w)\big)\\
&\qquad + \cos\xi(1+t_v)\big((t_v+t_w^2)(1-t_vt_w)^2 + t_vt_w(1-t_w)^2(1+t_v)\big)\Big]\\
&+ c^4(1-\cos\xi)^2(1-t_w)t_v(1+t_w)^2(1-t_v^2t_w),\\
\mathrm{Im}(\alpha^{\circ\circ}_{1,0}) = &\, c\sin\xi(1+t_v)(1+t_w)\Big[(1-t_vt_w)^2(1+t_vt_w-\cos\xi(t_v+t_w))\\
&\qquad - c^2(1-\cos\xi)(t_v(1-t_w)(1-t_vt_w^2) + t_w(1-t_v)(1-t_v^2t_w))\Big],\\
\alpha^{\circ\circ}_{1,1} = &\,(1-t_vt_w)^2\big((1+t_vt_w)^2 - \cos^2(\xi)(t_v+t_w)^2\big)\\
&- c^2(1-\cos^2(\xi))t_w^2(1-t_v^2)^2 - c^4(1-\cos\xi)^2t_v^2(1-t_w^2)^2\\
&- 2c^2\cos\xi(1-\cos\xi)t_v(1-t_w^2)(1-t_vt_w)(t_v+t_w).
\end{aligned}$$

The first derivative of φ°_2 reads:

$$\mathrm{d}\varphi^{\circ}_2/\mathrm{d}X = \varphi^{\circ\bullet}_1(X;\xi,c,t_v,t_w) = \alpha^{\circ\bullet}_{1,0} + \alpha^{\circ\bullet}_{1,1}X, \tag{28}$$

with

$$\begin{aligned}
\mathrm{Re}(\alpha^{\circ\bullet}_{1,0}) &= -(1-t_v^2t_w^2) + c^2(1+t_w)(1-\cos\xi)(1-t_v^2t_w)\\
&\qquad + \big(t_v(1-t_w^2) + t_w(1-t_v^2)\big)\cos\xi,\\
\mathrm{Im}(\alpha^{\circ\bullet}_{1,0}) &= c(1-t_vt_w^2)(1+t_v)\sin(\xi),\\
\alpha^{\circ\bullet}_{1,1} &= 2(1-t_v^2t_w^2).
\end{aligned}$$

Applying the Cohn-Schur algorithm on $\varphi^{\bullet}_2$ yields:

$$\mathscr{C}(\varphi^{\bullet}_2) = \varphi^{\bullet\circ}_1(X;\xi,c,t_v,t_w) = \alpha^{\bullet\circ}_{1,0} + \alpha^{\bullet\circ}_{1,1}X, \tag{29}$$

with

$$\begin{aligned}
\mathrm{Re}(\alpha^{\bullet\circ}_{1,0}) = &-2(1-\cos\xi(t_v+t_w))(3-t_vt_w+\cos\xi(t_v+t_w)) + 2(1-\cos\xi)c^2\big(\\
&\qquad 3(1+t_w) + \cos\xi(t_v+t_w^2)(1+t_v) - t_v(1+t_w)^2 - t_w(1+t_v)^2\\
&\big) + 2c^4t_v(1-\cos\xi)^2(1+t_w)^2,\\
\mathrm{Im}(\alpha^{\bullet\circ}_{1,0}) = &\, 2c\sin\xi(1+t_v)\big(3 + t_w(1+t_v) - (c^2+\cos\xi(1-c^2))(t_v+t_w)(1+t_w)\big),\\
\alpha^{\bullet\circ}_{1,1} = &\, 9 - (\cos\xi(t_v+t_w) - t_vt_w)^2 + (1-\cos\xi)c^2\big(\\
&\qquad 2t_v(1+t_w)(t_wt_v - \cos\xi(t_v+t_w)) - (1+\cos\xi)t_w^2(1+t_v)^2\\
&\big) - c^4(1-\cos\xi)^2(1+t_w)^2t_v^2.
\end{aligned}$$

The first derivative of $\varphi_2^\bullet$ reads:

$$\mathrm{d}\varphi_2^\bullet/\mathrm{d}X = \varphi_1^{\bullet\bullet}(X;\xi,c,t_v,t_w) = \alpha_{1,0}^{\bullet\bullet} + \alpha_{1,1}^{\bullet\bullet} X, \tag{30}$$

with

$$\begin{aligned}
\mathrm{Re}(\alpha_{1,0}^{\bullet\bullet}) &= -2 + 2c^2(1+t_w)(1-\cos\xi) + 2\cos\xi(t_v+t_w),\\
\mathrm{Im}(\alpha_{1,0}^{\bullet\bullet}) &= 2c(1+t_v)\sin\xi,\\
\alpha_{1,1}^{\bullet\bullet} &= 6.
\end{aligned}$$

References

[1] Denise Aregba-Driollet. Convergence of Lattice Boltzmann methods with overrelaxation for a nonlinear conservation law. *ESAIM: Mathematical Modelling and Numerical Analysis*, 58(5):1935–1958, 2024.

[2] Denise Aregba-Driollet and Thomas Bellotti. Monotonicity and Convergence of Two-Relaxation-Times Lattice Boltzmann Schemes for a Nonlinear Conservation Law. *SIAM Journal on Numerical Analysis*, 64(1):251–276, 2026.

[3] Mapundi K Banda, Wen-An Yong, and Axel Klar. A stability notion for lattice Boltzmann equations. *SIAM Journal on Scientific Computing*, 27(6):2098–2111, 2006.

[4] Thomas Bellotti. Monotonicity for genuinely multi-step methods: results and issues from a simple lattice Boltzmann scheme. In *International Conference on Finite Volumes for Complex Applications*, pages 33–41. Springer, 2023.

[5] Thomas Bellotti. Truncation errors and modified equations for the lattice boltzmann method via the corresponding finite difference schemes. *ESAIM: Mathematical Modelling and Numerical Analysis*, 57(3):1225–1255, 2023.

[6] Thomas Bellotti. Initialisation from lattice Boltzmann to multi-step Finite Difference methods: modified equations and discrete observability. *Journal of Computational Physics*, 504:112871, 2024.

[7] Thomas Bellotti. The influence of parasitic modes on stable lattice Boltzmann schemes and weakly unstable multi-step Finite Difference schemes. *Computers & Mathematics with Applications*, 174:397–416, 2024.

[8] Thomas Bellotti. Stability of lattice Boltzmann schemes for initial boundary value problems in raw formulation. *arXiv preprint arXiv:2507.13108*, 2025.

[9] Thomas Bellotti, Benjamin Graille, and Marc Massot. Finite Difference formulation of any lattice Boltzmann scheme. *Numerische Mathematik*, 152(1):1–40, September 2022.

[10] Philip Brenner, Vidar Thomée, and Lars B Wahlbin. *Besov spaces and applications to difference methods for initial value problems*, volume 434. Springer, 2006.

[11] James E Broadwell. Shock structure in a simple discrete velocity gas. *The Physics of Fluids*, 7(8):1243–1247, 1964.

[12] Filipa Caetano, François Dubois, and Benjamin Graille. A result of convergence for a mono-dimensional two-velocities lattice Boltzmann scheme. *Discrete and Continuous Dynamical Systems - S*, 17(11):3129–3154, 2024.

[13] Torsten Carleman. *Problèmes mathématiques dans la théorie cinétique des gaz*. Number 2 in Publications scientifiques de l'Institut Mittag-Leffler. Almqvist & Wiksell, Uppsala, 1957.

[14] David Coulette, Clémentine Courtès, Emmanuel Franck, and Laurent Navoret. Vectorial kinetic relaxation model with central velocity. application to implicit relaxations schemes. *Communications in Computational Physics*, 27(4), 2020.

[15] Jean-François Coulombel. Stability of finite difference schemes for hyperbolic initial boundary value problems, 2013. HCDTE Lecture Notes. Part I. Nonlinear Hyperbolic PDEs, Dispersive and Transport Equations.

[16] Stéphane Dellacherie. Construction and analysis of lattice Boltzmann methods applied to a 1D convection-diffusion equation. *Acta Applicandae Mathematicae*, 131(1):69–140, 2014.

[17] Dominique d'Humières and Irina Ginzburg. Viscosity independent numerical errors for Lattice Boltzmann models: From recurrence equations to "magic" collision numbers. *Computers & Mathematics with Applications*, 58(5):823–840, September 2009.

[18] François Dubois. Equivalent partial differential equations of a lattice Boltzmann scheme. *Computers & Mathematics with Applications*, 55(7):1441–1449, 2008.

[19] François Dubois. Nonlinear fourth order Taylor expansion of lattice Boltzmann schemes. *Asymptotic Analysis*, 127(4):297–337, 2022.

[20] François Dubois, Benjamin Graille, and S. V. Raghurama Rao. A notion of non-negativity preserving relaxation for a mono-dimensional three velocities scheme with relative velocity. *Journal of Computational Science*, 47:101181, November 2020.

[21] Irina Ginzburg. Consistent lattice Boltzmann schemes for the Brinkman model of porous flow and infinite Chapman-Enskog expansion. *Physical Review E—Statistical, Nonlinear, and Soft Matter Physics*, 77(6):066704, 2008.

[22] Irina Ginzburg, Dominique d'Humières, and Alexander Kuzmin. Optimal stability of advection-diffusion lattice Boltzmann models with two relaxation times for positive/negative equilibrium. *Journal of Statistical Physics*, 139(6):1090–1143, 2010.

[23] Anne Greenbaum, Ren-cang Li, and Michael L Overton. First-order perturbation theory for eigenvalues and eigenvectors. *SIAM review*, 62(2):463–482, 2020.

[24] Zhaoli Guo and Chang Shu. *Lattice Boltzmann method and its applications in engineering*, volume 3. World Scientific, 2013.

[25] Michael Junk and Zhaoxia Yang. Convergence of lattice Boltzmann methods for Navier–Stokes flows in periodic and bounded domains. *Numerische Mathematik*, 112(1):65–87, 2009.

[26] Michael Junk and Wen-An Yong. Weighted L^2-Stability of the Lattice Boltzmann Method. *SIAM Journal on Numerical Analysis*, 47(3):1651–1665, 2009.

[27] Tosio Kato. *Perturbation theory for linear operators*. Springer Science & Business Media, 1995.

[28] A Kuzmin, Irina Ginzburg, and AA Mohamad. The role of the kinetic parameter in the stability of two-relaxation-time advection–diffusion lattice Boltzmann schemes. *Computers & Mathematics with Applications*, 61(12):3417–3442, 2011.

[29] Pierre Lallemand and Li-Shi Luo. Theory of the lattice Boltzmann method: Dispersion, dissipation, isotropy, Galilean invariance, and stability. *Physical Review E*, 61:6546–6562, Jun 2000.

[30] Peter Lax and Burton Wendroff. Systems of conservation laws. *Communications on Pure and Applied Mathematics*, 13(2):217–237, 1960.

[31] Morris Marden. *Geometry of Polynomials*. Number 3 in Mathematical surveys and monographs. American Mathematical Society, 2nd edition, 1966.

[32] John J. H. Miller. On the Location of Zeros of Certain Classes of Polynomials with Applications to Numerical Analysis. *IMA Journal of Applied Mathematics*, 8(3):397–406, 1971.

[33] Yue-Hong Qian, Dominique d'Humières, and Pierre Lallemand. Lattice BGK models for Navier-Stokes equation. *EPL (Europhysics Letters)*, 17(6):479–484, 1992.

[34] Martin Rheinländer. On the stability structure for lattice Boltzmann schemes. *Computers & Mathematics with Applications*, 59(7):2150–2167, 2010.

[35] Eugène Rouché. Mémoire sur la série de Lagrange. *Journal de l'École Impériale Polytechnique*, 39:193–224, 1862.

[36] James D Sterling and Shiyi Chen. Stability analysis of lattice Boltzmann methods. *Journal of Computational Physics*, 123(1):196–206, 1996.

[37] John C. Strikwerda. *Finite difference schemes and partial differential equations*. Society for Industrial and Applied Mathematics, Philadelphia, 2nd ed edition, 2004.

[38] Sauro Succi. *The lattice Boltzmann equation: for fluid dynamics and beyond*. Oxford University Press, 2001.

[39] The Sage Developers. *SageMath, the Sage Mathematics Software System (Version 9.7.beta7)*, 2022. `https://www.sagemath.org`.