

Pour se remettre dans le bain

## Pour se remettre dans le bain

1. Citer les quatre estimateurs bayésiens les plus couramment utilisés.
2. Qu'est-ce qu'une région de crédibilité ?
3. Comment prédire quand on est bayésien(ne) ?
4. Décrire l'approche quasi-bayésienne.
5. Pourquoi l'approche quasi-bayésienne peut-elle être vue comme une généralisation de l'apprentissage bayésien ?
6. Illustrer la provenance du quasi-posterior au moyen d'une formulation variationnelle.
7. Citer les quatre estimateurs quasi-bayésiens les plus couramment utilisés.
8. Rappeler le principe de l'ERM et de l'agrégation à poids convexe (EWA). Quel est le lien entre EWA et apprentissage quasi-bayésien ?

## Assessing the performance: the oracle approach

Oracle:

$$\phi^* \in \arg \min_{\phi \in \mathcal{Y}^x} R(\phi).$$

Ultimate goal: do almost as well as the oracle.

# Assessing the performance: the oracle approach

Oracle:

$$\phi^* \in \arg \min_{\phi \in \mathcal{Y}^x} R(\phi).$$

Ultimate goal: do almost as well as the oracle.

Excess risk:

$$\mathcal{E}(\cdot) = R(\cdot) - R^* \geq 0, \quad R^* = R(\phi^*).$$

# PAC oracle inequalities

## PAC oracle inequalities

Let  $R^*$  denote the oracle risk. For any  $\epsilon > 0$ ,

$$\mathbb{P} \left( R(\hat{\phi}_\lambda) - R^* \leq \spadesuit \inf_{\phi \in \mathcal{F}} \left\{ R(\phi) - R^* + \frac{\Delta(\phi, \epsilon)}{n^\alpha} \right\} \right) \geq 1 - \epsilon,$$

where  $\spadesuit \geq 1$  and  $\lambda \propto n$ . If  $\spadesuit = 1$ , the inequality is *exact* or *sharp*.

## PAC oracle inequalities

Let  $R^*$  denote the oracle risk. For any  $\epsilon > 0$ ,

$$\mathbb{P} \left( R(\hat{\phi}_\lambda) - R^* \leq \spadesuit \inf_{\phi \in \mathcal{F}} \left\{ R(\phi) - R^* + \frac{\Delta(\phi, \epsilon)}{n^\alpha} \right\} \right) \geq 1 - \epsilon,$$

where  $\spadesuit \geq 1$  and  $\lambda \propto n$ . If  $\spadesuit = 1$ , the inequality is *exact* or *sharp*.

Key argument: concentration inequalities (e.g., Bernstein, Hoeffding) + Legendre transform of the Kullback-Leibler divergence.

## PAC oracle inequalities

Let  $R^*$  denote the oracle risk. For any  $\epsilon > 0$ ,

$$\mathbb{P} \left( R(\hat{\phi}_\lambda) - R^* \leq \spadesuit \inf_{\phi \in \mathcal{F}} \left\{ R(\phi) - R^* + \frac{\Delta(\phi, \epsilon)}{n^\alpha} \right\} \right) \geq 1 - \epsilon,$$

where  $\spadesuit \geq 1$  and  $\lambda \propto n$ . If  $\spadesuit = 1$ , the inequality is *exact* or *sharp*.

Key argument: concentration inequalities (e.g., Bernstein, Hoeffding) + Legendre transform of the Kullback-Leibler divergence.

Typical rates in the literature

- ▶  $\alpha = \frac{1}{2}$  (slow rate)
- ▶  $\alpha = 1$  (fast rate)



## PAC oracle inequalities

Let  $R^*$  denote the oracle risk. For any  $\epsilon > 0$ ,

$$\mathbb{P} \left( R(\hat{\phi}_\lambda) - R^* \leq \spadesuit \inf_{\phi \in \mathcal{F}} \left\{ R(\phi) - R^* + \frac{\Delta(\phi, \epsilon)}{n^\alpha} \right\} \right) \geq 1 - \epsilon,$$

where  $\spadesuit \geq 1$  and  $\lambda \propto n$ . If  $\spadesuit = 1$ , the inequality is *exact* or *sharp*.

Key argument: concentration inequalities (e.g., Bernstein, Hoeffding) + Legendre transform of the Kullback-Leibler divergence.

Typical rates in the literature

- ▶  $\alpha = \frac{1}{2}$  (slow rate)
- ▶  $\alpha = 1$  (fast rate)

Let  $d = \dim(\mathcal{X})$

- ▶  $\Delta(\phi, \epsilon) \propto d + \log \frac{1}{\epsilon}$
- ▶  $\Delta(\phi, \epsilon) \propto \log d + \log \frac{1}{\epsilon}$

## PAC oracle inequalities

Let  $R^*$  denote the oracle risk. For any  $\epsilon > 0$ ,

$$\mathbb{P} \left( R(\hat{\phi}_\lambda) - R^* \leq \spadesuit \inf_{\phi \in \mathcal{F}} \left\{ R(\phi) - R^* + \frac{\Delta(\phi, \epsilon)}{n^\alpha} \right\} \right) \geq 1 - \epsilon,$$

where  $\spadesuit \geq 1$  and  $\lambda \propto n$ . If  $\spadesuit = 1$ , the inequality is *exact* or *sharp*.

Key argument: concentration inequalities (e.g., Bernstein, Hoeffding) + Legendre transform of the Kullback-Leibler divergence.

Typical rates in the literature

▶  $\alpha = \frac{1}{2}$  (slow rate)

▶  $\alpha = 1$  (fast rate)

Let  $d = \dim(\mathcal{X})$

▶  $\Delta(\phi, \epsilon) \propto d + \log \frac{1}{\epsilon}$

▶  $\Delta(\phi, \epsilon) \propto \log d + \log \frac{1}{\epsilon}$

The remainder term grows with  $d$  and the size of  $\mathcal{F}$ . It decreases with  $n$ .

## Hoeffding inequality

Let  $V_1, \dots, V_n$  be independent real-valued random variables such that  $a_i \leq V_i \leq b_i$  a.s. Let  $\bar{V}_n = \frac{1}{n} \sum_{i=1}^n V_i$ .

# Hoeffding inequality

Let  $V_1, \dots, V_n$  be independent real-valued random variables such that  $a_i \leq V_i \leq b_i$  a.s. Let  $\bar{V}_n = \frac{1}{n} \sum_{i=1}^n V_i$ . Then

$$\mathbb{P}(\bar{V}_n - \mathbb{E}\bar{V}_n > t) \leq \exp\left(-\frac{2n^2 t^2}{\sum_{i=1}^n (b_i - a_i)^2}\right), \forall t > 0.$$

### Lemma (Csiszar, 1975 ; Catoni, 2004)

*Let  $(A, \mathcal{A})$  be a measurable space. For any probability  $\mu$  on  $(A, \mathcal{A})$  and any measurable function  $h : A \rightarrow \mathbb{R}$  such that  $\int (\exp \circ h) d\mu < \infty$ ,*

$$\log \int (\exp \circ h) d\mu = \sup_{m \in \mathcal{M}_\pi(A, \mathcal{A})} \left\{ \int h dm - \mathcal{K}(m, \mu) \right\},$$

*with the convention  $\infty - \infty = -\infty$ . Moreover, as soon as  $h$  is upper-bounded on the support of  $\mu$ , the supremum with respect to  $m$  on the right-hand side is reached for the Gibbs distribution  $g$  given by*

$$\frac{dg}{d\mu}(a) = \frac{\exp \circ h(a)}{\int (\exp \circ h) d\mu}, \quad a \in A.$$

# The PAC-Bayesian theory

# The PAC-Bayesian theory

...consists in producing PAC inequalities of quasi-Bayesian estimators.

# The PAC-Bayesian theory

...consists in producing PAC inequalities of quasi-Bayesian estimators.

▣ Shawe-Taylor and Williamson (1997). A PAC analysis of a Bayes estimator, *COLT*

▣ McAllester (1998). Some PAC-Bayesian theorems, *COLT*

▣ McAllester (1999). PAC-Bayesian model averaging, *COLT*

▣ Catoni (2004). Statistical Learning Theory and Stochastic Optimization, Springer

▣ Audibert (2004). Une approche PAC-bayésienne de la théorie statistique de l'apprentissage, *Ph.D. thesis*,  
*Université Pierre & Marie Curie*

▣ Catoni (2007). PAC-Bayesian Supervised Classification: The Thermodynamics of Statistical Learning, IMS

▣ Dalalyan and Tsybakov (2008). Aggregation by exponential weighting, sharp PAC-Bayesian bounds and sparsity,  
*Machine Learning*

▣ Alquier and Lounici (2011). PAC-Bayesian theorems for sparse regression estimation with exponential weights,  
*Electronic Journal of Statistics*



## A flexible and powerful framework

# A flexible and powerful framework

Numerous models addressed by the PAC-Bayesian literature.

▣ Alquier and Wintenberger (2012). Model selection for weakly dependent time series forecasting, *Bernoulli*

▣ Seldin, Lavolette, Cesa-Bianchi, Shawe-Taylor and Auer (2012). PAC-Bayesian inequalities for martingales, *IEEE Transactions on Information Theory*

▣ Alquier and Biau (2013). Sparse Single-Index Model, *Journal of Machine Learning Research*

▣ Guedj and Alquier (2013). PAC-Bayesian Estimation and Prediction in Sparse Additive Models, *Electronic Journal of Statistics*

▣ Guedj and Robbiano (2017). PAC-Bayesian High Dimensional Bipartite Ranking, *Journal of Statistical Planning and Inference*

▣ Alquier and Guedj (2017). An Oracle Inequality for Quasi-Bayesian Non-Negative Matrix Factorization, *Mathematical Methods of Statistics*

▣ Li, Guedj and Loustau (2018). A Quasi-Bayesian Perspective to Online Clustering, *arXiv preprint*

# A flexible and powerful framework

Numerous models addressed by the PAC-Bayesian literature.

- Alquier and Wintenberger (2012). Model selection for weakly dependent time series forecasting, *Bernoulli*
- Seldin, Laviolette, Cesa-Bianchi, Shawe-Taylor and Auer (2012). PAC-Bayesian inequalities for martingales, *IEEE Transactions on Information Theory*
- Alquier and Biau (2013). Sparse Single-Index Model, *Journal of Machine Learning Research*
- Guedj and Alquier (2013). PAC-Bayesian Estimation and Prediction in Sparse Additive Models, *Electronic Journal of Statistics*
- Guedj and Robbiano (2017). PAC-Bayesian High Dimensional Bipartite Ranking, *Journal of Statistical Planning and Inference*
- Alquier and Guedj (2017). An Oracle Inequality for Quasi-Bayesian Non-Negative Matrix Factorization, *Mathematical Methods of Statistics*
- Li, Guedj and Loustau (2018). A Quasi-Bayesian Perspective to Online Clustering, *arXiv preprint*

Towards (almost) no assumptions to derive powerful results

- Bégin, Germain, Laviolette and Roy (2016). PAC-Bayesian bounds based on the Rényi divergence, *AISTATS*
- Alquier and Guedj (2017). Simpler PAC-Bayesian bounds for hostile data, *Machine Learning*

# Sampling

# Monte Carlo integration

Objective: approximation of an integral

$$\mathcal{J} = \int h(x)f(x)dx.$$

# Monte Carlo integration

Objective: approximation of an integral

$$\mathcal{J} = \int h(x)f(x)dx.$$

**Key idea:** exploit the fact that  $\mathcal{J} = \mathbb{E}_{X \sim f}[h(X)]$ .

# Monte Carlo principle

# Monte Carlo principle

Sample a sequence  $x_1, \dots, x_m \sim f$ .



# Monte Carlo principle

Sample a sequence  $x_1, \dots, x_m \sim f$ .

Then use

$$\hat{\mathcal{J}}_m = \frac{1}{m} \sum_{i=1}^m h(x_i)$$

as an estimator of  $\mathcal{J}$ .

## Monte Carlo principle

Sample a sequence  $x_1, \dots, x_m \sim f$ .

Then use

$$\hat{\mathcal{J}}_m = \frac{1}{m} \sum_{i=1}^m h(x_i)$$

as an estimator of  $\mathcal{J}$ .

Justification: by the Strong Law of Large Numbers,

$$\hat{\mathcal{J}}_m \rightarrow \mathcal{J}.$$

# Approximation evaluation

## Approximation evaluation

Estimate the variance with

$$\nu_m = \frac{1}{m} \frac{1}{m-1} \sum_{i=1}^m (h(x_i) - \hat{\mathcal{J}}_m)^2,$$

## Approximation evaluation

Estimate the variance with

$$\nu_m = \frac{1}{m} \frac{1}{m-1} \sum_{i=1}^m (h(x_i) - \hat{\mathcal{J}}_m)^2,$$

and recall that for  $m$  large,

$$\frac{\hat{\mathcal{J}}_m - \mathbb{E}[h(X)]}{\sqrt{\nu_m}} \approx \mathcal{N}(0, 1).$$

# Importance sampling

# Importance sampling

Simulating from  $f$  might not be the best idea: difficult/impossible, not optimal, ...

# Importance sampling

Simulating from  $f$  might not be the best idea: difficult/impossible, not optimal, ...

An alternative to direct sampling is importance sampling, with the following trick:



# Importance sampling

Simulating from  $f$  might not be the best idea: difficult/impossible, not optimal, ...

An alternative to direct sampling is importance sampling, with the following trick:

$$\begin{aligned}\mathbb{E}_{X \sim f}[h(X)] &= \int h(x)f(x)dx = \int h(x)\frac{f(x)}{g(x)}g(x)dx \\ &= \mathbb{E}_{X \sim g}\left[h(X)\frac{f(X)}{g(X)}\right],\end{aligned}$$

which allows us to use other distributions.

# Importance sampling

# Importance sampling

Sample a sequence  $x_1, \dots, x_m \sim g$ .

# Importance sampling

Sample a sequence  $x_1, \dots, x_m \sim g$ .

Then use

$$\hat{\mathcal{J}}_m = \frac{1}{m} \sum_{i=1}^m \frac{f(x_i)}{g(x_i)} h(x_i)$$

as an estimator of  $\mathcal{J}$ .

Justification: by the Strong Law of Large Numbers,

$$\hat{\mathcal{J}}_m \rightarrow \mathcal{J}.$$

## Justification

1. Converges for any choice of the distribution  $g$  as long as the support of  $g$  contains the support of  $f$ .

# Justification

1. Converges for any choice of the distribution  $g$  as long as the support of  $g$  contains the support of  $f$ .
2. Instrumental distribution  $g$  may be chosen among distributions easy to simulate.

## Justification

1. Converges for any choice of the distribution  $g$  as long as the support of  $g$  contains the support of  $f$ .
2. Instrumental distribution  $g$  may be chosen among distributions easy to simulate.
3. The same sample generated from  $g$  can be used repeatedly, not only for different functions  $h$  but also for different densities  $f$ .

## Choice of importance function



## Choice of importance function

The optimal choice is

$$g^*(x) = \frac{|h(x)|f(x)}{\int |h(x)|f(x)dx},$$

but the integral is unknown (obviously).

## Choice of importance function

The optimal choice is

$$g^*(x) = \frac{|h(x)|f(x)}{\int |h(x)|f(x)dx},$$

but the integral is unknown (obviously).

In practice, pick a density  $g$  which is close enough to  $|h|f$  and for which  $|h|f/g$  is bounded.

## Choice of importance function

The optimal choice is

$$g^*(x) = \frac{|h(x)|f(x)}{\int |h(x)|f(x)dx},$$

but the integral is unknown (obviously).

In practice, pick a density  $g$  which is close enough to  $|h|f$  and for which  $|h|f/g$  is bounded.

Beware: importance sampling suffers from the curse of dimensionality.

## Choice of importance function

The optimal choice is

$$g^*(x) = \frac{|h(x)|f(x)}{\int |h(x)|f(x)dx},$$

but the integral is unknown (obviously).

In practice, pick a density  $g$  which is close enough to  $|h|f$  and for which  $|h|f/g$  is bounded.

Beware: importance sampling suffers from the curse of dimensionality.

If  $\sup f/g = M < +\infty$ , the accept-reject algorithm may be used.

## Choice of importance function

The optimal choice is

$$g^*(x) = \frac{|h(x)|f(x)}{\int |h(x)|f(x)dx},$$

but the integral is unknown (obviously).

In practice, pick a density  $g$  which is close enough to  $|h|f$  and for which  $|h|f/g$  is bounded.

Beware: importance sampling suffers from the curse of dimensionality.

If  $\sup f/g = M < +\infty$ , the accept-reject algorithm may be used.

The instrumental function may be  $\pi$  (the prior). But often inefficient if data informative, and impossible if  $\pi$  is improper...

# Sampling random variables

The fundamental theorem of sampling:

## Sampling random variables

The fundamental theorem of sampling: if  $U \sim \mathcal{U}(0, 1)$ , let  $F$  denote a CDF, then  $F^{-1}(U)$  is a random variable with distribution  $dF$ .

## Sampling random variables

The fundamental theorem of sampling: if  $U \sim \mathcal{U}(0, 1)$ , let  $F$  denote a CDF, then  $F^{-}(U)$  is a random variable with distribution  $dF$ .

In practice, this theorem has a very limited scope since the pseudo-inverse  $F^{-}$  is usually unknown/not analytically tractable.



## Accept-reject

Goal: sample  $x \sim f$ .

# Accept-reject

Goal: sample  $x \sim f$ .

Ingredients: a density  $g$  such that

- ▶  $f/g$  is upper-bounded by  $M$  (known)
- ▶ sampling from  $g$  is easy

# Accept-reject

Goal: sample  $x \sim f$ .

Ingredients: a density  $g$  such that

- ▶  $f/g$  is upper-bounded by  $M$  (known)
- ▶ sampling from  $g$  is easy

The algorithm:

1. Sample  $z \sim g$  and  $u \sim \mathcal{U}(0, Mg(z))$
2. If  $u \leq f(z)$ , take  $x = z$ , otherwise go back to 1.

## How should we choose $g$ ?

The mean number of samples to obtain 1 realization is  $M$ !  
Efficient algorithm =  $M$  close to 1.

## How should we choose $g$ ?

The mean number of samples to obtain 1 realization is  $M$ !  
Efficient algorithm =  $M$  close to 1.

In practice we pick a  $g$  similar to  $f$ .

## How should we choose $g$ ?

The mean number of samples to obtain 1 realization is  $M$ !  
Efficient algorithm =  $M$  close to 1.

In practice we pick a  $g$  similar to  $f$ .

Nice fact: no need to know the normalizing constant of  $f$ !