

# LINEAR REGRESSION MODELS W4315

## HOMEWORK 1 QUESTIONS

January 26, 2010

Instructor: Frank Wood

**1. (20 points)** Let  $Y_i = \beta_0 + \beta_1 X_i + \epsilon_i$  be a linear regression model with distribution of error terms unspecified (but with mean  $E(\epsilon) = 0$  and variance  $V(\epsilon_i) = \sigma^2$  ( $\sigma^2$  finite) known). Show that  $s^2 = MSE = \frac{\sum(Y_i - \hat{Y}_i)^2}{n-2}$  is an unbiased estimator for  $\sigma^2$ .  $\hat{Y}_i = b_0 + b_1 X_i$  where  $b_0 = \bar{Y} - b_1 \bar{X}$  and  $b_1 = \frac{\sum_i((X_i - \bar{X})(Y_i - \bar{Y}))}{\sum_i(X_i - \bar{X})^2}$

**2. (20 points)** Derive the maximum likelihood estimators  $\hat{\beta}_0, \hat{\beta}_1$ , and  $\hat{\sigma}^2$  for parameters  $\beta_0, \beta_1$ , and  $\sigma^2$  for the normal linear regression model (i.e.  $\epsilon_i \sim_{iid} N(0, \sigma^2)$ ).

**3. (10 points)** File "unif.txt" (you can find in on professor Wood's website) contains 200 numbers randomly generated from uniform distribution  $U(2,5)$ . Read these numbers into MATLAB (using command say, "textread" or simply copy paste from the .txt file) and do the followings:

- Take these 200 number as a population. Use command 'randsample' of MATLAB, draw 100 numbers out of this population randomly with replacement and plot one histogram of these 100 numbers.
- Use command 'rand' in MATLAB, draw 100 samples directly from  $U(2,5)$  and plot another histogram of these numbers.
- Compare the two histograms, what can you say about the difference between the distribution of samples from the population and from the uniform distribution itself? For the convenience of comparison, you may want to overlay two histograms onto one graph and to see if any apparent difference.

**4. (10 points)** File "normal.txt" (professor's website) contain 200 randomly generated numbers from Normal distribution  $N(-1, 2^2)$ . Like in problem 3, do the followings:

- Take these 200 number as a population. Use command 'randsample' of MATLAB, draw 100 numbers out of this population randomly with replacement and plot a histogram of these 100 numbers.
- Use command 'rand' in MATLAB, draw 100 samples directly from  $N(-1, 2^2)$  and plot a histogram of these numbers.
- Compare the two histograms, can you get the similar conclusion as that of problem 3?

**5. (40 points) Copier maintenance.**<sup>1</sup> The Tri-City Office Equipment Corporation sells an imported copier on a franchise basis and performs preventive maintenance and repair services on this copier. The data below have been collected from 45 recent calls on users to perform routine preventive number of minutes spent by the service person. Assume that first-order regression model( $Y_i = b_0 + b_1X_i + \epsilon_i$ ) is appropriate.

i:	1	2	3	...	43	44	45
$X_i$	2	4	3	...	2	4	5
$Y_i$	20	60	46	...	27	61	77

- Obtain estimated regression function.
- Plot the estimated regression function and the data. How well does the estimated regression function fit the data?
- interpret  $b_0$  in your estimated regression function. Does  $b_0$  provide any relevant information here? Explain.
- Obtain a point estimate of the mean service time when  $X = 5$  copiers are serviced.

Notice: You can get data of this problem on [www.mhhe.com/KutnerALRM4e](http://www.mhhe.com/KutnerALRM4e). Use MATLAB to finish this problem, do not use any other program language. Only basic MATLAB commands are allowed, do not use any built-in command to run regression automatically, e.g. 'regress' in MATLAB is forbidden.

---

<sup>1</sup>This is problem 1.20 in "Applied Linear Regression Models(4th edition)" by Kutner etc. )