

Problem 1

- (a) Prove $-(-x) = x$.
- (b) Prove $-(xy) = (-x)y$.

Solution**Part (a)**

$$0 + -(-x) = -(-x) \quad (\text{A3})$$

$$[x + -x] + -(-x) = -(-x) \quad (\text{A4})$$

$$x + [-x + -(-x)] = -(-x) \quad (\text{A2})$$

$$x + 0 = -(-x) \quad (\text{A4})$$

$$x = -(-x) \quad (\text{A3})$$

Part (b)

$$(-x)y + xy = (-x + x)y \quad (\text{D})$$

$$(-x)y + xy = (0)y \quad (\text{A4})$$

In class it was proved that $0 \cdot x = 0$ for all x . By this result I get

$$(-x)y + xy = 0$$

$$(-x)y + xy + -(xy) = 0 + -(xy) \quad \text{add } -(xy)$$

$$(-x)y + 0 = 0 + -(xy) \quad (\text{A4})$$

$$(-x)y = -(xy) \quad (\text{A3})$$

Problem 2

- (a) Prove if $x > y, z < 0$ then $xz < yz$.
- (b) Prove if $x > y > 0, z > w > 0$ then $xz > yw$.
- (c) Prove if $x > 0$ then $x^{-1} > 0$.

Solution**Part (a)**

In order to prove this I will first prove consequent 7 introduced in class, that $(-x) \cdot (y) = (-xy) = x \cdot (-y)$.

$$x + [(-1) \cdot x] = [1 \cdot x] + [(-1) \cdot x] \quad (\text{M3})$$

$$x + [(-1) \cdot x] = (1 + (-1)) \cdot x \quad (\text{D})$$

$$x + [(-1) \cdot x] = 0 \cdot x \quad (\text{A4})$$

$$x + [(-1) \cdot x] = 0 \quad (\text{consequent 3})$$

$$-x + x + [(-1) \cdot x] = -x + 0$$

$$0 + [(-1) \cdot x] = -x + 0 \quad (\text{A4})$$

$$[(-1) \cdot x] = -x \quad (\text{A3})$$

Using the equivalence established above;

$$(-x)(y) = (-1 \cdot x) \cdot (y)$$

$$(-x)(y) = -1 \cdot (x \cdot y) \quad (\text{D})$$

$$(-x)(y) = -(xy)$$

This shows that $(-x)(y) = -(xy)$. The argument that $-(xy) = (x)(-y)$ has an identical structure.

$z < 0$ and $-1 < 0$ so

$$0 \cdot -1 < z \cdot (-1) \quad (\text{O7})$$

$$0 < z \cdot (-1) \quad \text{consequent 3 proved in class}$$

$$0 < -(z \cdot 1) \quad \text{consequent 7}$$

$$0 < -z \quad \text{M3}$$

$$-z > 0 \quad \text{definition of } >$$

Having $-z > 0$ and $x > y$ I use (O6) to get $x(-z) > y(-z)$

$$-(xz) > -(yz) \quad \text{consequent 7}$$

$$-(xz) + ((xz) + (yz)) > -(yz) + ((xz) + (yz)) \quad (\text{O4})$$

$$-(xz) + ((xz) + (yz)) > -(yz) + ((yz) + (xz)) \quad (\text{A1})$$

$$(-(xz) + (xz)) + (yz) > (-(yz) + (yz)) + (xz) \quad (\text{M2})$$

$$0 + (yz) > 0 + (xz) \quad (\text{A4})$$

$$(yz) > (xz) \quad (\text{A3})$$

By the definition of $>$ this is the same as saying $xz < yz$.

Part (b)

It is given that $z > w$ and $w > 0$. By (O2) $z > 0$. It is also given that $x > y$. Then by (O6) $xz > yz$. It is also given that $y > 0$. By (O6) again $zy > wy$. Then by (M1) $yz > yw$ and finally by (O2) $xz > yw$.

Part (c)

First assume that $x^{-1} < 0$. Then by (O7) proved in class:

$$x \cdot x^{-1} < 0 \cdot x^{-1}$$

It was also proved in class that $0 \cdot x = 0$ for all x . Thus,

$$\begin{aligned} x \cdot x^{-1} &< 0 \\ 1 &< 0 \end{aligned} \qquad \text{by (M4)}$$

This is a contradiction with (O1) because I know that $1 > 0$, so $x^{-1} > 0$.

Problem 3

Prove that there does not exist an $x \in \mathbb{Z}$ such that $0 < x < 1$.

$$\mathbb{Z} = \{x \in \mathbb{R} \mid x \in \mathbb{N} \vee x = 0 \vee -x \in \mathbb{N}\}.$$

Solution

Consider any arbitrary $x \in \mathbb{R}$. There are three possible cases.

(a) Case 1: $x \in \mathbb{N}$

It was proven in class that for all x in \mathbb{N} , $x \geq 1$. Thus it is impossible that $x < 1$.

(b) Case 2: $x = 0$

If $x = 0$ then it is impossible that $x > 0$.

(c) Case 3: $-x \in \mathbb{N}$

By the same fact used in case 1, $-x \geq 1 \implies x \leq -1$. So it is impossible that $x > 0$.

There is no case in which it is possible that $0 < x < 1$.

Problem 4

Prove that it is impossible to define inequalities in \mathbb{C} such that (O1)-(O4) hold.

Solution

The proof given in the book that for any nonzero $a \in \mathbb{R}$, $a^2 > 0$ depends only on axioms (O1)-(O4). Thus if these axioms held in \mathbb{C} then it would have to be the case that the square of any nonzero element of \mathbb{C} was greater than 0. However, i is defined such that $i^2 = -1$. Using the fact introduced in class that $1 > 0$ I can say

$$\begin{aligned} 1 + (-1) &> 0 + (-1) \\ 0 &> -1 \end{aligned} \qquad \text{(A4)}$$

By axiom (O1) it is impossible for it also to be the case that $0 < -1$. Thus this is a contradiction. Therefore it is impossible to define inequalities in \mathbb{C} in such a way that axioms (O1)-(O4) hold.

Problem 5

- (a) Let $x, y \in \mathbb{R}$. Prove $x \leq y$ if and only if $x - \epsilon < y + \epsilon \forall \epsilon > 0$.
- (b) Let $x, y \in \mathbb{R}$ with $x < y$. Prove there exists $z \in \mathbb{R}$ with $x < z < y$.
- (c) Let $a, x, b \in \mathbb{R}$ with $a < x < b$. Prove there exists $\epsilon > 0$ such that $a < x - \epsilon < x + \epsilon < b$.
Deduce that $(x - \epsilon, x + \epsilon) \subset (a, b)$.

Solution**Part (a)**

First let $x \leq y$. By part (i) of theorem 1.9 proved in the textbook I know that $x < y + \epsilon$ for all $\epsilon > 0$. As done with z in problem 2.a I can show that for any ϵ , $-\epsilon < 0$. Thus by (O5) $x - \epsilon > y + \epsilon$.

Now let $x - \epsilon < y + \epsilon$ for all $\epsilon > 0$. Assume that $x > y$. Then $x - y > 0$ so I can set $\epsilon_0 = \frac{x-y}{3}$. Then plugging in I get $x - \frac{x-y}{3} < y + \frac{x-y}{3}$. $\epsilon_0 > 0$ so by (O5)

$$x < y + \epsilon_0 + \epsilon_0$$

$$x + \epsilon_0 < y + \epsilon_0 + \epsilon_0 + \epsilon_0 \quad (\text{O5})$$

$$x + \epsilon_0 < y + (x + (-y))$$

$$x + \epsilon_0 < y + ((-y) + x) \quad (\text{A1})$$

$$x + \epsilon_0 < (y + (-y)) + x \quad (\text{A2})$$

$$x + \epsilon_0 < 0 + x \quad (\text{A4})$$

$$x + \epsilon_0 < x \quad (\text{A3})$$

$$(-x) + x + \epsilon_0 < (-x) + x \quad (\text{O4})$$

$$0 + \epsilon_0 < 0 \quad (\text{A4})$$

$$\epsilon_0 < 0 \quad (\text{A3})$$

This is a contradiction with (O1) because I know that $\epsilon_0 > 0$. Therefore $x \leq y$.

Part (b)

Let n be the largest natural number such that $\frac{1}{n} < y - x$. Let k be the largest natural number such that $\frac{k}{n} \leq x$. Then by our selection of k , $\frac{k+1}{n} > x$.

Now assume that $y \leq \frac{k+1}{n}$. Then I have that $\frac{k+1}{n} \geq y$ and $-\frac{k}{n} \geq -x$ so by (O5):

$$\frac{1}{n} = \frac{k+1}{n} - \frac{k}{n} \geq y - x$$

. This is a contradiction so it must be the case that $y > \frac{k+1}{n}$. Thus $z = \frac{k+1}{n}$ is a number satisfying $x < z < y$.

Part (c)

First I prove consequent 5 introduced in class that $-(x - y) = y - x$.

$$\begin{array}{ll}
 -(x - y) = -(x + (-y)) & \text{def. of -} \\
 -(x - y) = -1 \cdot (x + (-y)) & \text{see prob. 2} \\
 -(x - y) = -1 \cdot x + -1 \cdot (-y) & \text{(D)} \\
 -(x - y) = -x + -(-y) & \text{see prob. 2} \\
 -(x - y) = -x + y & \text{proved in class} \\
 -(x - y) = y + (-x) & \text{(A1)} \\
 -(x - y) = y - x & \text{def. of -}
 \end{array}$$

Let y be the smaller value of $b - x$ and $x - a$, both of which are positive. If $y = x - a$ then

$$\begin{array}{ll}
 x - y = x + (-(x - a)) & \\
 x - y = x + (a + (-x)) & \text{consequent 5} \\
 x - y = x + ((-x) + a) & \text{(A1)} \\
 x - y = (x + (-x)) + a & \text{(A2)} \\
 x - y = 0 + a & \text{(A4)} \\
 x - y = a & \text{(A3)}
 \end{array}$$

$a \leq a$ by definition so in this case $a \leq x - y$.

The other case is when $y = b - x$. By our selection of y I know $x - a > y$ so $(x - a) - y > 0$ and I also know from the first case that $x - (x - a) \geq a$. So by (O5),

$$\begin{array}{ll}
 x + (-(x - a)) + ((x - a) + (-y)) \geq a & \\
 x + ((-(x - a)) + (x - a)) + (-y) \geq a & \text{(A2)} \\
 x + 0 + (-y) \geq a & \text{(A4)} \\
 x + (-y) \geq a & \text{(A3)}
 \end{array}$$

so in both cases $a \leq x - y$. It is given that both $b - x$ and $x - a$ are positive so in either case $y > 0$. By (O4) $x + y > x$. Application of (O5) obtains $x > x - y$.

Now I want to show that $x + y \leq b$. In the case when $y = b - x$

$$\begin{array}{ll}
 x + y = x + (b + (-x)) & \\
 x + y = x + ((-x) + b) & \text{(A1)} \\
 x + y = (x + (-x)) + b & \text{(A2)} \\
 x + y = 0 + b & \text{(A4)} \\
 x + y = b & \text{(A3)}
 \end{array}$$

so $x + y \leq b$ by definition.

In the case when $y = x - a$ I know by our selection of y that $y < (b - x)$ and thus $0 > y + (-(b - x))$ by (O4). I also know from the first case that $b \geq x + (b - x)$. Then by

(O5)'

$$b + 0 > x + (b - x) + (y + (-(b - x))) \quad (A1)$$

$$b + 0 > x + (b - x) + ((-(b - x)) + y) \quad (A2)$$

$$b + 0 > x + ((b - x) + (-(b - x))) + y \quad (A3)$$

$$b + 0 > x + 0 + y \quad (A4)$$

$$b > x + y \quad (A5)$$

So therefore $x + y \leq b$ by the definition of \leq .

Having shown that $x < x + y$ I can use the result of part b) to produce some number z such that $x < z < x + y$. Let $\epsilon = z - x$. To show that ϵ satisfies the desired characteristics it must be shown that $\epsilon > 0$, and that $a < x - \epsilon < x + \epsilon < b$. It is known that $x < z$ so by (O4) $\epsilon = z - x > 0$. By the same process as before, $0 > -\epsilon$. Then by (O2) $-\epsilon < \epsilon$ and by (O4) $x - \epsilon < x + \epsilon$.

It is known that $z < x + y$. Then by (O4) $z - x < y$ so $\epsilon < y$. It is also known $a \leq x - y$. Then by (O5)' $a + \epsilon < x$ and by (O4) $a < x - \epsilon_0$.

From above it is known that $x + y \leq b$ and that $\epsilon < y$. Then by (O5)' $x + y + \epsilon < b + y$ and by (O4) $x + \epsilon < b$.

Thus ϵ satisfies the desired properties and by definition $(x - \epsilon, x + \epsilon) \subset (a, b)$.

Problem 6

Prove that each of the following are metric spaces.

(a) $X = \mathbb{R}, d(x, y) = |y - x|$

(b) $X = \text{any set}, d(x, y) = 1 \text{ if } x \neq y \text{ and } d(x, y) = 0 \text{ if } x = y.$

(c) Give another example of a metric space.

Solution

Part (a)

This proof will use the fact that $-1 \cdot x = -x$. This was proven as an intermediate step in problem 2.

First I will prove that $-(x - y) = y - x$.

$$\begin{aligned}
 -(x - y) &= -1 \cdot (x + (-y)) && \text{see problem 2} \\
 -(x - y) &= -1 \cdot x + -1 \cdot -y && \text{(D)} \\
 -(x - y) &= -x + -(-y) && \text{see problem 2} \\
 -(x - y) &= -x + y && \text{proved in class} \\
 -(x - y) &= y + (-x) && \text{(A1)} \\
 -(x - y) &= y - x && \text{def. of -}
 \end{aligned}$$

i $d(x, y) = 0 \iff x = y$

First assume $x = y$. Then $|y - x| = |0| = 0$. Now assume that $|y - x| = 0$. Then either $y - x = 0$ or $x - y = 0$. In the first case $y - x + x = x$ so by (A4) $y = x$. In the second case $x - y + y = y$ so by (A4) $x = y$.

ii $d(x, y) = d(y, x)$

This would directly follow from a proof of property 2 of absolute values that states $|y - x| = |x - y|$. There are two cases.

(i) Case: $y - x > 0$.

By the definition of absolute value $|y - x| = y - x$. Then

$$\begin{aligned}
 y - x &> 0 \\
 y - x + x &> 0 + x && \text{O4} \\
 y + 0 &> 0 + x && \text{A4} \\
 y &> x && \text{A3} \\
 y + (-y) &> x + (-y) && \text{O4} \\
 0 &> x + (-y) && \text{A4} \\
 0 &> x - y && \text{def. of -}
 \end{aligned}$$

Thus by the definition of absolute value $|x - y| = -(x - y)$ which, as proved at the beginning of this problem, is equal to $y - x$.

Case: $y - x < 0$.

By the definition of absolute value $|y - x| = -(y - x)$. Using the same fact as above, this equals $x - y$.

$$\begin{aligned}
 y - x &< 0 \\
 y - x + x &< 0 + x && \text{O4} \\
 y + 0 &< 0 + x && \text{A4} \\
 y &< x && \text{A3} \\
 y + (-y) &< x + (-y) && \text{O4} \\
 0 &< x + (-y) && \text{A4} \\
 0 &< x - y && \text{def. of -}
 \end{aligned}$$

Thus $|x - y| = x - y$ by definition.

Case: $y - x = 0$

In this case $|y - x| = y - x = 0$ by definition.

$$\begin{aligned} y - x &= 0 \\ y - x + x &= 0 + x \\ y + 0 &= 0 + x \end{aligned} \tag{A4}$$

$$y = x \tag{A3}$$

$$\begin{aligned} y + (-y) &= x + (-y) \\ 0 &= x + (-y) \end{aligned} \tag{A4}$$

$$0 = x - y \quad \text{def. of -}$$

So $|x - y| = y - x = 0$ by definition.

$$\text{iii } d(x, z) \leq d(x, y) + d(y, z)$$

$|z - x| \leq |y - x| + |z - y|$ by the triangle inequality proved in class.

Part (b)

$$\text{i } d(x, y) = 0 \iff x = y$$

This is true by the definition of the function d .

$$\text{ii } d(x, y) = d(y, x)$$

In the case when $x = y$, $d(x, y) = 0 = d(y, x)$. In the case when $x \neq y$, $d(x, y) = 1 = d(y, x)$.

$$\text{iii } d(x, z) \leq d(x, y) + d(y, z)$$

$$\begin{aligned} \text{(i) Case: } x = y = z \\ 0 &\leq 0 \end{aligned}$$

$$\begin{aligned} \text{(ii) Case: } x \neq y \neq z \\ 1 &\leq 2 \end{aligned}$$

$$\begin{aligned} \text{(iii) Case: } x = y \neq z \\ 1 &\leq 1 \end{aligned}$$

$$\begin{aligned} \text{(iv) Case: } x \neq y = z \\ 1 &\leq 1 \end{aligned}$$

$$\begin{aligned} \text{(v) Case: } x = z \neq y \\ 0 &\leq 1 \end{aligned}$$

Part (c)

$$X = \mathbb{C}, d(x, y) = \sqrt{x^2 + y^2}.$$