
This report is for evaluation purposes only. There are 23 days remaining in your free trial (Expires on 4/20/2018).

Linear Regression Report

Dataset Untitled
Y = CYCLOMATIC_COMPLEXITY X = MAINTAINABILITY

Linear Regression Plot Section -



This report is for evaluation purposes only.

There are 23 days remaining in your free trial (Expires on 4/20/2018).

Linear Regression Report

Dataset Untitled
Y = CYCLOMATIC COMPLEXITY X = MAINTAINABILITY

Run Summary Section -

Parameter	Value	Parameter	Value
Dependent Variable 10	CYCLOMATIC_COMPLEX	ITY	Rows Processed
Independent Variable	MAINTAINABILITY	Rows Used in Estimation	10
Frequency Variable	None	Rows with X Missing	0
Weight Variable	None	Rows with Freq Missing	0
Intercept	-16571.4079	Rows Prediction Only	0
Slope	5.6289	Sum of Frequencies	10
R-Squared	0.9919	Sum of Weights	10.0000
Correlation	0.9959	Coefficient of Variation	0.0169
Mean Square Error	2393365	Square Root of MSE	1547.05

Summary Statement -

The equation of the straight line relating CYCLOMATIC_COMPLEXITY and MAINTAINABILITY is estimated as: CYCLOMATIC_COMPLEXITY = (-16571.4079) + (5.6289) MAINTAINABILITY using the 10 observations in this dataset. The y-intercept, the estimated value of CYCLOMATIC_COMPLEXITY when MAINTAINABILITY is zero, is -16571.4079 with a standard error of 3495.9922. The slope, the estimated change in CYCLOMATIC_COMPLEXITY per unit change in MAINTAINABILITY, is 5.6289 with a standard error of 0.1799. The value of R-Squared, the proportion of the variation in CYCLOMATIC_COMPLEXITY that can be accounted for by variation in MAINTAINABILITY, is 0.9919. The correlation between CYCLOMATIC_COMPLEXITY and MAINTAINABILITY is 0.9959.

A significance test that the slope is zero resulted in a t-value of 31.2915. The significance level of this t-test is 0.0000. Since 0.0000 < 0.0500, the hypothesis that the slope is zero is rejected.

The estimated slope is 5.6289. The lower limit of the 95% confidence interval for the slope is 5.2141 and the upper limit is 6.0437. The estimated intercept is -16571.4079. The lower limit of the 95% confidence interval for the intercept is -24633.1804 and the upper limit is -8509.6354.

This report is for evaluation purposes only. There are 23 days remaining in your free trial (Expires on 4/20/2018).

Linear Regression Report

Dataset Untitled
Y = CYCLOMATIC_COMPLEXITY X = MAINTAINABILITY

Regression Estimation Section -

	Intercept	Slope
Parameter	B(0)	B(1)
Regression Coefficients	-16571.4079	5.6289
Lower 95% Confidence Limit	-24633.1804	5.2141
Upper 95% Confidence Limit	-8509.6354	6.0437
Standard Error	3495.9922	0.1799
Standardized Coefficient	0.0000	0.9959
T Value	-4.7401	31.2915
Prob Level (T Test)	0.0015	0.0000
Reject H0 (Alpha = 0.0500)	Yes	Yes
Power (Alpha = 0.0500)	0.9846	1.0000
Regression of Y on X	-16571.4079	5.6289
Inverse Regression from X on Y	-17456.3998	5.6749
Orthogonal Regression of Y and X	-17429.5291	5.6735

Notes:

The above report shows the least-squares estimates of the intercept and slope followed by the corresponding standard errors, confidence intervals, and hypothesis tests. Note that these results are based on several assumptions that should be validated before they are used.

Estimated Model

(-16571.4079437042) + (5.62886537429478) * (MAINTAINABILITY)

This report is for evaluation purposes only. There are 23 days remaining in your free trial (Expires on 4/20/2018).

Linear Regression Report

Dataset Untitled
Y = CYCLOMATIC_COMPLEXITY X = MAINTAINABILITY

Correlation and R-Squared Section —

	Pearson		Spearman Rank
	Correlation		Correlation
Parameter	Coefficient	R-Squared	Coefficient
Estimated Value	0.9959	0.9919	1.0000
Lower 95% Conf. Limit (r dist'n)	0.9803		
Upper 95% Conf. Limit (r dist'n)	0.9989		
Lower 95% Conf. Limit (Fisher's z)	0.9823		1.0000
Upper 95% Conf. Limit (Fisher's z)	0.9991		1.0000
Adjusted (Rbar)		0.9909	
T-Value for H0: Rho = 0	31.2915	31.2915	
Prob Level for H0: Rho = 0	0.0000	0.0000	0.0000

Notes:

The confidence interval for the Pearson correlation assumes that X and Y follow the bivariate normal distribution. This is a different assumption from linear regression which assumes that X is fixed and Y is normally distributed.

Two confidence intervals are given. The first is based on the exact distribution of Pearson's correlation. The second is based on Fisher's z transformation which approximates the exact distribution using the normal distribution. Why are both provided? Because most books only mention Fisher's approximate method, it will often be needed to do homework. However, the exact methods should be used whenever possible.

The confidence limits can be used to test hypotheses about the correlation. To test the hypothesis that rho is a specific value, say r0, check to see if r0 is between the confidence limits. If it is, the null hypothesis that rho = r0 is not rejected. If r0 is outside the limits, the null hypothesis is rejected.

Spearman's Rank correlation is calculated by replacing the orginal data with their ranks. This correlation is used when some of the assumptions may be invalid.

Summary Matrices ————

	X'X	X'X	Χ'Y	X'X Inverse	X'X Inverse
Index	0	1	2	0	1
0	10	192434	917471	5.106603	-0.0002601725
1	192434	3.777049E+09	1.80716E+10	-0.0002601725	1.352009E-08
2 (Y'Y)			8.653794E+10		
Determina	ant	7.396401E+08			1.352009E-09

Variance - Covariance Matrix of Regression Coefficients ————

	VC(b)	VC(b)
Ind	lex 0	1
0	1.222196E+07	-622.6875
1	-622.6875	0.0323585

This report is for evaluation purposes only. There are 23 days remaining in your free trial (Expires on 4/20/2018).

Linear Regression Report

Dataset Untitled

Y = CYCLOMATIC_COMPLEXITY X = MAINTAINABILITY

Tests of Assumptions Section -

	Test	Prob	Is the Assumption Reasonable at the 0.2000
Assumption/Test	Value	Level	Level of Significance?
Residuals follow Normal Distribu	ution?		
Shapiro Wilk	0.9612	0.799688	Yes
Anderson Darling	0.2110	0.858692	Yes
D'Agostino Skewness	-0.7419	0.458132	Yes
D'Agostino Kurtosis	0.0812	0.935301	Yes
D'Agostino Omnibus	0.5570	0.756901	Yes
Constant Residual Variance?			
Modified Levene Test	0.8907	0.372915	Yes
Relationship is a Straight Line?			
Lack of Linear Fit F(0, 0) Test	0.0000	0.000000	No

No Serial Correlation?

Evaluate the Serial-Correlation report and the Durbin-Watson test if you have equal-spaced, time series data.

Notes

A 'Yes' means there is not enough evidence to make this assumption seem unreasonable. This lack of evidence may be because the sample size is too small, the assumptions of the test itself are not met, or the assumption is valid.

A 'No' means the that the assumption is not reasonable. However, since these tests are related to sample size, you should assess the role of sample size in the tests by also evaluating the appropriate plots and graphs. A large dataset (say N > 500) will often fail at least one of the normality tests because it is hard to find a large dataset that is perfectly normal.

Normality and Constant Residual Variance:

Possible remedies for the failure of these assumptions include using a transformation of Y such as the log or square root, correcting data-recording errors found by looking into outliers, adding additional independent variables, using robust regression, or using bootstrap methods.

Straight-Line:

Possible remedies for the failure of this assumption include using nonlinear regression or polynomial regression.


This report is for evaluation purposes only.

There are 23 days remaining in your free trial (Expires on 4/20/2018).

Linear Regression Report

Dataset Untitled
Y = CYCLOMATIC_COMPLEXITY X = MAINTAINABILITY


Residual Plots Section -




This report is for evaluation purposes only. There are 23 days remaining in your free trial (Expires on 4/20/2018).


Linear Regression Report

Dataset Untitled
Y = CYCLOMATIC_COMPLEXITY X = MAINTAINABILITY





7



This report is for evaluation purposes only. There are 23 days remaining in your free trial (Expires on 4/20/2018).

Linear Regression Report

Dataset Untitled

Y = CYCLOMATIC_COMPLEXITY X = MAINTAINABILITY

Procedure Input Settings —

Autosaved Template File

C:\Users\KASATLA\Documents\NCSS 12\Procedure Templates\Autosave\Linear Regression and Correlation -

Autosaved 2018_3_30-0_56_24.t153 Variables Tab -- Variables ---Y: Dependent Variable(s): CYCLOMATIC COMPLEXITY X: Independent Variable: MAINTAINABILITY Frequency Variable: <Empty> Weight Variable: <Empty> -- Model Specification -----Remove Intercept Unchecked -- Resampling (Increases computation time) ------Calculate Bootstrap C.I.'s Unchecked Run Randomization Tests Unchecked -- Alpha Levels -----Alpha for C.I.'s and Tests: 0.050 Alpha for Assumptions: 0.20 **Reports Tab** -- Select Report / Plot Group ------Select a Group of Reports and Plots: Display only those items that are CHECKED BELOW **Show Notes** Checked Show All Rows Checked -- Select Reports ------Run Summary Checked Summary Statement Checked Descriptive Statistics Unchecked Correlation and R-Squared Checked **Summary Matrices** Checked ·· Estimation ······ Regression Estimation Checked ·· ANOVA ······ ANOVA Unchecked ·· Assumptions ····· Assumptions Checked Levene Groups: 2

Durbin-Watson Unchecked **PRESS** Unchecked

This report is for evaluation purposes only. There are 23 days remaining in your free trial (Expires on 4/20/2018).

Linear Regression Report

Dataset Untitled

Y = CYCLOMATIC_COMPLEXITY X = MAINTAINABILITY

Procedure Input Settings (Continued)

Reports Tab (Continued)	
·· Prediction ······	
Predict Y at these X values:	<empty></empty>
Predicted Y - C.L.	Unchecked
Predicted Y - P.L.	Unchecked
·· Row-by-Row Lists ·····	
Original Ďata	Unchecked
Predicted Y Means	Unchecked
Predicted Y Individuals	Unchecked
Simultaneous Bands	Unchecked
Predicted X Means	Unchecked
Predicted X Individuals	Unchecked
1 Todioted / Marvidadio	Choncoked
·· Regression Diagnostics ·····	
Residuals	Unchecked
Residual Diagnostics	Unchecked
Leave One Row Out	Unchecked
Outlier Detection Chart	Unchecked
Influence Detection Chart	Unchecked
Outlier-Influence Chart	Unchecked
Outlier-Illinderice Chart	Officialization
Report Options Tab	
Report Options	
Precision:	Single
Variable Names:	Names
variable (varies).	Tumos
·· Decimal Places ······	
Probability:	4
Beta (Coefficients):	4
SE:	4
T:	4
R2:	4
X:	4
Λ. Υ:	4
Residuals:	4
Std Residuals:	4
	All
Sum Squares:	
Matrix:	All

This report is for evaluation purposes only. There are 23 days remaining in your free trial (Expires on 4/20/2018).

Linear Regression Report

Dataset Untitled

Y = CYCLOMATIC_COMPLEXITY X = MAINTAINABILITY

Procedure Input Settings (Continued)

Plots Tab Select Plots	
Y vs X	Checked
RStudent vs X	Checked
Histogram	Checked
Residuals vs X	Checked
Residuals vs Row	Checked
Probability Plot	Checked
Residuals vs X	Checked
Serial Correlation	Checked
Plot Options	
Y vs X Plot Size:	Medium
All Other Plot Sizes:	Small
Resampling Tab Bootstrap Calculation Options Sampling ······ Samples (N): Sampling Method: Retries:	3000 Observations 50
·· Estimation ······	
Percentile Type:	Ave X(p[n+1])
C.I. Method:	Reflection
Bootstrap Confidence Coefficients:	0.90 0.95 0.99
Randomization Test Options	
Monte Carlo Samples:	1000
Storage Tab Data Storage Options	
Storage Option:	Do not store data