


This report is for evaluation purposes only.
There are 23 days remaining in your free trial (Expires on 4/20/2018).

Linear Regression Report

Dataset Untitled
Y = Coupling X = no_of_Issues

Linear Regression Plot Section



This report is for evaluation purposes only.
There are 23 days remaining in your free trial (Expires on 4/20/2018).

Linear Regression Report

Dataset Untitled
 Y = Coupling X = no_of_Issues

Run Summary Section

Parameter	Value	Parameter	Value
Dependent Variable	Coupling	Rows Processed	10
Independent Variable	no_of_Issues	Rows Used in Estimation	10
Frequency Variable	None	Rows with X Missing	0
Weight Variable	None	Rows with Freq Missing	0
Intercept	416.9674	Rows Prediction Only	0
Slope	0.8178	Sum of Frequencies	10
R-Squared	0.9886	Sum of Weights	10.0000
Correlation	0.9943	Coefficient of Variation	0.0357
Mean Square Error	17673.15	Square Root of MSE	132.9404

Summary Statement

The equation of the straight line relating Coupling and no_of_Issues is estimated as: Coupling = (416.9674) + (0.8178) no_of_Issues using the 10 observations in this dataset. The y-intercept, the estimated value of Coupling when no_of_Issues is zero, is 416.9674 with a standard error of 132.6559. The slope, the estimated change in Coupling per unit change in no_of_Issues, is 0.8178 with a standard error of 0.0311. The value of R-Squared, the proportion of the variation in Coupling that can be accounted for by variation in no_of_Issues, is 0.9886. The correlation between Coupling and no_of_Issues is 0.9943.

A significance test that the slope is zero resulted in a t-value of 26.2881. The significance level of this t-test is 0.0000. Since $0.0000 < 0.0500$, the hypothesis that the slope is zero is rejected.

The estimated slope is 0.8178. The lower limit of the 95% confidence interval for the slope is 0.7461 and the upper limit is 0.8896. The estimated intercept is 416.9674. The lower limit of the 95% confidence interval for the intercept is 111.0624 and the upper limit is 722.8725.

This report is for evaluation purposes only.
There are 23 days remaining in your free trial (Expires on 4/20/2018).

Linear Regression Report

Dataset Untitled
 Y = Coupling X = no_of_Issues

Regression Estimation Section

Parameter	Intercept B(0)	Slope B(1)
Regression Coefficients	416.9674	0.8178
Lower 95% Confidence Limit	111.0624	0.7461
Upper 95% Confidence Limit	722.8725	0.8896
Standard Error	132.6559	0.0311
Standardized Coefficient	0.0000	0.9943
T Value	3.1432	26.2881
Prob Level (T Test)	0.0137	0.0000
Reject H0 (Alpha = 0.0500)	Yes	Yes
Power (Alpha = 0.0500)	0.7858	1.0000
Regression of Y on X	416.9674	0.8178
Inverse Regression from X on Y	378.6784	0.8273
Orthogonal Regression of Y and X	401.5788	0.8217

Notes:

The above report shows the least-squares estimates of the intercept and slope followed by the corresponding standard errors, confidence intervals, and hypothesis tests. Note that these results are based on several assumptions that should be validated before they are used.

Estimated Model

$(416.967438165739) + (0.817845942790729) * (\text{no_of_Issues})$

This report is for evaluation purposes only.
There are 23 days remaining in your free trial (Expires on 4/20/2018).

Linear Regression Report

Dataset Untitled
 Y = Coupling X = no_of_Issues

Correlation and R-Squared Section

Parameter	Pearson Correlation Coefficient	R-Squared	Spearman Rank Correlation Coefficient
Estimated Value	0.9943	0.9886	1.0000
Lower 95% Conf. Limit (r dist'n)	0.9722		
Upper 95% Conf. Limit (r dist'n)	0.9984		
Lower 95% Conf. Limit (Fisher's z)	0.9750		1.0000
Upper 95% Conf. Limit (Fisher's z)	0.9987		1.0000
Adjusted (Rbar)		0.9871	
T-Value for H0: Rho = 0	26.2881	26.2881	
Prob Level for H0: Rho = 0	0.0000	0.0000	0.0000

Notes:

The confidence interval for the Pearson correlation assumes that X and Y follow the bivariate normal distribution. This is a different assumption from linear regression which assumes that X is fixed and Y is normally distributed.

Two confidence intervals are given. The first is based on the exact distribution of Pearson's correlation. The second is based on Fisher's z transformation which approximates the exact distribution using the normal distribution. Why are both provided? Because most books only mention Fisher's approximate method, it will often be needed to do homework. However, the exact methods should be used whenever possible.

The confidence limits can be used to test hypotheses about the correlation. To test the hypothesis that rho is a specific value, say r_0 , check to see if r_0 is between the confidence limits. If it is, the null hypothesis that $\rho = r_0$ is not rejected. If r_0 is outside the limits, the null hypothesis is rejected.

Spearman's Rank correlation is calculated by replacing the original data with their ranks. This correlation is used when some of the assumptions may be invalid.

Summary Matrices

Index	X'X 0	X'X 1	X'Y 2	X'X Inverse 0	X'X Inverse 1
0	10	40442	37245	0.9957243	-0.0002214837
1	40442	1.818151E+08	1.655598E+08	-0.0002214837	5.476576E-08
2 (Y'Y)			1.510737E+08		
Determinant		1.825959E+08			5.476576E-09

Variance - Covariance Matrix of Regression Coefficients

Index	VC(b) 0	VC(b) 1
0	17597.59	-3.914315
1	-3.914315	0.0009678835

This report is for evaluation purposes only.
There are 23 days remaining in your free trial (Expires on 4/20/2018).

Linear Regression Report

Dataset Untitled
 Y = Coupling X = no_of_Issues

Tests of Assumptions Section

Assumption/Test	Test Value	Prob Level	Is the Assumption Reasonable at the 0.2000 Level of Significance?
Residuals follow Normal Distribution?			
Shapiro Wilk	0.8730	0.108287	No
Anderson Darling	0.6197	0.106750	No
D'Agostino Skewness	-1.4773	0.139593	No
D'Agostino Kurtosis	0.0486	0.961201	Yes
D'Agostino Omnibus	2.1848	0.335410	Yes
Constant Residual Variance?			
Modified Levene Test	0.0022	0.963422	Yes
Relationship is a Straight Line?			
Lack of Linear Fit F(0, 0) Test	0.0000	0.000000	No

No Serial Correlation?

Evaluate the Serial-Correlation report and the Durbin-Watson test if you have equal-spaced, time series data.

Notes:

A 'Yes' means there is not enough evidence to make this assumption seem unreasonable. This lack of evidence may be because the sample size is too small, the assumptions of the test itself are not met, or the assumption is valid.

A 'No' means the that the assumption is not reasonable. However, since these tests are related to sample size, you should assess the role of sample size in the tests by also evaluating the appropriate plots and graphs. A large dataset (say $N > 500$) will often fail at least one of the normality tests because it is hard to find a large dataset that is perfectly normal.

Normality and Constant Residual Variance:

Possible remedies for the failure of these assumptions include using a transformation of Y such as the log or square root, correcting data-recording errors found by looking into outliers, adding additional independent variables, using robust regression, or using bootstrap methods.

Straight-Line:





Possible remedies for the failure of this assumption include using nonlinear regression or polynomial regression.

This report is for evaluation purposes only.
There are 23 days remaining in your free trial (Expires on 4/20/2018).

Linear Regression Report

Dataset Untitled
Y = Coupling X = no_of_Issues




Residual Plots Section



This report is for evaluation purposes only.
There are 23 days remaining in your free trial (Expires on 4/20/2018).

Linear Regression Report

Dataset Untitled
Y = Coupling X = no_of_Issues



This report is for evaluation purposes only.
There are 23 days remaining in your free trial (Expires on 4/20/2018).

Linear Regression Report

Dataset Untitled
 Y = Coupling X = no_of_Issues

Procedure Input Settings

Autosaved Template File

C:\Users\KASATLA\Documents\NCSS 12\Procedure Templates\Autosave\Linear Regression and Correlation - Autosaved 2018_3_29-22_59_39.t153

Variables Tab

```
-- Variables -----
Y: Dependent Variable(s):      Coupling
X: Independent Variable:       no_of_Issues
Frequency Variable:           <Empty>
Weight Variable:              <Empty>

-- Model Specification -----
Remove Intercept               Unchecked

-- Resampling (Increases computation time) -----
Calculate Bootstrap C.I.'s     Unchecked
Run Randomization Tests       Unchecked

-- Alpha Levels -----
Alpha for C.I.'s and Tests:    0.050
Alpha for Assumptions:        0.20
```

Reports Tab

```
-- Select Report / Plot Group -----
Select a Group of Reports and Plots:  Display only those items that are CHECKED BELOW
Show Notes                           Checked
Show All Rows                         Checked

-- Select Reports -----
.. Summaries .....
Run Summary                          Checked
Summary Statement                    Checked
Descriptive Statistics                Unchecked
Correlation and R-Squared             Checked
Summary Matrices                     Checked

.. Estimation .....
Regression Estimation                 Checked

.. ANOVA .....
ANOVA                               Unchecked

.. Assumptions .....
Assumptions                          Checked
  Levene Groups:                      2
Durbin-Watson                        Unchecked
PRESS                               Unchecked
```


This report is for evaluation purposes only.
There are 23 days remaining in your free trial (Expires on 4/20/2018).

Linear Regression Report

Dataset Untitled
 Y = Coupling X = no_of_Issues

Procedure Input Settings (Continued)

Reports Tab (Continued)

.. Prediction	
Predict Y at these X values:	<Empty>
Predicted Y - C.L.	Unchecked
Predicted Y - P.L.	Unchecked
.. Row-by-Row Lists	
Original Data	Unchecked
Predicted Y Means	Unchecked
Predicted Y Individuals	Unchecked
Simultaneous Bands	Unchecked
Predicted X Means	Unchecked
Predicted X Individuals	Unchecked
.. Regression Diagnostics	
Residuals	Unchecked
Residual Diagnostics	Unchecked
Leave One Row Out	Unchecked
Outlier Detection Chart	Unchecked
Influence Detection Chart	Unchecked
Outlier-Influence Chart	Unchecked

Report Options Tab

-- Report Options -----	
Precision:	Single
Variable Names:	Names
.. Decimal Places	
Probability:	4
Beta (Coefficients):	4
SE:	4
T:	4
R2:	4
X:	4
Y:	4
Residuals:	4
Std Residuals:	4
Sum Squares:	All
Matrix:	All

This report is for evaluation purposes only.
There are 23 days remaining in your free trial (Expires on 4/20/2018).

Linear Regression Report

Dataset Untitled
 Y = Coupling X = no_of_Issues

Procedure Input Settings (Continued)

Plots Tab

-- Select Plots -----	
Y vs X	Checked
RStudent vs X	Checked
Histogram	Checked
Residuals vs X	Checked
Residuals vs Row	Checked
Probability Plot	Checked
Residuals vs X	Checked
Serial Correlation	Checked
-- Plot Options -----	
Y vs X Plot Size:	Medium
All Other Plot Sizes:	Small

Resampling Tab

-- Bootstrap Calculation Options -----	
.. Sampling	
Samples (N):	3000
Sampling Method:	Observations
Retries:	50
.. Estimation	
Percentile Type:	Ave X(p[n+1])
C.I. Method:	Reflection
Bootstrap Confidence Coefficients:	0.90 0.95 0.99
-- Randomization Test Options -----	
Monte Carlo Samples:	1000

Storage Tab

-- Data Storage Options -----	
Storage Option:	Do not store data