
This report is for evaluation purposes only. There are 23 days remaining in your free trial (Expires on 4/20/2018).

Linear Regression Report

Dataset Untitled
Y = CYCLOMATIC_COMPLEXITY X = MAINTAINABILITY

Linear Regression Plot Section -



This report is for evaluation purposes only.

There are 23 days remaining in your free trial (Expires on 4/20/2018).

Linear Regression Report

Dataset Untitled
Y = CYCLOMATIC_COMPLEXITY X = MAINTAINABILITY

Run Summary Section —

Parameter	Value	Parameter	Value
Dependent Variable	CYCLOMATIC_COMPLEX	ITY	Rows Processed
10			
Independent Variable	MAINTAINABILITY	Rows Used in Estimation	10
Frequency Variable	None	Rows with X Missing	0
Weight Variable	None	Rows with Freq Missing	0
Intercept	3810.8578	Rows Prediction Only	0
Slope	5.7986	Sum of Frequencies	10
R-Squared	0.9949	Sum of Weights	10.0000
Correlation	0.9974	Coefficient of Variation	0.0234
Mean Square Error	361053.9	Square Root of MSE	600.8776

Summary Statement -

The equation of the straight line relating CYCLOMATIC_COMPLEXITY and MAINTAINABILITY is estimated as: CYCLOMATIC_COMPLEXITY = (3810.8578) + (5.7986) MAINTAINABILITY using the 10 observations in this dataset. The y-intercept, the estimated value of CYCLOMATIC_COMPLEXITY when MAINTAINABILITY is zero, is 3810.8578 with a standard error of 585.6470. The slope, the estimated change in CYCLOMATIC_COMPLEXITY per unit change in MAINTAINABILITY, is 5.7986 with a standard error of 0.1472. The value of R-Squared, the proportion of the variation in CYCLOMATIC_COMPLEXITY that can be accounted for by variation in MAINTAINABILITY, is 0.9949. The correlation between CYCLOMATIC_COMPLEXITY and MAINTAINABILITY is 0.9974.

A significance test that the slope is zero resulted in a t-value of 39.3975. The significance level of this t-test is 0.0000. Since 0.0000 < 0.0500, the hypothesis that the slope is zero is rejected.

The estimated slope is 5.7986. The lower limit of the 95% confidence interval for the slope is 5.4592 and the upper limit is 6.1380. The estimated intercept is 3810.8578. The lower limit of the 95% confidence interval for the intercept is 2460.3533 and the upper limit is 5161.3622.

This report is for evaluation purposes only. There are 23 days remaining in your free trial (Expires on 4/20/2018).

Linear Regression Report

Dataset Untitled
Y = CYCLOMATIC_COMPLEXITY X = MAINTAINABILITY

Regression Estimation Section —

	Intercept	Slope
Parameter	B(0)	B(1)
Regression Coefficients	3810.8578	5.7986
Lower 95% Confidence Limit	2460.3533	5.4592
Upper 95% Confidence Limit	5161.3622	6.1380
Standard Error	585.6470	0.1472
Standardized Coefficient	0.0000	0.9974
T Value	6.5071	39.3975
Prob Level (T Test)	0.0002	0.0000
Reject H0 (Alpha = 0.0500)	Yes	Yes
Power (Alpha = 0.0500)	0.9999	1.0000
Regression of Y on X	3810.8578	5.7986
Inverse Regression from X on Y	3698.3705	5.8285
Orthogonal Regression of Y and X	3701.6036	5.8276

Notes:

The above report shows the least-squares estimates of the intercept and slope followed by the corresponding standard errors, confidence intervals, and hypothesis tests. Note that these results are based on several assumptions that should be validated before they are used.

Estimated Model

(3810.85775801676) + (5.79861901322685) * (MAINTAINABILITY)

This report is for evaluation purposes only. There are 23 days remaining in your free trial (Expires on 4/20/2018).

Linear Regression Report

Dataset Untitled
Y = CYCLOMATIC_COMPLEXITY X = MAINTAINABILITY

Correlation and R-Squared Section -

			Spearman
	Pearson		Rank
	Correlation		Correlation
Parameter	Coefficient	R-Squared	Coefficient
Estimated Value	0.9974	0.9949	1.0000
Lower 95% Conf. Limit (r dist'n)	0.9875		
Upper 95% Conf. Limit (r dist'n)	0.9993		
Lower 95% Conf. Limit (Fisher's z)	0.9888		1.0000
Upper 95% Conf. Limit (Fisher's z)	0.9994		1.0000
Adjusted (Rbar)		0.9942	
T-Value for H0: Rho = 0	39.3975	39.3975	
Prob Level for H0: Rho = 0	0.0000	0.0000	0.0000

Notes:

The confidence interval for the Pearson correlation assumes that X and Y follow the bivariate normal distribution. This is a different assumption from linear regression which assumes that X is fixed and Y is normally distributed.

Two confidence intervals are given. The first is based on the exact distribution of Pearson's correlation. The second is based on Fisher's z transformation which approximates the exact distribution using the normal distribution. Why are both provided? Because most books only mention Fisher's approximate method, it will often be needed to do homework. However, the exact methods should be used whenever possible.

The confidence limits can be used to test hypotheses about the correlation. To test the hypothesis that rho is a specific value, say r0, check to see if r0 is between the confidence limits. If it is, the null hypothesis that rho = r0 is not rejected. If r0 is outside the limits, the null hypothesis is rejected.

Spearman's Rank correlation is calculated by replacing the orginal data with their ranks. This correlation is used when some of the assumptions may be invalid.

Summary Matrices —

	X'X	X'X	Χ'Y	X'X Inverse	X'X Inverse
Index	0	1	2	0	1
0	10	37638	256357	0.949948	-0.0002258218
1	37638	1.58329E+08	1.061523E+09	-0.0002258218	5.999835E-08
2 (Y'Y)			7.135195E+09		
Determinan	nt	1.666713E+08			5.999835E-09

Variance - Covariance Matrix of Regression Coefficients ————

	VC(b)	VC(b)
Index	0	1
0	342982.4	-81.53384
1	-81.53384	0.02166264

This report is for evaluation purposes only. There are 23 days remaining in your free trial (Expires on 4/20/2018).

Linear Regression Report

Dataset Untitled

Y = CYCLOMATIC_COMPLEXITY X = MAINTAINABILITY

Tests of Assumptions Section -

	Test	Prob	Is the Assumption Reasonable at the 0.2000
Assumption/Test	Value	Level	Level of Significance?
Residuals follow Normal Distribu	ition?		
Shapiro Wilk	0.8911	0.174540	No
Anderson Darling	0.5107	0.196424	No
D'Agostino Skewness	-1.2308	0.218393	Yes
D'Agostino Kurtosis	-0.3071	0.758731	Yes
D'Agostino Omnibus	1.6092	0.447258	Yes
Constant Residual Variance?			
Modified Levene Test	0.0043	0.949283	Yes
Relationship is a Straight Line?			
Lack of Linear Fit F(0, 0) Test	0.0000	0.000000	No

No Serial Correlation?

Evaluate the Serial-Correlation report and the Durbin-Watson test if you have equal-spaced, time series data.

Notes:

A 'Yes' means there is not enough evidence to make this assumption seem unreasonable. This lack of evidence may be because the sample size is too small, the assumptions of the test itself are not met, or the assumption is valid.

A 'No' means the that the assumption is not reasonable. However, since these tests are related to sample size, you should assess the role of sample size in the tests by also evaluating the appropriate plots and graphs. A large dataset (say N > 500) will often fail at least one of the normality tests because it is hard to find a large dataset that is perfectly normal.

Normality and Constant Residual Variance:

Possible remedies for the failure of these assumptions include using a transformation of Y such as the log or square root, correcting data-recording errors found by looking into outliers, adding additional independent variables, using robust regression, or using bootstrap methods.


Straight-Line:

Possible remedies for the failure of this assumption include using nonlinear regression or polynomial regression.

Linear Regression Report


Dataset Untitled
Y = CYCLOMATIC_COMPLEXITY X = MAINTAINABILITY


Residual Plots Section -




Linear Regression Report

Dataset Untitled
Y = CYCLOMATIC_COMPLEXITY X = MAINTAINABILITY





7



Linear Regression Report

Dataset Untitled

Y = CYCLOMATIC_COMPLEXITY X = MAINTAINABILITY

Procedure Input Settings —

Autosaved Template File

Levene Groups:

Durbin-Watson

PRESS

C:\Users\KASATLA\Documents\NCSS 12\Procedure Templates\Autosave\Linear Regression and Correlation - Autosaved 2018 3 30-1 3 22.t153

Autosaved 2018_3_30-1_3_22.t153 Variables Tab --- Variables ---Y: Dependent Variable(s): CYCLOMATIC COMPLEXITY X: Independent Variable: MAINTAINABILITY Frequency Variable: <Empty> Weight Variable: <Empty> -- Model Specification -----Remove Intercept Unchecked -- Resampling (Increases computation time) ------Calculate Bootstrap C.I.'s Unchecked Run Randomization Tests Unchecked -- Alpha Levels -----Alpha for C.I.'s and Tests: 0.050 Alpha for Assumptions: 0.20 **Reports Tab** -- Select Report / Plot Group ------Select a Group of Reports and Plots: Display only those items that are CHECKED BELOW **Show Notes** Checked Show All Rows Checked -- Select Reports ------Run Summary Checked Summary Statement Checked Descriptive Statistics Unchecked Correlation and R-Squared Checked **Summary Matrices** Checked ·· Estimation ······ Regression Estimation Checked ·· ANOVA ······ ANOVA Unchecked ·· Assumptions ····· Assumptions Checked

2

Unchecked

Unchecked

Linear Regression Report

Dataset Untitled

Y = CYCLOMATIC_COMPLEXITY X = MAINTAINABILITY

Procedure Input Settings (Continued)

Reports Tab (Continued)	
·· Prediction ······	
Predict Y at these X values:	<empty></empty>
Predicted Y - C.L.	Unchecked
Predicted Y - P.L.	Unchecked
·· Row-by-Row Lists ·····	
Original Ďata	Unchecked
Predicted Y Means	Unchecked
Predicted Y Individuals	Unchecked
Simultaneous Bands	Unchecked
Predicted X Means	Unchecked
Predicted X Individuals	Unchecked
1 Todioted / Marvidadio	Choncoked
·· Regression Diagnostics ·····	
Residuals	Unchecked
Residual Diagnostics	Unchecked
Leave One Row Out	Unchecked
Outlier Detection Chart	Unchecked
Influence Detection Chart	Unchecked
Outlier-Influence Chart	Unchecked
Outlier-Illinderice Chart	Officialization
Report Options Tab	
Report Options	
Precision:	Single
Variable Names:	Names
variable (varies).	Tumos
·· Decimal Places ·····	
Probability:	4
Beta (Coefficients):	4
SE:	4
T:	4
R2:	4
X:	4
Λ. Υ:	4
Residuals:	4
Std Residuals:	4
	All
Sum Squares:	
Matrix:	All

Linear Regression Report

Dataset Untitled

Y = CYCLOMATIC_COMPLEXITY X = MAINTAINABILITY

Procedure Input Settings (Continued)

Plots Tab Select Plots	
Y vs X	Checked
RStudent vs X	Checked
Histogram	Checked
Residuals vs X	Checked
Residuals vs Row	Checked
Probability Plot	Checked
Residuals vs X	Checked
Serial Correlation	Checked
	• · · · · · · · · · · · · · · · · · · ·
Plot Options	
Y vs X Plot Size:	Medium
All Other Plot Sizes:	Small
Resampling Tab Bootstrap Calculation Options Sampling Samples (N): Sampling Method: Retries:	3000 Observations 50
Percentile Type:	Ave X(p[n+1])
C.I. Method:	Reflection
Bootstrap Confidence Coefficients:	0.90 0.95 0.99
·	
Monte Carlo Samples:	1000
Storage Tab Data Storage Options Storage Option:	 Do not store data
Otorage Option.	Do not store data